

# Detection of rtxA Gene as a Biomarker of Seafood-Borne Pathogen *Vibrio cholerae* using In Silico PCR Assay

*by* Stalis Norma Ethica

---

**Submission date:** 11-Feb-2021 10:48AM (UTC+0700)

**Submission ID:** 1506832499

**File name:** 2020\_Detection\_of\_rtx\_Gene.pdf (657.26K)

**Word count:** 3845

**Character count:** 21114



## Detection of *rtxA* Gene as a Biomarker of Seafood-Borne Pathogen *Vibrio cholerae* using *In Silico* PCR Assay

Stalis Norma Ethica<sup>1\*</sup>, Nur Hidayati<sup>1,2</sup>, Hayatun Fuad<sup>1</sup>, Chaerul Arham<sup>1,3</sup>, Rivana Ariyadi<sup>1,4</sup>, Ellyka Purwaningrum<sup>1,5</sup>, and Kazi M. Anmad Zillur Rahman<sup>6</sup>

<sup>1</sup>Magister Program of Medical Laboratory Science, Universitas Muhammadiyah Semarang, Jl. Kedungmundu Raya No.18, Semarang, Indonesia 50273

<sup>2</sup>Anugerah Pekalongan Hospital, Jl. Perintis Kemerdekaan no. 3, Pekalongan, Indonesia, 51145

<sup>3</sup>Politeknik Kesehatan Muhammadiyah Makassar, Jl. Ratulangi no. 101, Makassar, Indonesia, 90222

<sup>4</sup>Sekolah Tinggi Ilmu Kesehatan Muhammadiyah Ciamis, Jl. K.H Ahmad Dahlan no. 20, Ciamis, Indonesia, 46216

<sup>5</sup>Medical Faculty, Universitas Wahid Hasyim Semarang, Jl. Kolonel Warsito, Semarang, Indonesia, 50232

<sup>6</sup>Centre for Marine Science and Innovation, University of New South Wales, Level 5 East, Biological Sciences North (D26), Kensington, New South Wales, Australia, 2052

Article history:

Received: 4 March 2020; Revised: 17 July 2020; Accepted: 1 August 2020

### Abstract

Seafood-borne outbreaks caused by *Vibrio cholerae* have led to the increased need for food safety risk assessment of marine products. An *in silico* investigation about the potential of virulence gene of *V. cholerae*, *rtxA*, as a DNA biomarker of the toxigenic bacterium has been carried out. The aim of this study was to use the bacterial DNA biomarker sequence as a tool to facilitate early rapid detection of cholera infection. Five specific pairs of primers were designed from the *rtxA* open reading frame DNA of *V. cholerae* O1 biovar El Tor str. N16961 genomic DNA using Primer3Plus. Next, *in silico* Polymerase Chain Reaction (PCR) assay was carried out using the newly designed primers and 25 genomic DNA of vibrio spp. retrieved from the *in silico* database. One of the five designed pairs of primers, RtxAOF-RtxAOR: '5-CGCAAAACAGTTTCAGCCGA-3' and 5'-AGGTTGGTCTTTTGTGGCCA-3', could result in single DNA amplicon sized 518 bp only from *V. cholerae* species. No amplicon bands were produced from 17 other vibrio genomes studied using similar RtxAF-RtxAR primers. A further check showed that the amplicon was indeed part of the *rtxA* gene of *V. cholerae*. Based on this *in silico* study, *rtxA* gene appeared to be a DNA biomarker of *V. cholerae*, which is potential to facilitate rapid diagnosis of the virulence bacterium using *in silico* PCR assay.

**Keywords:** seafood-borne infection, *Vibrio cholerae*, DNA biomarker, primer design, *rtxA*

### 1. Introduction

Seafood products are consumed worldwide, therefore playing a significant role in the economic market (Bonnin-Jusserand et al., 2019). The proportion of globally produced-seafood commodities that is internationally-traded is higher than other products. It is also increasing mostly due to globalization (Guillen et al., 2019). However, several recent outbreaks of human gastroenteritis have been linked to the consumption of contaminated seafood (Elbashir et al., 2018).

Infectious diseases are classified as illnesses caused by pathogenic microorganisms, including bacteria. Such diseases have been the main threat

worldwide and have a significant impact on public health and the world's economy (Hwang, Hwang, & Bueno, 2018). Seafood-borne outbreaks caused by bacterium have led to the increased need for food safety risk assessment of marine products. Continuous monitoring of bacterial contamination in aquatic products and identification of risk factors are, therefore, crucial for assuring food safety (Xu, Wu, & Chen, 2019).

Investigation of seafood-borne illnesses caused by bacteria and viruses requires a concrete knowledge about the pathogenicity and virulence properties of the etiologic agents (Elbashir et al., 2018; Iwamoto, Ayers, Mahon, & Swerdlow, 2010). Among pathogenic bacteria- contaminating seafood, *V. cholerae* is the

\*Corresponding author.  
E-mail: norma@unimus.ac.id

main *Vibrio* species responsible for the dramatic increase of seafood-borne infections worldwide (Bonnin-Jusserand et al., 2019). *V. cholerae* is notorious as the causal agent of cholera, a severe diarrheal disease, which could be quickly fatal if untreated. It is commonly transmitted via contaminated water and person-to-person contact (Baker-Austin et al., 2018). Approximately 2.9 million cases of cholera and 95,000 deaths had occurred annually worldwide between 2008 and 2012 (Ali, Nelson, Lopez, & Sack, 2015).

To date, more than 200 serogroups of *V. cholerae* have been recognized, based on variable somatic O antigen composition; O1 was the only known epidemic serogroup of *V. cholerae* up to 1991. In 1992, serogroup O139 was identified as the second epidemic serogroup of *V. cholerae* (Albert et al., 1993; Bhattacharya et al., 1993). Within the *V. cholerae* O1 serogroup, there are two well-established biotypes: classical and El Tor. The El Tor biotype emerged in 1961 and subsequently displaced the classical biotype as a cause of cholera throughout the world (Pradhan, Baidya, Ghosh, Paul, & Chowdhury, 2010). Both classical and El Tor biotypes are differentiated from each other by several properties, including agglutination of chicken red blood cells, susceptibility to polymyxin B and to biotype-specific phages, haemolysis of sheep red blood cells, and Voges-Proskauer reaction (Kaper, Morris Jr, & Levine, 1995). *V. cholerae* strains of El Tor biotype initiated the seventh cholera pandemic, which is continuing to cause outbreaks worldwide with catastrophic effects.

Among the critical virulence factors of *V. cholerae* are the cholera toxin (CTX) genetic element and the repeats in toxin (RTX) cluster (Davis, Moyer, Boyd, & Waldor, 2000). CTX and RTX are the chief virulence gene clusters and are grouped together (Cheng, Zhou, Kan, Wang, & Rui, 2014). The RTX family of toxins generally produced by several pathogenic Gram-negative bacteria. The RTX toxins constitute a family of dominant virulence factors that have widely spread among Gram-negative bacteria. In *V. cholerae*, the RTX toxin gene cluster encodes the presumptive cytotoxin (*rtxA*), an acyltransferase (*rtxC*), and a related ATP-binding cassette transporter system (two proteins for toxin transportation, RtxB and RtxD) (Chou et al., 2001; Lin et al., 1999)

The search for the ideal biomarkers in infectious diseases with high sensitivity, specificity, and predictive capacity must initially be focused on detection and identification of the infectious agent (Mohan & Harikrishna, 2015). *V. cholerae* genomic sequence provides a starting point for understanding how a free-living, environmental organism emerged to become a significant human bacterial pathogen. The complete

genomic sequence of the Gram-negative, gamma-Proteobacterium *V. cholerae* El Tor N16961 has been reported to have a length of 4,033,460 base pairs (bp) (Heidelberg et al., 2000). Numerous mutations, however, have been reported in *V. cholerae* O1 strains encompassing genes, which code virulence factors including the repeat in toxins (*rtxA*). In the case of seafood-borne infectious disease, detection and identification of the infectious agent *V. cholerae* are therefore necessary to do. In this study, *rtxA*, a member of RTX chief virulence gene cluster, was tested for its potential as a DNA biomarker of *V. cholerae*. Such biomarker is beneficial to facilitate early rapid diagnosis of *V. cholerae* infection using *in vitro* PCR.

## 2. Material and Methods

A literature study was initially conducted to obtain information related to the unique phenotypic feature of *V. cholerae*, particularly on specific toxins the species produces. Next, the genotype feature, such as genes associated with proteins underlining the unique phenotype was targeted. Primers were then designed from the internal part of open reading frame (ORF) of the targeted gene sequences in FASTA format as input using Primer3Plus at <https://primer3plus.com/> (Untergasser et al., 2007; Ethica et al., 2013; Ethica, 2014; Ethica, Sulistyningtyas, & Darmawati, 2019; Ethica, Darmawati, Dewi, Nurrahman, & Sulistyningtyas, 2020). The input of the program is basically a FASTA format of a gene of interest. Pairs of primers having the least possibility of hairpin formation, self-complementarity, and dimerization were selected (Ethica, Semiarti, Widada, Oedjijono, & Raharjo, 2017). The newly designed primers were subsequently used as input for the web-based *in silico* PCR. The virtual PCR work was run from a web-based software at <http://insilico.ehu.es/PCR/> using *Vibrio* spp. genomes stored in the program database. *In silico* PCR can be done by choosing "PCR Amplification" from the main page of the website, and once it opens, inserts the primers, then click "amp30" (Bikandi, Millán, Rementeria, & Garaizar, 2004; San Millán, Martínez-Ballesteros, Rementeria, Garaizar, & Bikandi, 2013); Ethica, Sulistyningtyas, & Darmawati, 2019; Ethica, Darmawati, Dewi, Nurrahman, & Sulistyningtyas, 2020). The nucleotide database used in the *in silico* PCR website comes from National Center of Bioinformatics Institute (NCBI) database. An analysis was eventually carried out to confirm if the *in silico* PCR products (amplicons) were specific to *V. cholerae* species and were indeed part of genes targeted as DNA biomarkers. This analysis is conducted to check whether the *in silico* PCR amplicons virtually seen on gel electrophoresis were correct according to primer positions. The analysis

was carried out by “clicking” the amplicon band or size shown by the *in silico* PCR result window.

### 3. Results and Discussion

Severe infection outbreaks originated from fish and other foods often create a diagnostic challenge for clinicians. There is a limited official guidance to help clinicians decide which biomarkers that can aid with the diagnosis of bacterial infections (Rogers et al., 2011). Therefore, the development of bacterial DNA biomarkers to detect *V. cholerae* was carried out in this study. It was expected that in the case of diarrheal infection outbreaks, the involvement of *V. cholerae* could be determined by detecting the presence of their DNA biomarkers.

From a literature study, it is known that *rtxA* gene, a member of RTX gene cluster, is among distinct genotypic features of infectious bacterium *V. cholerae* (Lin et al., 1999). The uniqueness of the gene makes it potential to be targeted as a DNA biomarker of *V. cholerae* species. The *rtxA* gene sequen<sup>[22]</sup> was then retrieved from the genome sequence of *V. cholerae*

O1 biovar El Tor strain N16961 (NCBI Reference<sup>[21]</sup> sequence: NC\_002505.1). The sequence encodes RTX toxin RtxA [*Vibrio cholerae* O1 biovar El Tor str. N16961 in the NCBI GenBank] (NCBI Reference Sequence: NP\_231094.1). The 3-D structure of cholera RtxA protein (UniProt accession number: Q9KS12) as the product of *rtxA* gene is shown in Figure 1. Using primer3Plus (Untergasser et al., 2007), pairs of primers obtained by using *rtxA* DNA sequence as input (Figure 1) are listed in Table 1.

*In silico* PCR assay using Pair 2 primers and genomic DNA of 25 *Vibrio* species were run on <http://insilico.ehu.es/PCR/>. The assay resulted in eight single 518-bp DNA bands (Figure 2) belonging to all *V. cholerae* or species number 4 to 11 (Table 2, highlighted). According to Table 2, *V. cholerae* species number 4-11 are *V. cholerae*, *V. cholerae* str. IEC224, *V. cholerae* str. LMA3984-4, *V. cholerae* str. M66-2, *V. cholerae* str. MJ-1236, *V. cholerae* O1 str. 2010EL-1786, *V. cholerae* str. O395, and *V. cholerae* str. O395 chromosome 1. Table 1 shows that though primers were designed from similar *rtxA* sequence, not all primers could detect unique conserved regions of *rtxA*




Figure 1. The 3-D structure of cholera RtxA protein (*rtxA* gene product) visualized by Litemol (UniProt Accession Number: Q9KS12)

Table 1. Primers designed using Primer3Plus from *rtxA* DNA sequence

Primer	Forward Primer	Reverse Primer	Amplicon Size (bp)	DNA of <i>V. cholerae</i> strains amplified (genome code number)
Pair 1	GCGCGAACGTAATAACCCAC	ACCGACATGCCATCACCAAT	508	4,5, and 7-9
Pair 2	CGCAAAACAGTTTCAGCCGA	AGGTTGGTCTTTTGTGGCCA	518	4-11 (all <i>V. cholerae</i> )
Pair 3	GTGGCGCGAACGTAATAACC	AGATGTTGACGTTCCCACC	597	4,5, and 7-9
Pair 4	TAGCGGTGAAAGCTCAGGTG	GTTATTACGTTCCGCGCCACC	585	4,5, and 7-9
Pair 5	GCGCATTTACTGGCTTACGG	TGGTGAAGATGTTACCCGCC	559	4-9

of all strains of *V. cholerae*, which differentiate them from other *Vibrio* species (genome code number 1-3 and 12-25, see Table 2). Also, not all of these primers could detect all *V. cholerae* strains. Only Pair 2 primer that could indicate *rtxA* conserved regions of all *V. cholerae* strains. As seen in Table 1, Primer3Plus by default produces five best pairs of primers a output and automatically eliminates low quality primers that have the potential of mispriming, self-dimer, or cross dimer.

After carrying out the *in silico* PCR assay, it is necessary to check whether the amplicon products that were seen on the virtual gel electrophoresis

(Figure 2) were correct, based on the primer positions. The analysis was carried out easily by clicking the amplicon size that appeared on the *in silico* PCR result window. A new tab was then appeared, showing that the *in silico* amplicons were indeed part of the *rtxA* gene fragments (Figure 3). The DNA sequence seen in Figure 3 represents one of eight PCR amplicon run on *in silico* PCR using the selected Pair 2 primer (see Table 1) and a genome of vibrios (species number 4, see Table 2).

“Clicking” step on the amplicon size on *in silico* PCR (Figure 2) was necessary. The result is shown in Figure 3. This is a way to prove that even though

Table 2. List of genomic DNA of *Vibrio* spp. used as *in silico* PCR templates

**No. Complete list of *Vibrio* strains in the database used as *in silico* PCR templates**

- 1 *Vibrio alginolyticus* NBRC 15630 = ATCC 17749
- 2 *Vibrio anguillarum* 775
- 3 *Vibrio campbellii* ATCC BAA-1116
- 4 *Vibrio cholerae*
- 5 *Vibrio cholerae* IEC224
- 6 *Vibrio cholerae* LMA3984-4
- 7 *Vibrio cholerae* M66-2
- 8 *Vibrio cholerae* MJ-1236
- 9 *Vibrio cholerae* O1 str. 2010EL-1786
- 10 *Vibrio cholerae* O395
- 11 *Vibrio cholerae* O395 chromosome 1
- 12 *Vibrio fischeri* ES 114
- 13 *Vibrio fischeri* MJ11
- 14 *Vibrio furnissii* NCTC 11218
- 15 *Vibrio harveyi* ATCC BAA-1116
- 16 *Vibrio parahaemolyticus* BB22OP
- 17 *Vibrio parahaemolyticus* O1:K33 str. CDC\_K4557
- 18 *Vibrio parahaemolyticus* O1:Kuk str. FDA\_R31
- 19 *Vibrio parahaemolyticus* RIMD 2210633
- 20 *Vibrio* sp. EJY3
- 21 *Vibrio* sp. Ex25
- 22 *Vibrio splendidus* LGP32
- 23 *Vibrio vulnificus* CMCP6
- 24 *Vibrio vulnificus* MO6-24/O
- 25 *Vibrio vulnificus* YJ016

Note: Highlighted rows are genomic sequences of *Vibrio cholerae* strains among those of other *Vibrios* (25 strains) available that were retrieved from the *in silico* database at the time of this study.




Figure 2. *In silico* PCR assay using Pair 2 primer and 25 *Vibrio* genomic DNA resulted seven single 518-bp DNA bands belonging to all *V. cholerae* strains

Genome: ***Vibrio cholerae***  
 Start position: 1558124  
 End position: 1558641  
 Length: 518

**DNA sequence**

```
>NC_002505, from 1558124 to 1558641 (518 bp); Vibrio cholerae
CGCAAACAGTTTCAGCCGACTATATGACCAGCTAGAAGCCGCTAACTTGAAAGAGAGTAAGCACCTTTA
TCTGGATCAAATGGTGACTTTGTTACCAAAGGCAAAGGTAATCTTGCCAATATCGATCTGCTAGGTAGC
CGCGAAGCTGTGCTTGAAAAAGTGAAGTTAACAGTAAGTAACGAGTACGGTCAAACCGTTGCGGATACAA
TTTTTGCTGGATATCAGCCAAAGATCTTGCCAAAGACGGTAAAGGGTTGATATCGCGGGTTGAATAA
AGTACATCAAGCGATTGAACAGCATCTGTCACCTGTGACGCCACGTTGTACATTTGGAACCGAGTGTAT
CATAGCGCGCTAGGTCATGCCGATTGCAAATAGGCCAAGGTCGCACGCAACTTGAAGTCAAGCTGCAG
CTGATTTTAACCGCAAATACGTAAGCTGGTGGCCACTAGGCAGCAAGTCATCAATATCAGCAATAT
CTTGAATGTGGCCACAAAAGACCAACCT
```

- [Translate to protein](#)
- [Restriction digest](#)
- [BLAST](#)
- [Design primers with primer3](#)
- [Genome related info at NCBI](#)

**Gene(s) or part of gene(s) amplified:**

ORF. 1550108-1563784 [Sequence](#) **VC1451 - RTX toxin RtxA**

Figure 3. *In silico* PCR amplicon with *V. cholerae* genome sequence as the template using Pair 2 primer designed in this study

*Vibrio* spp. might have the same *rtxA* gene, our primers recognize specific DNA regions, which is part of *rtxA* of *V. cholerae* only, and not part of *rtxA* of other *Vibrio* species. Primer sequences used in this study were taken from the *rtxA* gene of *V. cholerae* precisely because we targeted all strains of this species. Not all primers designed from the *rtxA* gene could amplify

the targeted genome selectively and therefore the *in silico* PCR was conducted. In this case, checking the amplicon size was carried out to confirm if the amplicon was the genuine part of *rtxA*. Occasionally, *in silico* PCR resulted DNA amplicon, which is not the targeted gene. This step also confirmed that primers, other than Pair 2, that were designed from




Figure 4. The output of Primer3Plus showing complete parameters of Pair 2 Primer, including HP (hairpin possibility)

the targeted *rtxA* gene of a *V. cholerae* could produce positive *in silico* PCR bands using all of its strains' genome (in this study eight strains).

Primers designed to amplify the *rtxA* gene of *V. cholerae* have been reported before. According to the previous report, a pair of designed primers, *rtxA*-F 5'-GGGATACAATGCCCTCTGGCA-3' and *rtxA*-R 5'-TGGGTTGGCGTTGGATTTTAC-3', was successfully used to detect *rtxA* of *V. cholerae* (Xu et al., 2019; Rivera, Chun, Huq, Sack, & Colwell, 2001). Despite being able to amplify the targeted genes in most *V. cholerae* strains by showing 977 bp amplicon, the primers were failed to amplify that of *V. cholerae* strain LMA 3984-4. If a pair of primers theoretically failed to amplify a targeted DNA sequence in an *in silico* PCR, then they were most likely unable to amplify the same sequence in an *in vitro* PCR.

In this study, the potential of *rtxA* gene as a DNA biomarker for *V. cholerae* was tested by conducting *in silico* PCR assay. To prove the hypothesis, five pairs of primers were successfully designed using web-based Primer3Plus software using *rtxA* full-length gene sequence of *V. cholerae* O1 biovar El Tor str. N16961. Among five pairs of primers obtained, a new pair of primers (Pair 2 based on Table 1) namely RtxAOF-RtxAOR: '5-CGCAAAACAGTTTCAGCCGA-3' and 5'-AGGTTGGTCTTTGTGGCCA-3' passed the hairpin and other stability test results based on Primer3Plus output as seen in Figure 4. In addition, the selected

newly designed specific primer, the RtxAOF-RtxAOR, was able to selectively amplify the internal part of the *rtxA* gene fragments of all genomic sequences of *V. cholerae*, but not those of other Vibrios. [27] important to also underline that the *in silico* PCR assay developed in this study for *rtxA* could identify *rtxA*-producing *V. cholerae* strains, not only from the O1, but also the non-O1 serogroup.

Our results infer that the *rtxA* gene fragment of *V. cholerae* flanked by a pair of primers designed in this study that indicated sequences between QNSFSR and ATKDQPD conserved regions of protein sequences coded by the gene appeared as the biomarker of the bacterium. The *rtxA* gene fragment could differentiate the infectious bacterial species from other Vibrios.

The newly designed RtxAOF-RtxAOR (Pair 2 primer resulted from primer design work of Primer3Plus) is potential to be used as prospective primers to detect the presence of *V. cholerae* in bacterial samples using *in vitro* PCR.

Among limitation of this work is that the performance of the obtained primers is only on a theoretical basis, which means the results are predictive. Visual PCR simulation in this study also only utilized resources accessed from <http://insilico.ehu.es/PCR>. If an update on the genomic sequence in this study is released, results might slightly change due to the alterations on DNA

templates used for *in silico* PCR. Therefore, if a new genomic DNA sequence of *V. cholerae* strain is added on the database, then consequently new *in silico* PCR assay should be conducted. The aim is to check if the selectivity level of these primers stays the same. Moreover, results from this study could be more meaningful if they are applied for an *in vivo* PCR study. Thus, they could support early detection and identification to control *V. cholerae* as cholera infection agent directly from actual samples.

#### 4. Conclusion

Based on *in silico* PCR assay carried out in this study, the *rtxA* gene was theoretically proved to be a DNA biomarker of the seafood-borne pathogenic bacterium *V. cholerae*. The aforementioned gene could be used to differentiate it from other *Vibrio* species. The newly designed primers are potential to facilitate early rapid diagnosis of cholera infection.

#### References

- Albert, M. J., Ansaruzzaman, M., Bardhan, P. K., Faruque, G., Faruque, S. M., Islam, M. S., ... & Yunus, M. D. (1993). Large epidemic of cholerae-like disease in Bangladesh caused by *Vibrio cholerae* O139 synonym Bengal. *Lancet*, *342*(8868), 387-90. doi: 10.1016/0140-6736(93)92811-7.
- Ali, M., Nelson, A. R., Lopez, A. L., & Sack, D. A. (2015). Updated global burden of cholera in endemic countries. *PLoS Neglected Tropical Diseases*, *9*(6), e0003832. doi: 10.1371/journal.pntd.0003832.
- Baker-Austin, C., Oliver, J. D., Alam, M. A., Waldor, M. K., Qadri, F., & Martinez-Urtaza, J. (2018). *Vibrio* spp. infections. *Nature Reviews Disease Primers*, *4*(1), 1-19. doi: 10.1038/s41572-018-0005-8.
- Bhattacharya, M. K., Bhattacharya, S. K., Garg, S., Ha, P., Dutta, D., Nair, G. B., ... & Sack, R. B. (1993). Outbreak of *Vibrio cholerae* non-01 in India and Bangladesh. *Lancet*, *341*(8856), 1346-1347. doi: 10.1016/0140-6736(93)90855-B.
- Bikandi, J., Millán, R. S., Rementería, A., & Garaizar, J. (2004). In silico analysis of complete bacterial genomes: PCR, AFLP-PCR and nucleotide restriction. *Bioinformatics*, *20*(5), 798-799. doi: 10.1093/bioinformatics/btg491.
- Bonnin-Jusserand, M., Copin, S., Le Bris, C., Brauge, T., Gay, M., Brisabois, A., ... & Midelet-Bourdin, G. (2019). *Vibrio* species involved in seafood-borne outbreaks (*Vibrio cholerae*, *V. parahaemolyticus* and *V. vulnificus*): Review of microbiological versus recent molecular detection methods in seafood products. *Critical Reviews in Food Science and Nutrition*, *59*(4), 597-610. doi: 10.1080/10408398.2017.1384715.
- Chen, C., Zhou, Y., Kan, B., Wang, Q., & Rui, Y. (2014). Construction and characterization of a *Vibrio cholerae* serogroup O139 vaccine candidate by genetic engineering. *Molecular Medicine Reports*, *9*(6), 2239-2244. doi: 10.3892/mmr.2014.2065.
- Chow, K. H., Ng, T. K., Yuen, K. Y., & Yam, W. C. (2001). Detection of RTX toxin gene in *Vibrio cholerae* by PCR. *Journal of Clinical Microbiology*, *39*(7), 2594-2597. doi: 10.1128/JCM.39.7.2594-2597.2001.
- Davis, B. M., Royer, K. E., Boyd, E. F., & Waldor, M. K. (2000). CTX prophages in classical biotype *Vibrio cholerae*: functional phage genes but dysfunctional phage genomes. *Journal of Bacteriology*, *182*(24), 6992-6998. doi: 10.1128/JB.182.24.6992-6998.2000.
- Elbashir, S., Parveen, S., Schwarz, J., Rippen, T., Jahncke, M., & DePaola, A. (2018). Seafood pathogens and information on antimicrobial resistance: A review. *Food Microbiology*, *70*, 85-93. doi: 10.1016/j.fm.2017.09.011.
- Ethica, S. N. (2014). *Detection of genes involved in glycerol metabolism of Alcaligenes* sp. JG3 (Doctoral dissertation, Universitas Gadjah Mada). Retrieved from <https://repository.ugm.ac.id/id/eprint/126377>.
- Ethica, S. N., Hammi, M. K., Lestari, P., Semiarti, E., Widada, J., & Raharjo, T. J. (2013). Amplification of *Azospirillum* sp. JG3 *glpD* gene fragment using degenerate primers generated by web-based tools. *Journal of Microbiology, Biotechnology and Food Sciences*, *3*(3), 231.
- Ethica, S. N., Semiarti, E., Widada, J., Oedjijono, O., & Raharjo, T. J. (2017). Characterization of *moaC* and a nontarget gene fragments of food borne pathogen *Alcaligenes* sp. JG3 using degenerate colony and arbitrary PCRs. *Journal of Food Safety*, *37*(4), e12345. doi: 10.1111/jfs.12345.
- Ethica, S. N., Sulistyningtyas, A. R., & Darmawati, S. (2019). *In silico* specificity comparison between GMF-GMR and JMF-JMR primers for detecting *moaC* genes of food spoilage bacteria *Pseudomonas* spp. In *IOP Conference Series: Earth and Environmental Science*, *292*(1), 012033. IOP Publishing. doi: 10.1088/1755-1315/292/1/012033.
- Ethica, S. N., Darmawati, S., Dewi, S. S., Nurrahman, N., & Sulistyningtyas, A. R. (2020). Streptolysin encoding genes *sagC* and *sagD* as biomarkers of fish pathogen *Streptococcus iniae*: an *in silico* study. *Squalen Bulletin of Marine and Fisheries Postharvest and Biotechnology*, *15*(1), 31-39. doi: 10.15578/squalen.v15i1.416.
- Guillen, J., Natale, F., Carvalho, N., Casey, J., Hofherr, J., Druon, J. N., ... & Martinsohn, J. T. (2019). Global seafood consumption footprint. *Ambio*, *48*(2), 111-122. doi:10.1007/s13280-018-1060-9.
- Heidelberg, J. F., Eisen, J. A., Nelson, W. C., Clayton, R. G., Gwinn, M. L., Dodson, R. J., ... & Gill, S. R. (2000). DNA sequence of both chromosomes of the cholerae pathogen *Vibrio cholerae*. *Nature*, *406*(6795), 477-483. doi: 10.1038/35020000.
- Hwang, H., Hwang, B. Y., & Bueno, J. (2018). Biomarkers in infectious diseases. *Disease Markers*, 8509127. doi: 10.1155/2018/8509127.



# Detection of rtxA Gene as a Biomarker of Seafood-Borne Pathogen *Vibrio cholerae* using In Silico PCR Assay

## ORIGINALITY REPORT

11%

SIMILARITY INDEX

7%

INTERNET SOURCES

8%

PUBLICATIONS

4%

STUDENT PAPERS

## PRIMARY SOURCES

- |   |  |     |
|---|--|-----|
| 1 | <a href="http://publications.jrc.ec.europa.eu">publications.jrc.ec.europa.eu</a><br>Internet Source  | 1%  |
| 2 | Racault, Abdulaziz, George, Menon, C, Punathil, McConville, Loveday, Platt, Sathyendranath, Vijayan. "Environmental Reservoirs of <i>Vibrio cholerae</i> : Challenges and Opportunities for Ocean-Color Remote Sensing", Remote Sensing, 2019<br>Publication | 1%  |
| 3 | B. Bakhshi, M. Boustanshenas, A. Mahmoudi-aznaveh. " Emergence of O1 classical biotype in 2012 in Iran ", Letters in Applied Microbiology, 2014<br>Publication   | 1%  |
| 4 | Submitted to Avondale College of Higher Education<br>Student Paper   | <1% |
| 5 | Xiao Cao, Lichao Zhao, Jingfeng Zhang, Xun Chen, Lei Shi, Xiang Fang, Hui Xie, Yanlei  | <1% |

Chang, Li Wang. "Detection of viable but nonculturable *Vibrio parahaemolyticus* in shrimp samples using improved real-time PCR and real-time LAMP methods", *Food Control*, 2019

Publication

---

6

C. L. Cordero, S. Sozhamannan, K. J. F. Satchell. "RTX Toxin Actin Cross-Linking Activity in Clinical and Environmental Isolates of *Vibrio cholerae*", *Journal of Clinical Microbiology*, 2007

Publication

---

7

[sfx.library.newschool.edu](http://sfx.library.newschool.edu)

Internet Source

---

8

Sedef Tunca. "Transcriptional regulation of the desferrioxamine gene cluster of *Streptomyces coelicolor* is mediated by binding of DmdR1 to an iron box in the promoter of the *desA* gene : Regulation of the desferrioxamine gene cluster", *FEBS Journal*, 01/25/2007

Publication

---

9

[medalsf.blogspot.com](http://medalsf.blogspot.com)

Internet Source

---

10

Submitted to University of Ulster

Student Paper

---

11

[astutipage.wordpress.com](http://astutipage.wordpress.com)

Internet Source

---

<1%

<1%

<1%

<1%

<1%

<1%

12

Sara B. Hernández, Tobias Dörr, Matthew K. Waldor, Felipe Cava. "Modulation of peptidoglycan synthesis by recycled cell wall tetrapeptides", Cold Spring Harbor Laboratory, 2019

Publication

<1%

13

Submitted to University of Westminster

Student Paper

<1%

14

Sangita Pal, B Guhathakurta, D Sasmal, R Mallick, A Datta. "Purification and characterisation of a hemolysin with phospholipase C activity from *Vibrio cholerae* O139", FEMS Microbiology Letters, 2006

Publication

<1%

15

[scholars.unh.edu](https://scholars.unh.edu)

Internet Source

<1%

16

[ejournal.forda-mof.org](http://ejournal.forda-mof.org)

Internet Source

<1%

17

Submitted to University of Sydney

Student Paper

<1%

18

J. Yu, B. Zhu, T. Zhou, Y. Wei, X. Li, Y. Liu. "Species-specific Identification of sp. based on 16S-23S rRNA gene internal transcribed spacer", Journal of Applied Microbiology, 2020

Publication

<1%

19	<a href="http://www.coursehero.com">www.coursehero.com</a> Internet Source	<1%
20	Submitted to Armstrong Atlantic State University Student Paper	<1%
21	<a href="http://en.dogeno.us">en.dogeno.us</a> Internet Source	<1%
22	Hiroshi Asakura. "Gene expression profile of <i>Vibrio cholerae</i> in the cold stress-induced viable but non-culturable state", <i>Environmental Microbiology</i> , 4/2007 Publication	<1%
23	<a href="http://assets0.pubget.com">assets0.pubget.com</a> Internet Source	<1%
24	K. Bahrani, James D. Oliver. " Studies on the lipopolysaccharide of a virulent and an avirulent strain of ", <i>Biochemistry and Cell Biology</i> , 1990 Publication	<1%
25	<a href="http://publikasiilmiah.unwahas.ac.id">publikasiilmiah.unwahas.ac.id</a> Internet Source	<1%
26	<a href="http://www.cdc.gov">www.cdc.gov</a> Internet Source	<1%
27	Geevaretnam Jeyasekaran, Kannan Thirumalai Raj, Robinson Jeya Shakila, Albin Jemila Thangarani, Durairaj Sukumar. "Multiplex polymerase chain reaction-based assay for the	<1%

specific detection of toxin-producing *Vibrio cholerae* in fish and fishery products", *Applied Microbiology and Biotechnology*, 2011

Publication

28

[ejournal.upi.edu](http://ejournal.upi.edu)

Internet Source

<1%

29

[ijp.iranpath.org](http://ijp.iranpath.org)

Internet Source

<1%

30

Richard M. Mariita, Mohammad J. Hossain, Anthony G. Moss. " Comparative genomic analysis and characterization of sp. AOAB, isolated from a notoriously invasive gut revealed multiple antibiotic resistance determinants ", Cold Spring Harbor Laboratory, 2021

Publication

<1%

31

M S Islam. "Role of cyanobacteria in the persistence of *Vibrio cholerae* O139 in saline microcosms", *Canadian Journal of Microbiology*, 02/2004

Publication

<1%

32

[combrex.bu.edu](http://combrex.bu.edu)

Internet Source

<1%

33

J. Bikandi. "In silico analysis of complete bacterial genomes: PCR, AFLP-PCR and endonuclease restriction", *Bioinformatics*, 01/22/2004

Publication

<1%

---

Exclude quotes Off

Exclude bibliography Off

Exclude matches Off