

# Streptolysin Encoding Genes sagC and sagD as Biomarkers of Fish Pathogen *Streptococcus* *iniae*: An In Silico Study

*by* Ayu Rahmawati Sulistyaningsih

---

**Submission date:** 15-Oct-2020 10:57AM (UTC+0700)

**Submission ID:** 1415687414

**File name:** rnal-Streptolysin\_Encoding\_Genes\_sagC\_and\_sagD\_as\_Biomarkers.pdf (563.78K)

**Word count:** 4539

**Character count:** 27917



## Streptolysin Encoding Genes *sagC* and *sagD* as Biomarkers of Fish Pathogen *Streptococcus iniae*: An *In Silico* Study

Stalis Norma Ethica<sup>1</sup>, Sri Darmawati<sup>1</sup>, Sri Sinto Dewi<sup>2</sup>, Nur<sup>24</sup>nan<sup>2</sup>, and Ayu Rahmawati Sulistyanyingtyas<sup>2</sup>

<sup>1</sup>Magister Program of Medical Laboratory Science, Universitas Muhammadiyah Semarang, Jl. Kedungmundu Raya No. 18, Semarang, 50273, Indonesia

<sup>2</sup>Faculty of Nursing and Health Sciences, Universitas Muhammadiyah Semarang, Jl. Kedungmundu Raya No. 18, Semarang, 50273, Indonesia

39

Article history:

Received: 15 December 2019; Revised: 3 April 2020; Accepted: 7 April 2020

### Abstract

*Streptococcus iniae* has been notorious as a serious tilapia fish pathogen leading to many disease outbreaks in warm water marine aquaculture. An *in silico* investigation about the potential of virulence genes of *S. iniae*, *sagC* and *sagD*, as biomarkers of the bacterial species, has been carried out. The aim was to determine bacterial biomarkers, which are important to facilitate early rapid diagnosis of *S. iniae* streptococcal infection in fish and also in humans. First, specific primers were designed from *sagC* and *sagD* genes of *S. iniae* SF1 genomic DNA using Primer3Plus. Next, *in silico* PCR (Polymerase Chain Reaction) analysis was carried out using the newly designed primers and 117 genomic DNA of streptococci (all species) retrieved from the database. Primers designed from *sagC* and *sagD* genes (SagCF: '5- TGCTGACCTCCTAAAAGGGC -3' and SagCR: '5- CTATGCGGCGGGTTAAGGT -3' as well as SagDF: '5- GCCAATCCAATCCTGTATGC -3' and SagDR: '5- TGCAGCTCCATAACCCCTC -3') could result in a single band of each matching to 558-bp and 590-bp PCR products only from *S. iniae*. From 116 other streptococcal genomes studied using similar primers have resulted in no amplicon bands. A further check showed that the amplicons were truly part of *sagC* and *sagD* genes of *S. iniae*. These results showed that *sagC* and *sagD* genes appeared to be biomarkers of *S. iniae*, which are potential to be used to facilitate rapid diagnostic of the pathogenic bacterium.

**Keywords:** *Streptococcus iniae*, *sagC*, *sagD*, *in silico* PCR

### 1. Introduction

*Streptococcus iniae* has been notorious as the most severe aquatic pathogen in the last decade. Disease outbreaks caused by the bacterium had most likely been occurring for several decades in marine aquaculture leading to high losses in farmed marine in warmer regions. *Streptococcus iniae* is known as the endemic pathogen responsible for the development of streptococcosis in fish. Yet, the bacterial species is also known as a rising pathogen in human for causing infection on soft tissues (Agnew & Barnes, 2007; Fuller et al., 2002; Nho et al., 2009; Rajoo et al., 2015). *Streptococcus iniae* infection in human generally occurs through injuries associated with preparing whole fresh fish for cooking (Baiano & Barnes, 2009).

<sup>37</sup> *S. iniae* had been isolated from culture of red hybrid (Nile Tilapia *Oreochromis niloticus* x Mozambique Tilapia *O. mossambicus*) in Malaysia, blue tilapia (*Oreochromis aureus*) in Mexico and olive flounders (*Paralichthys olivaceus*) in South Korea. The isolated Gram-positive *S. iniae* appeared punctiform, transparently white, catalase and oxidase negative and produced complete  $\beta$ -hemolysis on blood agar. The isolate was sensitive to tetracycline, vancomycin, and bacitracin but was resistant to streptomycin, ampicillin, penicillin, and erythromycin (Ortega et al., 2018; Rahmatullah et al., 2017; Rajoo et al., 2015).

The current understanding of virulence mechanisms of *Streptococcus iniae* in fish is lacking. Although, it is known that in the case of fish infection by *S. iniae*, the virulence factor is B-hemolysin S (Locke et al., 2007). Hemolysins, or cytolytins, are well-recognized

\*Corresponding author.  
E-mail: ayurs@unimus.ac.id

features of many bacterial species, including streptococci, and are generally associated with damage to the membranes of a variety of mammalian cell types. Examples of streptococcal cytolytins include streptolysin O (SLO) and streptolysin S (SLS) group A streptococci (GAS) as well as the b-hemolysin/cytolysin of group B streptococci (GBS) implicated as a virulence factor in animal models of infection (Le Breton & McIver, 2013).

The pathogen group A Streptococcus (GAS) produces a wide spectrum of infections including necrotizing fasciitis (NF). Streptolysin S (SLS) produces the hallmark b-haemolytic phenotype produced by GAS. The nine-gene GAS locus (*sagA-sagI*) resembling a bacteriocin biosynthetic operon is vital and sufficient for SLS production. Datta et al. (2005) reported that for production of SLS, all of *sagA*, *sagB*, *sagC*, and *sagD*, are needed. Being members of *S. iniae* SLS-associated gene cluster responsible for streptolysin production, the *sagC* and *sagD* encoded predicted products with significant amino acid identity to corresponding *sag* products of GAS, yet shared no homology with other proteins in the GenBank database (Fuller et al., 2002). Both genes encode two streptolysin associated proteins, SagC and SagD, respectively.

Identification of biomarkers is critical for the diagnosis of fish bacterial infection early during invasion. However, the identification of reliable biomarkers is often hampered by a low concentration of microbes. Shanks et al. (2007) stated that the development of specific PCR assay targeting bacterial DNA as markers is important for the identification and diagnosis of bacterial fish infection with low concentration of bacterium. Various PCR methods designed for *S. iniae* detection had been studied (Zhou et al., 2011). However, none of them targeting *sagC* and *sagD* sequences of *S. iniae*, while it had been revealed the connection between the *sagC* and *sagD* genes with the invasive fish-infection agent (Fuller et al., 2002).

The term *in silico* referred to “computational” experiment conducted virtually using computers complementing both *in vitro* and *in vivo* terms, which refer to works done in living organisms. In genomics area of study, the intention of *in silico* PCR is achieving predicted PCR results (theoretical) using updated sequences of bacterial genome allowing amplification of targeted DNA sequences (Kalendar et al., 2017; Moretti, 2011). In addition, *in silico* PCR is useful for computing ability prediction of primers to amplify selected gene fragments by using up-to-date bacterial genomic DNA sequences retrieved from a database.

It allows amplification of specific DNA sequences supporting successful DNA amplification using *in vivo* PCR (Bikandi et al., 2004; Ethica et al., 2013; Ethica, 2014; Kalendar et al., 2017; Priyadharsini et al., 2018).

*In silico* PCR method has been widely known as capable to be used for investigating genes in *A. baumannii* (an emerging fish pathogen) and its associated species. Moreover, accomplished development of a real-time PCR analysis supported by *in silico* PCR to amplify signature genes to quantify a group of bacteria that causes sepsis has also been widely used (Bellemain et al., 2010; Gupta et al., 2017). In this investigation, *sagC* and *sagD* genes of *S. iniae* are tested for their potential as DNA biomarkers of the bacterial species. The aim is to seek if they could be used as DNA biomarkers, which are important to facilitate early rapid diagnosis of *S. iniae* streptococcal infection.

## 2. Material and Methods

Literature study was initially conducted to get information about the most unique phenotypic feature of *S. iniae*, particularly on specific toxins it produces. Sequence of these genes were then retrieved from genbank NCBI (National Center for Biotechnology Information) under accession number of NZ\_CP010783.1 (*sagC*, Gene ID: 35765491) and NZ\_CP010783.1 (*sagD*, Gene ID: 35766625) (Rajoo et al., 2015). DNA sequences of *sagC* gene used as the basis of SagCF-SagDR primer design along with its 3-D protein model are shown in Figure 1 and 2.

Next, the genotype feature related with genes associated with proteins underlining the unique phenotype was targeted. Primers were then designed from internal part of open reading frame of the targeted gene sequences using Primer3Plus (Ethica et al., 2013; 2014; 2019; Untergasser et al., 2007). Two pairs of primers that have the least possibility of hairpin formation, self-complementarity, and dimerization was selected (Ethica et al., 2017). The newly designed primers were used as input for the web-based *in silico* PCR, which was run from <http://insilico.ehu.es/PCR/> using all streptococcal genomes retrieved from its source of database (Bikandi et al., 2004; Ethica et al., 2019; San Millán et al., 2013). A sequence analysis was finally conducted by clicking DNA band appeared as PCR product in program output or using 'Amplify Show Result' tool. This step was needed to clarify if the *in silico* PCR products (amplicons) were specific to *S. iniae* genome and were truly part of signature genes targeted as biomarker.

**Streptococcus iniae strain YSFST01-82, complete genome**  
 NCBI Reference Sequence: NZ\_CP010783.1  
[GenBank Graphics](#)  
 >NZ\_CP010783.1:c1242370-1241306 Streptococcus iniae strain YSFST01-82, complete genome  
 ATGAGATACCAATTAATAGTAATGTTCGTATTGTGCAATTTGATGATACCTTTTGTTTAGAAAAAGGAT  
 TATGGGATTTAATGAAGCTGTATTAGATTGT CACAAGAACCACAAATGTTAAAAGAAGTTTACCAGCA  
 AATTGTTGCTGACCTCCTAAAAGGGC TAGTGGTCGATACAGATGATTGAAAAAGAGTTAGAAGCTGAA  
 TTATTTGCTAAAGTGATGGAAGTCATGACAGCTCTATATTATAAATGATGTACTCATGTTTGAAGATGATT  
 ACGCCCTTGAAGAAAATGTTATGAAAATGATTGATGGGAAATTTCCGCTTTATGGCCCAAGAAGGGCATA  
 TAAGAATACGGACCCTGTCTTGTTCATCAGTGA CTCTAGTTATGTTAATGACTCAGCAAAGTTACTTGCA  
 GAACATTTGAATTTGAACCTTGTGTGGCAAGT GATGAACATAAAATGCTGATTCAAAGATGATGTTA  
 CCTCTCGTTAGATGCAT TAGAACACCATCGCAATATGAAGCGTTTATCTGGAAAAGTTAATGCCTTATCA  
 AAGTATTGTAGTTTGCCAGGAGCGTTTGAACAT AATGATGCTTCGTCATCTTAATGAAATCAGTGTGCT  
 TTGAAAAAGCAAATGGTA CTGTGGTTTGTAGATGGTCTTTTATTCATGCTTGTACCTTAAACCCGCCG  
 ATAGCCGAGATTTTGATA GTTTAGAAAGAAGGGTTTATAGCTCGTTGCAAGATCATGCTCTTTATCAGCA  
 TTTTGCCAGCCAAATGTTGCCAACGACGCAACCAAGTTAGTCATTCCTATCTGCCTTTATGAAATATTTA  
 ATGAATTTAGTTGTAGT GAGGCCCTTATTTATGCACAACAGGAAAGTTCCAAATTTGAAGGGCGATTGT  
 TGAGTATTTATTTGCCAACACTTTGAGATTCAGT TCAAGATATTTGAAAATGTTCAAATTCCAAACCA  
 AAGTGCACCTGCGAAAATTAAGGTATGAGGACCACAAATTCGACACGAGAAATGTCAAACCTCCTA  
 AAGAAGATTCCTAA

Figure 1. Gene sequence of *sagC* used as the basis of SagCF-SagCR primer design




Figure 2. The 3-D structure of V4QUH6\_STRIN V4QUH6 Streptolysin S biosynthesis protein *SagC* of *Streptococcus iniae* visualized by SWISS-MODEL (Created on December 9, 2019)

### 3. Results and Discussion

From literature studies, it is known that *sagC* and *sagD* genes are among the unique genotypic features of virulence bacterium *Streptococcus iniae*, the main cause of streptococcal infection in tilapia fish (Datta et al., 2005). Sequences of these genes (both genes were under accession number of NZ\_CP010783.1) were used as the basis of primer design (the genes and their encoded protein are displayed in Figure 1 & 2 respectively). Both 3D structures of *SagC* and *SagD* are displayed as validity test if the genomic DNA sequences used to design primer (*sagC* and *sagD*) produce folded (functional) proteins. Figure 1 and 2 showed that both proteins are folded, and hence functional. This opens possibility to use both *SagC* and *SagD* proteins as bacterial protein markers or to develop vaccines using both proteins as target. DNA sequences of *sagD* gene from NCBI Genbank used as the basis of SagDF-SagDR primer design along with its 3-D protein model are shown in Figure 3 and 4. Using primer3Plus, pairs of primers obtained by using *sagC* and *sagD* DNA sequences as input (Figure 1 and 2 respectively) are listed in Table 1.

*In silico* PCR using SagCF-SaGR primers and genomic DNA of 117 streptococci resulted in single 558-bp DNA band (Figure 5) belonging to *S. iniae* SF1

or species no 29 (Table 2). That using SagDF-SagDR primers resulted in also a single amplicon sized 590 bp (Figure 5) on *S. iniae* SF1 or species no. 29 according to Streptococci list in Table 2. Using *in silico* PCR run on <http://insilico.ehu.es/PCR/>, the resulted amplicon could be seen on Figure 5 and Figure 6.

*In silico* PCR conducted in this study was run directly from <http://insilico.ehu.es/PCR/>. Table 2 shows all current Streptococci stored in the program database (by choosing the menu APPLY TO ALL Streptococcus when selecting genomic templates of *in silico* PCR) by the time this study is conducted. Currently, 3 genome sequences of *S. iniae* other than SF1 strain have been released (GenBank accession nos.: BANM00000000, AMO00000000, and AOCT00000000). Yet, the only representative of *S. iniae* species sequence available in the program is strain SF1 (fish isolate of sero<sup>35</sup>e 1), which is the cause <sup>35</sup>in epidemic broke out in a fish farm in north China (Cheng et al., 2010). Although not all strains of *S. iniae* could be tested using the current <http://insilico.ehu.es> database, BLAST test showed that among other *S. iniae* strains, the *SagC* and *SagD* are highly conserved (99-100% homology level) in the species group, compared with other *Streptococcus* species (starting from *S. equi* by 84%).

Streptococcus iniae strain YSFST01-82, complete genome  
 NCBI Reference Sequence: NZ\_CP010783.1  
[GenBank Graphics](#)  
 >NZ\_CP010783.1:c1241293-1239935 Streptococcus iniae strain YSFST01-82, complete genome  
 ATGTTACATTATTACCGTCCTTTAACCATATTTAGATGAACCTCAAAGGTTTGAGTGGCAATAGAACAG  
 GGATTTTAAATCAATCACAAAGTTCAGTTTGTAAACCATCAACATGATGTACTTAAAAAGTGTGACTGG  
 TCAGATACCTGATTATCACAAACAATATATTGATGAATTGAGTCAAGTTAGCTATCATATTATTGTTAT  
 GGCAGTCATTATGAAGAAGCAC TTATTAATATTTAGGTGAGAGTATTGAACGCTATGCAACTATTATTG  
 CAGGTGATTTGCTTTTCAGATCGCA TTGTTTACGCTTCTTATAAGGAACTCAGCCAAATCCAATCCTGTCTAT  
 GCCATTAGAATATTTACAGGTTTT TACACAGGAACAATTTGATCAGTCTGCGAAAATTCATATGCATATG  
 TGTGAAAAAATGGTGACTGAAGAT GATGTTTTAGGTTGGGTGAAATGCCCATGTTTTTTGAAGATAAGG  
 AACTGTATGTGCTTATTCAGATGCTTTGCATTGGCTATAAGCCATATCGTGAAGTAGGTGAACAGTATGT  
 TATTCCTGGATTTTCACTGGTACGGCTTCTATAAAACACTTGAAGCAGCTATGTGTAATAGTCTAATT  
 GAATACATTCAAATGACTCAATGATGCTTAGCTGGCATACCAAAAAAGCTTGTCCAAGAACTATTATTG  
 ATGATCCAGACATTCAAGCTATTT TGGAGAAGCAAGACTTGGTAAAGCAGCTTGTATGAGATTATCCC  
 AATTGACATGACTGTTGGAGAGGA CAATCCCTTATATACCTTTGGTATTATCCTTAAAAATAAATGAA  
 GAAGGTCCTACCTTTTATTGGT GTTCAGGCTGGTTAGATCCCAAGCATACTACTAAGAGGGGTTA  
 TGGAACTGCAGCCATTAGTTTACA GTTACTATTATAATCTTTTATCAAAGGAATCTTTAGCAAATAT  
 TGAGAGGATTCGCTTTGTTCTT GGACTTAGATAGCAATGTCTTTTATTACGCTCATCCTAAGGATCAA  
 GAGCATAAATGGAAAGCTTTTGAACCTTTAATTTTCAGGTGAGATTGTCTGAGTGATTTAAAAATAATG  
 CAGGTCAGATAAAGAAAGGAAAAATC TGAATAATCTATTAGCTTATACTAAAAAAGTGAAGTCTAATGCAAT  
 ATTTTATAGACATTACACCTCCAGAGCTTCTGAAAAAGGTTGGTACCTCACACAGAGTGTAAACACCTGAA  
 CTTCTTGAATGTGTAATCCAGCAATTCCTATTGCTAATCACCCAGAAATGAGACAGTTTGGAGGTGTTA  
 CCAATGAATTTGTCCACCCAATGCCTTAA

Figure 3. Gene sequence of *sagD* used as the basis of SagDF-SagDR primer design




Figure 4. The 3-D structure of V4R003\_STRIN V4R003 Streptolysin S biosynthesis protein SagD of *S. iniae* visualized by SWISS-MODEL (Created on December 9, 2019)

Table 1. Primers designed using Primer3Plus from *sagC* and *sagD* gene sequences

Gene Source	42 Primer Name	Forward Primer	Reverse Primer	Amplicon Size (bp)
<i>sagC</i>	SagCF-SagCR	TGCTGACCTCCTAAAAGGGC	CTATGCGCGGGTTTAAGGT	558
<i>sagD</i>	SagDF-SagDR	GCCAATCCAATCCTGTCTATGC	TGCAGCTTCATAACCCCTC	590

After carrying out the *in silico* PCR, it was necessary to check whether amplicon products, which were seen on virtual gel electrophoresis (Figure 5 and 6), were correct based on primer positions. The analysis was carried out easily by clicking amplicon size appeared on the *in silico* PCR result window. New tab was then appeared showing that *in silico* amplicons were truly parts of *sagC* and *sagD*. Results appeared on the new tabs from two PCR runs (for SagCF-SagCR and SagDF-SagDR primer pairs of each) are displayed in Figure 7 and 8.

Infectious diseases are a group as ailments caused by virulence microbes, such as viruses, bacteria, or

parasites. Those types of diseases have been major issues worldwide affecting public health and economy of many countries. Investigation on the ideal biomarkers for diagnostics of infectious diseases (with high sensitivity, specificity, and predictive capacity) in the beginning must be focused on detection and identification of the virulence agents (Hwang et al., 2018; Mohan & Harikrishna, 2015). Serious infection outbreaks often present from fish and other foods creating a diagnostic challenge for clinicians. There are not many official guidelines to help medical scientists in determining which biomarkers are useful to diagnose the presence of

Table 2. List of genomic DNA of Streptococcus species used as *in silico* PCR templates

Complete list of Streptococcus strains in the Database of <i>in silico</i> PCR	
1 - Streptococcus agalactiae 09mas018883	60 - Streptococcus pneumoniae R6
2 - Streptococcus agalactiae 2-22	61 - Streptococcus pneumoniae SPNA45
3 - Streptococcus agalactiae 2603V/R	62 - Streptococcus pneumoniae ST556
4 - Streptococcus agalactiae A909	63 - Streptococcus pneumoniae TCHB431/19A
5 - Streptococcus agalactiae GD201008-001	64 - Streptococcus pneumoniae TIGR4
6 - Streptococcus agalactiae LRD05	65 - Streptococcus pneumoniae Taiwan19F-14
7 - Streptococcus agalactiae LRH12	66 - Streptococcus pneumoniae gamPNO373
8 - Streptococcus agalactiae NEM316	67 - Streptococcus pseudopneumoniae IS7493
9 - Streptococcus agalactiae SA20-06	68 - Streptococcus pyogenes A20
10 - Streptococcus anginosus C1051	69 - Streptococcus pyogenes Alab49
11 - Streptococcus anginosus C238	70 - Streptococcus pyogenes HSC5
12 - Streptococcus constellatus subsp. pharyngis C1050	71 - Streptococcus pyogenes MI 476 DNA
13 - Streptococcus constellatus subsp. pharyngis C232	72 - Streptococcus pyogenes MI GAS
14 - Streptococcus constellatus subsp. pharyngis C818	73 - Streptococcus pyogenes MGAS10270
15 - Streptococcus dysgalactiae subsp. equisimilis 167	74 - Streptococcus pyogenes MGAS10394
16 - Streptococcus dysgalactiae subsp. equisimilis AC-2713	75 - Streptococcus pyogenes MGAS10750
17 - Streptococcus dysgalactiae subsp. equisimilis ATCC 12394	76 - Streptococcus pyogenes MGAS15252
18 - Streptococcus dysgalactiae subsp. equisimilis GGS_124	77 - Streptococcus pyogenes MGAS1882
19 - Streptococcus dysgalactiae subsp. equisimilis RE378	78 - Streptococcus pyogenes MGAS2096
20 - Streptococcus equi subsp. equi 4047	79 - Streptococcus pyogenes MGAS315
21 - Streptococcus equi subsp. zooepidemicus	80 - Streptococcus pyogenes MGAS5005
22 - Streptococcus equi subsp. zooepidemicus ATCC 35246	81 - Streptococcus pyogenes MGAS6180
23 - Streptococcus equi subsp. zooepidemicus str. MGCS10565	82 - Streptococcus pyogenes MGAS9429
24 - Streptococcus gallolyticus UCN84	83 - Streptococcus pyogenes NZ131
25 - Streptococcus gallolyticus subsp. gallolyticus ATCC 43143	84 - Streptococcus pyogenes SSI-1
26 - Streptococcus gallolyticus subsp. gallolyticus ATCC BAA-	85 - Streptococcus pyogenes str. Manfredo
27 - Streptococcus gordonii str. Challis substr. CH1	86 - Streptococcus pyogenes strain MGA S8232
28 - Streptococcus infantarius subsp. infantarius CJ18	87 - Streptococcus salivarius 57.1
29 - Streptococcus iniae SF1	88 - Streptococcus salivarius CCHSS3
30 - Streptococcus intermedius B196	89 - Streptococcus salivarius JMB777
31 - Streptococcus intermedius C270	90 - Streptococcus sanguinis SK36
32 - Streptococcus intermedius JTH08	91 - Streptococcus sp. I-G2
33 - Streptococcus lutetiensis 033	92 - Streptococcus sp. I-P16
34 - Streptococcus macedonicus ACA-DC 198	93 - Streptococcus suis 05ZYH33
35 - Streptococcus mitis B6	94 - Streptococcus suis 98HAH33
36 - Streptococcus mutans GS-5	95 - Streptococcus suis A7
37 - Streptococcus mutans LJ23	96 - Streptococcus suis BM407
38 - Streptococcus mutans NN2025	97 - Streptococcus suis D12
39 - Streptococcus mutans UA159	98 - Streptococcus suis D9
40 - Streptococcus oligofermentans AS 1.3089	99 - Streptococcus suis GZ1
41 - Streptococcus oralis Uo5	100 - Streptococcus suis JS14
42 - Streptococcus parasanguinis ATCC 15912	101 - Streptococcus suis P1/7
43 - Streptococcus parasanguinis FW213	102 - Streptococcus suis S735
44 - Streptococcus parauberis KCTC 11537	103 - Streptococcus suis SC070731
45 - Streptococcus pasteurianus ATCC 43144	104 - Streptococcus suis SC84
46 - Streptococcus pneumoniae 670-6B	105 - Streptococcus suis SS12
47 - Streptococcus pneumoniae 70585	106 - Streptococcus suis ST1
48 - Streptococcus pneumoniae A026	107 - Streptococcus suis ST3
49 - Streptococcus pneumoniae AP200	108 - Streptococcus suis T15
50 - Streptococcus pneumoniae ATCC 700669	109 - Streptococcus suis TL13
51 - Streptococcus pneumoniae CGSP14	110 - Streptococcus suis YB51
52 - Streptococcus pneumoniae DB9	111 - Streptococcus thermophilus CNRZ1066
53 - Streptococcus pneumoniae C64	112 - Streptococcus thermophilus JM 8232
54 - Streptococcus pneumoniae Hungary19A-6	113 - Streptococcus thermophilus LMD-9
55 - Streptococcus pneumoniae NV104	114 - Streptococcus thermophilus LMG 18311
56 - Streptococcus pneumoniae NV200	115 - Streptococcus thermophilus MN-ZLW-002
57 - Streptococcus pneumoniae JJA	116 - Streptococcus thermophilus ND03
58 - Streptococcus pneumoniae OXC141	117 - Streptococcus uberis 0140J
59 - Streptococcus pneumoniae PI031	

Note: Yellow highlighted is genomic sequence of *S. iniae* strain from database used as template of both primer design and *in silico* PCR amplification




Figure 5. *In silico* PCR using SagCF-SaGR primers and genomic DNA of streptococci resulted in a single 558-bp DNA band belonging to strain no. 29 or *S. iniae* SF1 species




Figure 6. *In silico* PCR using SagDF-SagDR primers resulted in also a single amplicon sized 590 bp on strain no. 29 or *S. iniae* SF1 species

infections by bacteria (Rogers et al., 2011). Yet, the initial step of bacterial biomarker development to detect *S. iniae* had been carried out in this study. It was expected that in case of streptococcal infection outbreaks, the involvement of *S. iniae* could be determined by detecting the presence of their DNA biomarkers.

44

*In silico* PCR is a computational polymerase chain reaction allowing amplification of targeted DNA sequences using updated sequenced bacterial genome

available in the program database. *In silico* PCR aimed to predict ability of primers to amplify targeted DNA segments, which could support accurate *in vitro* DNA amplification.

In this *in silico* PCR study, *sagC* and *sagD* were predicted to be potential DNA biomarkers for *S. iniae*. This is based on initial literature study (Fuller et al., 2002) and NCBI search showing that both genes encoded predicted products with significant amino acid identity to corresponding *sag* products of GAS, yet

```
>NC_021314, from 1246525 to 1247082 (558 bp); Streptococcus iniae SF1
CTATGCGGCGGGTTAAGGTACAAGCATGAAATAAAGGACCATCTACAAAACCAAGTACCATTTGCTTTT
TCAAAGCAACACTGATTTTCATTAAGATGACGAAGCATCATATGTTCAAACGCTCCTGGCAAAC TACAAT
ACTTTGATAAAGGCATTAACCTTCCAGATAAA CGC TTCATATGCGATGGTGTCTAATGCATCTAAACGA
GAGGTAACATCAATCTTTTGAATCAGCATTTT TTAGTTCATCAC TTGCCACAACAAGGTTCAAAT TCAAAT
GTTCTGCAAGTAACTTTGCTGAGTCATTAACATAACTAGAGTCACTGATAAACAAGACAGGGTCCGTATT
CTTAGTATGCCCTTCTTGGGCCATAAAGCGGAAA TTCCCATCAATACTTTTCATAACATTTTCTTCAAGG
CGGTAATCATCTTCAAACATGAGTACATCAT TATAATATAGAGTGTGCATGACTTCCATCACTTTAGCAA
ATAATTCAGCTTCTAACTCTTTTTCATAATCATCTGTATC GACCAC TAGCCCTTTTAGGAGGTCAGCA
streptolysin associated protein SagC [Streptococcus iniae]
NCBI Reference Sequence: YP_008056992
```

Figure 7. *In silico* PCR amplicons using SagCF-SagCR primers

```
>NC_021314, from 1245231 to 1245820 (590 bp); Streptococcus iniae SF1
TGCAGCTTCCATAACCCCTCTTAGTAGTGTATGCTTGGGATCTAAACCAGCCTGAACACCAAATAAAAGG
TAGGACCTTCTTCATATTTATTTTAAGGATAATACCAAAGGTATATAGGGGATTTGCTCTC CAACAG
TCATGTC AATTGGGATAATCTCATAAAGCTGCTTTACCAAGTCTTGCTTCTTCCAAAATAGCTTGAAT
GTCTGGATCATCAATAATGATTCTTGGACAAGCTTTT TGGTATGCCAGCTAAGCATATTGAGTCAATT
TGAA TGTATCAATTAGACTATTACACATAGCTGCTT CAAGTGT TTTATGAGAAGCCGTACCAGTTGAAA
ATCCAGGAATAACATACTGTTCACTTACACGATTAGGCTTATAGCCAATGCAAAGCATCTGAATAGG
CACATACAGTTCCTTATCTTCAAAAACATGGGGCATTTCAACC CAACCTAAAACATCATCTTCAAGTCACC
ATTTTTTTCACACATATGCATATGAAGTTTGCACGACTGATCAATTTGTTCTGTGTAAAAACCTGTAAT
ATTCTAATGGCATGACAGGATTGGATTGGC
streptolysin associated protein SagD [Streptococcus iniae]
NCBI Reference Sequence: YP_008056991
```

Figure 8. *In silico* PCR amplicons using SagDF-SagDR primers

shared no homology with other proteins in the GenBank database. Both genes also showed high homology level (99-100%) among *S. iniae* group of species. To prove the hypothesis, 2 pairs of primers, SagCF-SagCR and SagDF-SagDR, were successfully designed using web-based Primer3Plus software based on *S. iniae sagC* and *sagD* gene sequences.

The first designed specific pair of primers, SagCF: '5- TGCTGACCTCCTAAAAGGGC-3' and SagCR: '5- CTATGCGGCGGGTTAAGGT-3' were able to selectively amplify internal part of *sagC* gene fragment of only from genomic sequence of bacterial species *S. iniae* SF1, but not those of 116 other streptococci. The second designed specific pair of primers SagDF: '5- GCCAATCCAATCCTGTGCATGC-3' and SagDR: '5- TGCAGCTTCCATAACCCCTC-3' could also selectively amplify internal part of *sagD* gene fragment of only from genomic sequence of bacterial species *S. iniae* SF1, but not those of 116 other streptococci. The results infer that:

1. The *sagC* and *sagD* are biomarkers for bacterial species, *Streptococcus iniae*, which differentiate it from other streptococci. As bacterial biomarkers, both genes have potential to facilitate early rapid diagnosis of *S. iniae* streptococcal infection;

2. The newly designed primers SagCF-SagCR and SagDF-SagDR are potential to be used as primers to detect the presence of *S. iniae* in a streptococcal samples using *in vitro* PCR.

Results from this work is based on *in silico* or prediction study and are limited using resource available from <http://insilico.ehu.es/PCR>. As soon as the new database is updated, particularly when new genomic DNA sequence of *S. iniae* new strains are added, new *in silico* PCR tests could be consequently conducted using primers designed in this study to test their specificity. However, these results could be used to support successful DNA amplification using an *in vivo* PCR as a way to early detect and identify the causing agent of streptococcosis, thus help in controlling the *S. iniae* infection.

#### 4. Conclusion

Based on *in silico* PCR carried out in this study, *sagC* and *sagD* genes were theoretically proved to be DNA biomarkers of the fish and human pathogen *S. iniae*, which could be used to selectively differentiate the species from other streptococci. The newly



designed primers designed from these genes in this study bear the potential as *in vitro* PCR tools to facilitate early rapid diagnosis of *S. iniae* streptococcal infection.

### Acknowledgment

This work is part of collaboration output between two institutions, Universitas Muhammadiyah Semarang and Research Center for Marine and Fisheries Product Processing and Biotechnology, Ministry of Marine Affairs and Fisheries of the Republic of Indonesia. The authors owe Dr. Dewi Seswita Zilda deep gratitude for taking a keen interest in this work and guiding manuscript preparation.

### References

- 7 Agnew, W., & Barnes, A. C. (2007). *Streptococcus iniae*: an aquatic pathogen of global veterinary significance and a challenging candidate for reliable vaccination. *Veterinary Microbiology*, 122(1-2), 1-15.
- Baiano, J. C., & Barnes, A. C. (2009). Towards control of *Streptococcus iniae*. *Emerging Infectious Diseases*, 15(12), 1891.
- Bellemain, E., Carlsen, T., Brochmann, C., Coissac, E., Taberlet, P., & Kausrud, H. (2010). ITS as an environmental DNA barcode for fungi: an *in silico* approach reveals potential PCR biases. *BMC Microbiology*, 10(1), 189.
- 6 Bikandi, J., Millán, R. S., Rementeria, A., & Garaizar, J. (2004). *In silico* analysis of complete bacterial genomes: PCR, AFLP-PCR and endonuclease 10 restriction. *Bioinformatics*, 20(5), 798-799.
- Cheng, S., Hu, Y. H., Jiao, X. D., & Sun, L. (2010). Identification and immunoprotective analysis of a *Streptococcus iniae* subunit vaccine candidate. *Vaccine*, 28(14), 2636-2641.
- Datta, V., Myskowski, S. M., Kwinn, L. A., Chiem, N., Varki, N., Kansal, R. G., Kotb, M. and Nizet, V. (2005). Mutational analysis of the group A streptococcal operon encoding streptolysin S and its virulence role in invasive infection. *Molecular Microbiology*, 56(3), pp.681-695.
- Ethica, S. N. (2014). Detection of genes involved in glycerol metabolism of *Alcaligenes* sp. JG3 (Doctoral dissertation, Universitas Gadjah Mada).
- Ethica, S. N., Hammi, M. K., Lestari, F., Semiarti, E., Widada, J., & Raharjo, T. J. (2013). Amplification of *Azospirillum* sp. JG3 *glpd* gene fragment using degenerate primers generated by web-based tools. *The Journal of Microbiology, Biotechnology and Food Sciences*, 3(3), 231.
- Ethica, S. N., Sulis, I., Ingtyas, A. R., & Darmawati, S. (2019, June). *In-silico* Specificity Comparison between GMF-GMR and JMF-JMR Primers for Detecting *moaC* Gene of Food Spoilage Bacteria *Pseudomonas* spp. In *IOP Conference Series: Earth and Environmental Science* 292(1), 012033. IOP Publishing.
- Ethica, S. N., Semiarti, E., Widada, J., Oedjiono, O., & Joko Raharjo, T. (2017). Characterization of *moaC* and a nontarget gene fragments of food borne pathogen *Alcaligenes* sp. JG3 using degenerate colony and arbitrary PCRs. *Journal of Food Safety*, 37(4), e12345.
- 15 Fuller, J. D., Camus, A. C., Duncan, C. L., Nizet, V., Bast, D. J., Threlkeld, R. L., Low, D. E. and De Azavedo, J. C., (2002). Identification of a streptolysin S-associated gene cluster and its role in the pathogenesis of *Streptococcus iniae* disease. *Infection and Immunity*, 70(10), 5730-5739.
- Gu, S., Dongre, A., Saxena, J., & Jyoti, A. (2017). Computation and *in silico* validation of a real-time PCR array for quantitative detection of pathogens isolated from blood sample in sepsis patients. *IJBT*, 16(3), 505-508.
- 32 Hwang, H., Hwang, B. Y. & Bueno, J. (2018). Biomarkers in infectious diseases. *Disease Markers*, 2018.
- 13 Kalendar, R., Muterko, A., Shamekova, M., & Zhambakin, K. (2017). *In Silico* PCR Tools for a Fast Primer, Probe, and Advanced Searching. In PCR (pp. 1-31). Springer, New York, NY.
- 14 Le Breton, Y., & McIver, K. S. (2013). Genetic manipulation of *Streptococcus pyogenes* (the Group A Streptococcus, GAS). *Current Protocols in Microbiology*, 30(1), 9D-3.
- 5 Locke, J. B., Colvin, K. M., Varki, N., Vicknair, M. R., Nizet, V., & Buchanan, J. T. (2007). *Streptococcus iniae*  $\beta$ -hemolysin streptolysin S is a virulence factor in fish infection. *Diseases of Aquatic Organisms*, 76(1), 17-26.
- 9 Mohan, A., & Harikrishna, J. (2015). Biomarkers for the diagnosis of bacterial infections: in pursuit of the 'Holy Grail'. *The Indian Journal of Medical Research*, 141(3), 271.
- 29 Moretti, S. (2011). *In silico* experiments in scientific papers on molecular biology. *Science & Technology Studies*, 24(2).
- 27 Nho, S. W., Shin, G. W., Park, S. B., Hwang, H. B., Cha, I. S., Ha, M. A., ... & Joh, S. J. (2009). Phenotypic characteristics of *Streptococcus iniae* and *Streptococcus parauberis* isolated from olive flounder (*Paralichthys olivaceus*). *FEMS Microbiology Letters*, 293(1), 20-27.
- 3 Ortega, C. E. S. A. R., García, I., Irgang, R., Fajardo, R., Tapia Cammas, D., Acosta, J., & Avendaño Herrera, R. (2018). First identification and characterization of *Streptococcus iniae* obtained from tilapia (*Oreochromis aureus*) farmed in Mexico. *Journal of Fish Diseases*, 41(5), 773-782.
- 12 Priyadharsini, J. V., Girija, A. S., & Paramasivam, A. (2018). *In silico* analysis of virulence genes in an emerging dental pathogen *A. baumannii* and related species. *Archives of Oral Biology*, 94, 93-98.

11

Rahmatullah, M., Ariff, M., Kahieshesfandari, M., Daud, H. M., Zamri-Saad, M., Sabri, M. Y., ... & Ina-Salwany, M. Y. (2017). Isolation and pathogenicity of *Streptococcus iniae* in cultured red hybrid tilapia in Malaysia. *Journal of Aquatic Animal Health*, 29(4), 208-213.

Rajoo, S., Jeon, W., F25, K., Yoo, S., Yoon, I., Lee, H., & Ahn, J. (2015). Complete genome sequence of *Streptococcus iniae* YSFST01-82, isolated from olive flounder in Jeju, South Korea. *Genome Announc.*, 3(2), e00319-15.

Rogers, G. B., Hoffman, L. R., Johnson, M. W., Mayer-Hamblett, N., Schwarze, J., Carroll, M. P., & Bruce, K. D. (2011). Using bacterial biomarkers to identify early indicators of cystic fibrosis pulmonary exacerbation onset. *Expert Review of Molecular Diagnostics*, 11(2), 197-206.

2

San Millán, R. M., Martínez-Ballesteros, I., Rementería, A., Garaizar, J., & Bikandi, J. (2013). Online exercise for the design and simulation of PCR and PCR-RFLP experiments. *BMC Research Notes*, 6(1), 513.

Shanks, O. C., San20omingo, J. W., Lu, J., Kelty, C. A., & Graham, J. E. (2007). Identification of bacterial DNA markers for the detection of human fecal pollution in water. *Appl. Environ. Microbiol.*, 73(8), 2416-2422.

1

Untergasser, A., Nijveen, H., Rao, X., Bisseling, T., Geurts, R., & Leunissen, J. A. (2007). Primer3Plus, an enhanced web interface to Primer3. *Nucleic Acids Research*, 35(38: ppl\_2), W71-W74.

Zhu315. M., Fan, Y., Zhu, X. Q., Xie, M. Q., & Li, A. X. (2011). Rapid identification of *Streptococcus iniae* by specific PCR assay utilizing genetic markers in ITS rDNA. *Journal of Fish Diseases*, 34(4), 265-271.

# Streptolysin Encoding Genes sagC and sagD as Biomarkers of Fish Pathogen Streptococcus iniae: An In Silico Study

## ORIGINALITY REPORT

20%

SIMILARITY INDEX

13%

INTERNET SOURCES

11%

PUBLICATIONS

9%

STUDENT PAPERS

## PRIMARY SOURCES

1

[digitalcommons.library.umaine.edu](https://digitalcommons.library.umaine.edu)

Internet Source

1%

2

A.S. Machado, S. Lyon, F. Rocha-Silva, C.B. Assunção et al. " Novel PCR primers for improved detection of and diagnosis of leprosy ", Journal of Applied Microbiology, 2020

Publication

1%

3

Herng Yih Tan, Sai-Wei Chen, Shao-Yang Hu. "Improvements in the growth performance, immunity, disease resistance, and gut microbiota by the probiotic Rummeliibacillus stabekisii in Nile tilapia (Oreochromis niloticus)", Fish & Shellfish Immunology, 2019

Publication

1%

4

[appliedbiolchem.springeropen.com](https://appliedbiolchem.springeropen.com)

Internet Source

1%

5

[www.currentprotocols.com](https://www.currentprotocols.com)

Internet Source

1%

[studentsrepo.um.edu.my](https://studentsrepo.um.edu.my)

6	Internet Source	1%
7	fa.wikipedia.org Internet Source	1%
8	www.ecfs.eu Internet Source	1%
9	Submitted to Chester College of Higher Education Student Paper	1%
10	Khalid Shahin, Kim D. Thompson, Neil F. Inglis, Kevin Mclean et al. " Characterisation of the outer membrane proteome of subsp. ", Journal of Applied Microbiology, 2018 Publication	1%
11	Md. Shirajum Monir, Md Sabri bin Mohd Yusoff, Zarirah binti Mohamed Zulperi, Hasliza binti Abu Hassim et al. "Haemato-immunological responses and effectiveness of feed-based bivalent vaccine against Streptococcus iniae and Aeromonas hydrophila infections in hybrid red tilapia (Oreochromis spp.)", Research Square, 2020 Publication	1%
12	Submitted to Monash University Student Paper	1%

Submitted to Flinders University

13

Student Paper

1%

14

Submitted to Liverpool John Moores University

Student Paper

1%

15

homepage.mac.com

Internet Source

1%

16

es.scribd.com

Internet Source

1%

17

www.blalbiotech.com

Internet Source

&lt;1%

18

microbiology.mtsinai.on.ca

Internet Source

&lt;1%

19

acadstaff.ugm.ac.id

Internet Source

&lt;1%

20

Submitted to East Tennessee State University

Student Paper

&lt;1%

21

Submitted to Curtin University of Technology

Student Paper

&lt;1%

22

ichestech.unimus.ac.id

Internet Source

&lt;1%

23

Submitted to Nottingham Trent University

Student Paper

&lt;1%

24

journal.ui.ac.id

Internet Source

&lt;1%

25	<a href="http://www.genome.jp">www.genome.jp</a> Internet Source	<1%
26	<a href="http://www.limnofish.org">www.limnofish.org</a> Internet Source	<1%
27	Submitted to South Dakota Board of Regents Student Paper	<1%
28	Makio Yokono, Soichirou Satoh, Ayumi Tanaka. "Comparative analyses of whole-genome protein sequences from multiple organisms", Scientific Reports, 2018 Publication	<1%
29	<a href="http://sciencetechnologystudies.journal.fi">sciencetechnologystudies.journal.fi</a> Internet Source	<1%
30	Merlyn D. Manurung, Reiny A. Tumbol, Henneke D. Pangkey, Deiske A. Sumilat, Remy E. P. Mangindaan. "The use of Sponge Crude Extract to Increase Growth and Immune Response of Nile Tilapia Oreochromis niloticus", JURNAL ILMIAH PLATAX, 2019 Publication	<1%
31	<a href="http://www.ajas.info">www.ajas.info</a> Internet Source	<1%
32	Submitted to University of Sunderland Student Paper	<1%

Stalis Norma Ethica, Sakti Imam Muchlissin,

33 Ragil Saptaningtyas, Agus Sabdono. "Protease Producers Predominate Cultivable Hydrolytic Bacteria Isolated from Liquid Biomedical Waste", Asian Journal of Chemistry, 2018  
Publication <1%

---

34 Nizet, V.. "Streptococcal @b-hemolysins: genetics and role in disease pathogenesis", Trends in Microbiology, 20021201  
Publication <1%

---

35 Mo-fei Li, Chun-Lin Wang, Li Sun. "A pathogenic Vibrio harveyi lineage causes recurrent disease outbreaks in cultured Japanese flounder (Paralichthys olivaceus) and induces apoptosis in host cells", Aquaculture, 2011  
Publication <1%

---

36 [ejournal-balitbang.kkp.go.id](http://ejournal-balitbang.kkp.go.id)  
Internet Source <1%

---

37 Abdel-Fattah M El-Sayed. "Alternative dietary protein sources for farmed tilapia, Oreochromis spp.", Aquaculture, 1999  
Publication <1%

---

38 [www.degruyter.com](http://www.degruyter.com)  
Internet Source <1%

---

39 [tc.copernicus.org](http://tc.copernicus.org)  
Internet Source <1%

---

40 [muqtasid.iainsalatiga.ac.id](http://muqtasid.iainsalatiga.ac.id)

<1%

41

Jasper J. Koehorst, Edoardo Saccenti, Peter J. Schaap, Vitor A. P. Martins dos Santos, Maria Suarez-Diez. "Protein domain architectures provide a fast, efficient and scalable alternative to sequence-based methods for comparative functional genomics", F1000Research, 2016

Publication

<1%

42

[arthritis-research.biomedcentral.com](http://arthritis-research.biomedcentral.com)

Internet Source

<1%

43

[springerplus.springeropen.com](http://springerplus.springeropen.com)

Internet Source

<1%

44

Yongxi Zhao, Feng Chen, Qian Li, Lihua Wang, Chunhai Fan. "Isothermal Amplification of Nucleic Acids", Chemical Reviews, 2015

Publication

<1%

45

Sohn, Michael B, Lingling An, Naruekamol Pookhao, and Qike Li. "Accurate genome relative abundance estimation for closely related species in a metagenomic sample", BMC Bioinformatics, 2014.

Publication

<1%

46

Ying Liu, Changhan Lee, Fengyang Li, Janja Trček et al. "A Cyclic Di-GMP Network Is Present in Gram-Positive Streptococcus and

<1%



## Gram-Negative Proteus Species", American Chemical Society (ACS), 2020

Publication

---

47

Sangeeta Raut, Soumya Satpathy, Sudip Kumar Sen. "Screening of Potential Bioremediation Enzymes from Hot Spring Isolates and Characterization of Amylase Protein Sequences of Bacteria and Fungi: An In Silico Study", Journal of Hazardous, Toxic, and Radioactive Waste, 2020

Publication

---

<1%

48

"Lactic Acid Bacteria", Springer Nature, 2014

Publication

---

<1%

49

Rahwa Taddese, Daniel R. Garza, Lilian N. Ruitter, Marien I. de Jonge et al. "Growth rate alterations of human colorectal cancer cells by 157 gut bacteria", Cold Spring Harbor Laboratory, 2019

Publication

---

<1%

---

Exclude quotes

Off

Exclude matches

Off

Exclude bibliography

Off