

Mask Protection Against Lead Exposure and Its Correlation with Erythropoiesis in Automotive Body Painters at Ligu District, Semarang, Indonesia

by Fitriani Nur Damayanti

Submission date: 12-Apr-2022 12:26AM (UTC-0400)

Submission ID: 1808528527

File name: Mask_Protection_Against_Lead_Exposure.pdf (344.06K)

Word count: 4048

Character count: 21572

Mask Protection Against Lead Exposure and Its Correlation with Erythropoiesis in Automotive Body Painters at Ligu District, Semarang, Indonesia

Budi Santosa¹, Ali Rosidi², Herlisa Anggraini³, Zahra Maharani Latrobdiba², Fitriani Nur Damayanti⁴, Heru Santoso Wahito Nugroho⁵

¹Department of Medical Laboratory Science, Universitas Muhammadiyah Semarang, Semarang, Indonesia; ²Department of Nutrition, Faculty of Nursing and Health Science, Universitas Muhammadiyah Semarang, Semarang, Indonesia; ³Department of Medical Laboratory Technology, Faculty of Nursing and Health Science, Universitas Muhammadiyah Semarang, Semarang, Indonesia; ⁴Department of Midwifery, Faculty of Nursing and Health Science, Universitas Muhammadiyah Semarang, Semarang, Indonesia; ⁵Department of Health, Poltekkes Kemenkes Surabaya, Surabaya, Indonesia

Correspondence: Budi Santosa, Department of Medical Laboratory Science, Universitas Muhammadiyah Semarang, Jl. Kedungmundu Raya 18, Semarang, 50273, Central Java, Indonesia Email budisantosa@unimus.ac.id

Introduction: Automotive body painters are at risk of excessive lead exposure that may compromise their health. This study examined the protective effects of masks against lead exposure and its correlation with blood profile of automotive body painters at Ligu district, Semarang, Indonesia.

Material and Methods: This cross-sectional study included 61 painters at Ligu district, Semarang, Indonesia (mean age 45 years, 100% male). Mask use was categorized into “frequent” and “infrequent”. Blood samples were taken to measure lead exposure and haematological parameters (haemoglobin/HGB, erythrocytes/RBC, haematocrit/HCT, mean corpuscular haemoglobin/MCH, mean corpuscular volume/MCV, mean corpuscular haemoglobin concentration/MCHC). Effectivity of mask use against lead exposure was analysed with Fisher’s exact test, while lead exposure and haematopoietic parameters were evaluated with Pearson’s correlation tests.

Results: There were 26 automotive painters (42.6%) classified as infrequent mask users, and all of them had high blood lead levels. Most haematopoiesis parameters were in normal range. Workers with infrequent mask use were 1.269 more likely to be exposed to lead compared to those with frequent mask use (CI 95%, 1.082–1.552). Significant associations were found between lead exposure and RBC ($p=0.0$, $r=-0.53$), HGB ($p=0$, $r=-0.61$), and HCT ($p=0.00$, $r=-0.61$). No significant correlations were observed between lead exposure and MCV, MCH, MCHC.

Conclusion: Lead exposure was significantly associated with haematological parameters RBC, HGB, and HCT, confirming the effects of prolonged exposure on blood profile. However, mask use provided significant protection against lead exposure in automotive body painters and should be an obligatory gear for workers.

Keywords: mask, lead, haematopoiesis, automotive painters

Introduction

Lead is a common material found in various industries. Automotive painting, for instance, uses lead to improve paint quality in terms of brightness, solubility, and drying time.¹ As a highly toxic non-essential metal, lead comes with a number of environmental risks and health hazards.^{2–6} Regular exposure to lead may cause anaemia, kidney and liver damage.^{7,8}

The impact of lead exposure depends on the route, duration, and dose. Lead can enter the human body through ingestion, absorption, and most predominantly, inhalation (80%).⁹ Approximately half of lead exposure in children is obtained through ingestion, while it contributes to only 10–15% in adults due to more advanced hygiene practices.^{10–12}

Car painters are exposed to lead during the entire painting process, starting from sanding, priming with epoxy, mixing paint with thinner, paint spraying until finishing with varnish. Lead may enter the body through inhalation, absorption

through the skin, or ingestion by unhygienic hand-to-mouth transmissions. Lead is mostly absorbed through skin after 12 hours of constant exposure.¹³

In the body, lead is metabolized and circulated in blood, tissues, and bones. As a result, biomarkers for lead exposure include blood, soft tissues, bones, hair, urine, sweat, teeth, nails, and bile.¹⁴ The majority (95%) of absorbed lead is carried by the erythrocytes, making it the best marker to measure lead exposure.² Lead interferes with erythropoietin, a hormone that promotes erythropoiesis, thus causing hampered erythrocyte production.¹⁵ Lead also inhibits delta-aminolevulinic acid dehydratase (ALAD) enzyme which is involved in the formation of heme, consequently reducing haemoglobin levels and altering erythrocyte volume, as seen from decreased haematocrit levels. Those changes can be observed from erythrocyte indices, particularly mean corpuscular haemoglobin (MCH), mean corpuscular volume (MCV), and mean corpuscular haemoglobin concentration (MCHC). Erythrocyte indices are useful to identify anaemia.^{16–18}

For automotive body painters, mask usage may play a big role in the amount of lead exposure. Masks serve as a protective item against lead inhalation, but its use varies between individuals from regular, occasional, to never. The objective of this study is to examine the degree of protection from wearing masks against lead exposure and its correlation with haematopoietic systems in automotive body painters in Ligu district, Semarang, Indonesia. Ligu district is in the northern area of Semarang which is a heavily populated with automotive body painting as the main home industry of the region.

13 Materials and Methods

Design and Subjects

In this cross-sectional study, data was collected in automotive body painting workshops at Ligu district, Semarang, from December 2020 to March 2021. Subjects were randomly chosen according to the following inclusion criteria: (i) provided written consent, (ii) currently works as an auto body painter in workshops at Ligu district, Semarang. Exclusion criteria were (i) has worked in the workshop for less than 5 years (because they might get lead exposure from other places), (ii) has working hours less than 8 hours in total daily work load, and (iii) is consuming multivitamin supplements. Based on the inclusion and exclusion criteria, then 61 research subjects were obtained.

Ethical Consideration

As a scientific study involving humans, this study received ethical approval prior to data collection. In order to avoid conflicts of interest, ethical approval was obtained from the Health research ethics committee from other university, namely the Health Research Ethics Committee, Faculty of Medicine, Universitas Sultan Agung Semarang, with the number: 183/VII/2121/Komisi Bioetik. The ethical approval was given based on the results of an evaluation conducted by the ethics examiner team, which showed that this study had applied ethical principles of medical and health research, especially the Declaration of Helsinki which upholds autonomy, does not harm, maintains justice and benefits for the participants, such as: counseling based on the results of related laboratory examinations. All participants also agreed to be involved as research subjects by signing an informed consent, having previously been clearly informed about the purpose and procedures of this study.

Measurements

General Characteristics of Subjects

The general characteristics of subjects (age, sex, work period, working hours/day, room ventilation, exercise habit) were obtained using a questionnaire.

Mask Use

Mask use of automotive body painters at workshops in Ligu district were observed daily for a week. Afterwards, subjects were interviewed about their average mask use during work: always, sometimes, or never. The result from the interview is later categorized as “frequent” and “infrequent” (for workers who sometimes or never wear masks). Observations and interviews were conducted simultaneously as a cross check. This was done in order to minimize bias in data collection.

Lead Levels, Haemoglobin (HGB), Erythrocytes/Red Blood Cell (RBC), Haematocrit (HCT), MCV, MCH, MCHC

Blood samples were taken from the superficial veins of antecubital fossa and then stored in purple evacuated tubes (with K2EDTA). The samples were stored in a plastic transport box and sent to the BioMedical Laboratory Universitas Muhammadiyah Semarang for analysis using a haematology analyzer by photometry and electrical impedance. The results of laboratory tests were based on the following normal limits: Lead levels (<40 µg/dL), HGB (13.3–16.7 g/dL), RBC ($4.5\text{--}5.7 \times 10^{12}$ /L), HCT (35–53%), MCV (77–98 fL), MCH (26–33 pg), MCHC (33–37 g/dL). The measurement of blood lead was done at GAKI Laboratory Universitas Diponegoro using atomic absorption spectrophotometer (AAS).

Other Data

Additional data that were collected consisted of age, gender, work period, working hours/day, ventilation, and exercise habits. Ventilation was measured based on room area and window width which should be around 10–25%. Exercise habits were categorized as “active” when done twice a week and “not active” when only done once or not at all in a week.

Statistical Analysis

Categorical variables (age, gender, work period, working hours/day, ventilation, exercise habits) were described in numbers and percentage. The degree of protection from masks were analysed using Fisher’s exact test, while the association between lead exposure and haematopoietic system were evaluated with Pearson’s correlation tests.

Results

The characteristics of subjects are presented in Table 1. Long-term work period (>5 years) and long working hours (>8 hours) were specifically used as an inclusion and exclusion criteria to determine subjects. The 61 subjects were chosen as they filled the inclusion criteria while the remaining four were excluded. We found that subjects with infrequent mask use had higher blood lead levels (100%, n=26) than those with frequent mask use (Table 2). In this case, the limit for high blood lead levels is 40 micrograms per deciliter.¹⁹ The risk of lead exposure was 1.296 times higher for infrequent mask users in comparison to frequent users (p=0.016).

Table 1 Characteristics of Subjects

Mean Age (Years)	44.23±10.33
Gender	
Male	61 (100%)
Female	0 (0%)
Work period	
≤5 years	4 (6.2%)
>5 years	61 (93.8%)
Working hours/day	
≤8 hours	4 (6.2%)
>8 hours	61 (93.4%)
Room ventilation	
Inadequate	44 (72.1%)
Adequate	17 (27.9%)
Exercise habit	
Not active	37 (60.7%)
Active	24 (39.3%)

Table 2 Relation of Blood Lead Concentrations and Mask Use in Automotive Body Painters

Mask Use	Blood Lead Levels				Total		p	RR	95% CI
	Normal (<40 µg/dL)		High (≥40 µg/dL)		n	%			
	n	%	n	%					
Infrequent	0	0	26	100	26	100	0.016	1.296	1.082–1.552
Frequent	8	22.9	27	77.1	35	100			

Most haematological parameters for all subjects were still within normal range, including MCHC (Table 3). Average blood lead levels were above normal values although it was still generally acceptable (Table 4). This refers to Decharat et al that the blood lead level that can still be tolerated is <60 micrograms per deciliter.¹⁹

Blood lead levels were significantly and strongly correlated with HGB, RBC, and HCT (−0.61, −0.53, −0.61), suggesting that higher lead exposure caused decreased levels of HGB, RBC, and HCT. On the contrary, no significant correlation was observed between blood lead levels and MCV, MCH, and MCHC.

Discussion

Results from this study affirmed that mask use largely affected the extent of lead exposure. Automotive body painters at Ligu district Semarang with infrequent mask use were 1.296 times more likely to be exposed to lead compared to those

Table 3 Haematological Parameters of Automotive Body Painters in Ligu District, Semarang, Indonesia

Parameters				Normal Limit
	Low	Normal	High	
HGB	5 (8.2%)	53 (86.9%)	3 (4.9%)	13.3–16.7 g/dL
RBC	20 (32.8%)	30 (49.2%)	11 (18.0)	4.5–5.7 × 10 ¹² /L
HCT	18 (29.5%)	43 (70.5%)	0 (0%)	35–53%
MCV	14 (23%)	41 (67.2%)	6 (9.8%)	77–98 fL
MCH	19 (31.1%)	34 (55.7%)	8 (13.1%)	26–33 pg
MCHC	0 (0%)	61 (100%)	0 (0%)	33–37 pg/dL

Table 4 Relation of Haematological Parameters with Blood Lead Levels

	Normal Limit	Average	Max	Min	Analysis		
					Correlation	p	r
Blood Pb	<40 µg/dL	42.99±4.53	52.3	27.4			
HGB	13.3–16.7 g/dL	14.01±1.33	16.1	11.3	Lead-HGB	0.00	−0.61
RBC	4.5–5.7 × 10 ¹² /L	4.94±0.61	5.98	3.99	Lead-RBC	0.00	−0.53
HCT	35–53%	42.14±4.11	48.6	34	Lead-HCT	0.00	−0.61
MCV	77–98 fL	28.53±2.42	33.3	22.7	Lead-MCV	0.63	0.06
MCH	26–33 pg	85.77±7.28	100.90	68.4	Lead-MCH	0.58	0.01
MCHC	33–37 pg/dL	33.25±0.39	35.9	32.5	Lead-MCHC	0.46	0.09

who frequently wore masks. This study also found a significant difference ($p=0.016$) between employees with frequent mask use and employees with infrequent use against lead exposure, further proving the importance of masks as a protective gear against pollutants including lead that can be absorbed through inhalation. Kuruvilla et al also reported that the use of activated carbon fabric (ASF) masks that successfully reduced lead exposure.²⁰

Our study had several limitations, particularly the type of masks. There is a wide variety of mask types, including disposable face masks, bandana masks, pre-shaped face masks, and full-face respirators.^{21,22} Each type provides different levels of protection against airborne contaminants, thus longitudinal studies with specified type of masks are required. The findings in this study are independent of our previous work where mask use had no significant correlation with cognitive deficits from lead exposure. Cognitive decline is considerably harder to measure as detectable signs may only appear after a long period of time even though lead exposure may exceed the tolerable level at present.²³

We found a strong negative correlation between blood lead concentrations with HGB, RBC, HCT. Inhalation is a major exposure route for lead since more than 40% of lead is absorbed into the respiratory system before getting distributed in blood. Approximately 95% of lead is attached to red blood cells and the rest is dissolved in plasma. Some lead is stored in tissues and bones while the excretion is primarily through kidney and digestive system.²⁴ Lead has been proven to disrupt heme synthesis and shorten the lifespan of erythrocytes, consequently decreasing HGB, RBC, and HCT concentrations. Lead interferes with the enzymatic activity in the beginning, middle, and end of heme biosynthesis. At the initial steps, lead inhibits δ ALAD enzyme, an enzyme that triggers heme biosynthesis.²⁵ As a result, the transformation of δ ALA into porphobilinogen (PBG) is blocked²⁶ and the concentration of δ ALA is increased in blood and urine. Lead also hinders coproporphyrinogen oxidase in the midway process of heme synthesis, leading to elevated coproporphyrinogen levels. The last enzyme affected by lead is ferrochelatase²⁷ which resulted in increased protoporphyrin concentration in red blood cells or known as free erythrocyte protoporphyrin.²⁸ In addition, lead has pro-oxidant activity that may incite oxidative stress in vulnerable erythrocyte membranes, thus reducing erythrocytes lifespan and eventually causing hemolytic anaemia.^{27,29}

No significant association was found between lead exposure with MCV, MCH, and MCHC. Blood lead levels in automotive body painters were generally high but still within tolerable range (40 μ g/dl-60 μ g/dl). Value threshold for blood lead range 40 g/dL. Similarly, all haematological parameters were also within normal range. Erythrocyte indices (MCV, MCH, MCHC) can be calculated from HGB, RBC, and HCT. The calculation of erythrocyte indices revealed the mean volume of erythrocytes and mean haemoglobin concentration per erythrocyte. Clinically, erythrocyte indices are used to diagnose and identify anaemia.

The results of this study are expected to provide justification that the use of masks as personal protective equipment for automotive painters is very important, because it has been proven to have an effect on lead levels in the blood which in turn can have an impact on HGB, RBC, HCT levels. Indeed, there were also changes in MCV, MCH, and MCHC levels; but still within a tolerable level. Logically, if the concentration of lead exposure studied was higher, it is possible that changes in MCV, MCH, and MCHC levels would exceed tolerable limits. Of course, to prove this, further research is needed involving automotive paint workers with a high duration and frequency of work or who are too ignorant to use masks as personal protective equipment.

Conclusion

In conclusion, the results of this study showed that automotive body painters who occasionally or never wear masks while working had higher risks for lead exposure than workers who always wear masks. We also affirmed that lead exposure significantly affected blood profile, particularly HGB, RBC, and HCT. Further longitudinal studies with specified mask types are required to be able to make recommendations of protective gears for automotive body painters.

Acknowledgments

We thank all subjects who have participated in our study, our colleagues at the laboratory who provided great help in blood analysis, and Universitas Muhammadiyah Semarang for the full support.

Disclosure

The authors declare no potential **14** conflict of interest regarding the study and the publication of this paper.

References

1. Gottesfeld P. Time to ban lead in industrial paints and coatings. *Front Public Health*. 2015;3:144. doi:10.3389/fpubh.2015.00144
2. Wani AL, Ara A, Usmani JA. Lead toxicity: a review. *Interdiscip Toxicol*. 2015;8(2):55–64. doi:10.1515/intox-2015-0009
3. Halmo L, Nappe TM. Lead toxicity. In: *StatPearls*. Treasure Island (FL): StatPearls Publishing; 2021:1–25.
4. Gidlow DA. Lead toxicity. *Occup Med*. 2015;65(5):348–356. doi:10.1093/occmed/kqv018
5. Rajkumar V, Gupta V. Heavy metal toxicity. In: *StatPearls*. Treasure Island (FL): StatPearls Publishing; 2020:1–33.
6. Gump BB, Hruska B, Parsons PJ, et al. Dietary contributions to increased background lead, mercury, and cadmium in 9–11 year old children: accounting for racial differences. *Env Res*. 2020;185:109308. doi:10.1016/j.envres.2020.109308
7. Santosa B, Iswara A, Ernanto AR. IR-bagendit leaves water extract as preventing agent in hematopoiesis, degeneration and necrosis in kidney tubules of lead-exposed rats. *Pakistan J Med Health Sci*. 2019;13(3):899–904.
8. Kwon S, Bae O, Noh J, et al. Erythrophagocytosis of lead-exposed erythrocytes by renal tubular cells: possible role in lead-induced nephrotoxicity. *Environ Health Perspect*. 2015;(2):120–127. doi:10.1289/ehp.1408094
9. Kastury F, Smith E, Lombi E, Donnelley MW, Cmielewski PL, Parsons DW. Dynamics of lead bioavailability and speciation in indoor dust and X-ray spectroscopic investigation of the link between ingestion and inhalation pathways. *Env Sci Technol*. 2019;53(19):11486–11495. doi:10.1021/acs.est.9b03249
10. Assi MA, Noor M, Hezmee M, Haron AW, Yusof M, Sabri M. The detrimental effects of lead on human and animal health. *Vet World*. 2016;9:2231–2916. doi:10.14202/vetworld.2016.660-671
11. Kumar A, Kumar A, Chaturvedi AK. Lead toxicity: health hazards, influence on food chain, and sustainable remediation approaches. *Int J Environ Res Public Health*. 2020;17:1–33. doi:10.3390/ijerph17072179
12. Leonas R, Noor Z, Rasyid HN, Madjid TH, Tanjung F. Effect of lead nanoparticles inhalation on bone calcium sensing receptor, hydroxyapatite crystal and receptor activator of nuclear factor-kappa B in rats. *Acta Inf Med*. 2016;24(5):343–346. doi:10.5455/aim.2016.24.343-346
13. Kim H, Jang T, Chae H, et al. Evaluation and management of lead exposure. *Ann Occup Environ Med*. 2015;27(1):1–9. doi:10.1186/s40557-015-0085-9
14. Charkiewicz AE, Backstrand JR. Lead toxicity and pollution in Poland. *Int J Environ Res Public Health*. 2020;17(12):1–14. doi:10.3390/ijerph17124385
15. Bunn HF. Erythropoietin. *Cold Spring Harb Perspect Med*. 2013;3:a011619. doi:10.1101/cshperspect.a011619
16. Hsieh NH, Chung SH, Chen SC, et al. Anemia risk in relation to lead exposure in lead-related manufacturing. *BMC Public Health*. 2017;17(1):1–12. doi:10.1186/s12889-017-4315-7
17. Kalahasthi R, Barman T. Effect of lead exposure on the status of reticulocyte count indices among workers from lead battery manufacturing plant. *Toxicol Res*. 2016;32(4):281–287. doi:10.5487/TR.2016.32.4.281
18. Santosa B, Angraini H, Dyah I, Muliati N, Nugroho A. Erythrocyte index of residents exposed to lead in Tambaklorok, Semarang, Indonesia. *J Teknol Lab*. 2020;9(2):151–158. doi:10.29238/teknolabjournal.v9i2.213
19. Decharat S, Kongtip P, Thampoophasiam P, Thetkathuek A. An examination of blood lead levels in Thai Nielloware workers. *Saf Health Work*. 2012;3(3):216–223. doi:10.5491/SHAW.2012.3.3.216
20. Kuruvilla A, Pillay VV, Adhikari P, et al. Role of activated carbon fabric mask to prevent lead absorption—a short report. *Toxicol Ind Health*. 2008;24(9):569–572. doi:10.1177/0748233708100367
21. Bowen LE. Does that face mask really protect you? *Appl Biosaf*. 2010;15(2):67–71. doi:10.1177/153567601001500204
22. Gutierrez AM, Galang MD, Seva RR, Lu MC, Ty DRS. Designing an improved respirator for automotive painters. *Int J Ind Ergon*. 2014;44(1):131–139. doi:10.1016/j.ergon.2013.11.004
23. Laksmidewi AA, Suputra G, Widyadharna IPE. High serum lead levels increase the incidence of cognitive impairment of public fueling station operators. *Open Access Maced J Med Sci*. 2019;7(4):599–602. doi:10.3889/OAMJMS.2019.127
24. Jain RB. Lead and kidney: concentrations, variabilities, and associations across the various stages of glomerular function. *J Trace Elem Med Biol*. 2019;54:36–43. doi:10.1016/j.jtemb.2019.03.007
25. da Cunha Martins A, Mazzaron Barcelos GR, Jacob Ferreira ALB. Effects of lead exposure and genetic polymorphisms on ALAD and GPx activities in Brazilian battery workers. *J Toxicol Env Heal A*. 2015;78(15):1073–1081. doi:10.1080/15287394.2015.1055527
26. La-llave-león O, Méndez-Hernández EM, Castellanos-Juárez FX, et al. Association between blood lead levels and delta-aminolevulinic acid dehydratase in pregnant women. *Int J Environ Res Public Health*. 2017;14(4):432. doi:10.3390/ijerph14040432
27. Flora G, Gupta D, Tiwari A. Toxicity of lead: a review with recent updates. *Interdiscip Toxicol*. 2012;5(2):47–58. doi:10.2478/v10102-012-0009-2
28. Dongre NN, Suryakar AN, Patil AJ, Ambekar JG, Rathu DB. Biochemical effects of lead exposure on systolic & diastolic blood pressure, heme biosynthesis and hematological parameters in automobile workers of North Karnataka (India). *Indian J Clin Biochem*. 2011;26(4):400–406. doi:10.1007/s12291-011-0159-6
29. Corradi M, Goldoni M, Sabbadini F, Mutti A. [Acute Lead Poisoning: A Singular Case of Hemolytic Anemia and Lead Colic]. *Med Lav*. 2011;102(3):243–249. Italian.

Journal of Blood Medicine

Dovepress

Publish your work in this journal

The Journal of Blood Medicine is an international, peer-reviewed, open access, online journal publishing laboratory, experimental and clinical aspects of all aspect pertaining to blood based medicine including but not limited to: Transfusion Medicine; Blood collection, Donor issues, Transmittable diseases, and Blood banking logistics; Immunohematology; Artificial and alternative blood based therapeutics; Hematology; Biotechnology/nanotechnology of blood related medicine; Legal aspects of blood medicine; Historical perspectives. The manuscript management system is completely online and includes a very quick and fair peer-review system. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <http://www.dovepress.com/journal-of-blood-medicine-journal>

Mask Protection Against Lead Exposure and Its Correlation with Erythropoiesis in Automotive Body Painters at Ligu District, Semarang, Indonesia

ORIGINALITY REPORT

10%

SIMILARITY INDEX

9%

INTERNET SOURCES

5%

PUBLICATIONS

2%

STUDENT PAPERS

PRIMARY SOURCES

1	researcher.life Internet Source	2%
2	Heni Setyowati Esti Rahayu, N. Nasruddin, Laela Hayu Nurani, Sri Darmawati et al. "Ethanol extract of the natural product of Daun sirih (Piper betle) leaves may impede the effectiveness of the plasma jet contact style for acute wounds", Clinical Plasma Medicine, 2019 Publication	1%
3	WWW.DOVEPRESS.COM Internet Source	1%
4	www.science.gov Internet Source	1%
5	oamjms.eu Internet Source	1%
6	teknolabjournal.com Internet Source	1%

7	es.scribd.com Internet Source	1 %
8	Stenveld, Harma. "Allergic to Pool Water", Safety and Health at Work, 2012. Publication	<1 %
9	smujo.id Internet Source	<1 %
10	njafe.org Internet Source	<1 %
11	Submitted to De Montfort University Student Paper	<1 %
12	johe.rums.ac.ir Internet Source	<1 %
13	Homegnon A. F. Bah, Ana Laura S. dos Anjos, Erival A. Gomes-Júnior, Matheus J. Bandeira et al. "Delta-Aminolevulinic Acid Dehydratase, Low Blood Lead Levels, Social Factors, and Intellectual Function in an Afro-Brazilian Children Community", Biological Trace Element Research, 2021 Publication	<1 %
14	www.jjbs.hu.edu.jo Internet Source	<1 %
15	Masahiro Hashizume, Momoko Chiba, Atsuko Shinohara, Shigehiro Iwabuchi et al. "Anaemia, iron deficiency and vitamin A	<1 %

status among school-aged children in rural
Kazakhstan", Public Health Nutrition, 2007

Publication

16

R S Iswari, M Dafip, A I Kartika, I R Apriliana, I
N Chamidah, M Abduh. "Lipid profile
comparison of tomato extract and
atorvastatin supplementation in
atherosclerosis rats", Journal of Physics:
Conference Series, 2020

Publication

<1 %

17

occmmed.oxfordjournals.org

Internet Source

<1 %

18

www.mdpi.com

Internet Source

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On

Mask Protection Against Lead Exposure and Its Correlation with Erythropoiesis in Automotive Body Painters at Ligu District, Semarang, Indonesia

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7
