

Susceptibility of *Aedes albopictus* Larvae to the Larvicidal Activity of Three Types of *Derris elliptica* Extract

by Sayono Sayono

Submission date: 15-Mar-2022 09:47AM (UTC+0700)

Submission ID: 1784550762

File name: 263-269.pdf (610.98K)

Word count: 4423

Character count: 23286

<http://www.pjbs.org>

PJBS

ISSN 1028-8880

**Pakistan
Journal of Biological Sciences**

ANSI*net*

Asian Network for Scientific Information
308 Lasani Town, Sargodha Road, Faisalabad - Pakistan



Research Article

Susceptibility of *Aedes albopictus* Larvae to the Larvicidal Activity of Three Types of *Derris elliptica* Extract

¹Sayono Sayono, ²Risyandi Anwar, ³Didik Sumanto, ⁴Eni Nurmalasari and ⁵Fajar Fauzi Abdullah

¹Faculty of Public Health, Universitas Muhammadiyah Semarang, Semarang, Indonesia

²Herbal medicine research of Dentistry Faculty, Universitas Muhammadiyah Semarang, Semarang, Indonesia

³Laboratory of Epidemiology and Tropical Diseases, Faculty of Public Health, Universitas Muhammadiyah Semarang, Semarang, Indonesia

⁴Student of Public Health Program, Faculty of Public Health, Universitas Muhammadiyah Semarang, Semarang, Indonesia

⁵Laboratory of Natural Chemical of Sciences and Mathematics Faculty of Garut University, West Java Province, Indonesia

Abstract

Background and Objective: The methanol, ethyl acetate and n-hexane extracts of *D. elliptica* root have high larvicidal activity against *Aedes aegypti* larvae, the primary vector of dengue but have not been understood the potential against *Ae. albopictus* larvae, the secondary vector of dengue that also transmits Chikungunya and Zika viruses. This *in vitro* study aims to understand the larvicidal activity of the 3 extract types of *D. elliptica* root against *Ae. albopictus* larvae. **Materials and Methods:** The tuba root extract types were obtained from the sequential extraction process with 3 steps of liquid-liquid partition as described in the previous report. Six concentrations were occupied in this experiment ranging of 0.5, 1.0, 2.0, 4.0, 10.0 and 15.0 mg L⁻¹ each concentration was 5 times replicated and tested in 250 mL plastic cups. As many as 20 of 3rd instar larvae of *Ae. albopictus* were subjected in each treatment cup and larval mortality was observed after 24 and 48 hrs of exposure. **Results:** Larval mortality rates based on concentration range of 13.75-97.00 and 43.75-100%, 14.00-44.00, 34.00-90.00%, 12.00-47.00 and 28.00-88.00%, with the LC₅₀ after 24 and 48 hrs of exposure were 0.325 and 0.414, 16.184, 2.900, 15.789 and 4.380 mg L⁻¹, respectively for methanol, ethyl acetate and n-hexane extracts. **Conclusion:** The methanol, ethyl acetate and n-hexane extract of tuba root have high larvicidal activity against *Ae. albopictus* larvae. Further study on prototype formulation of larvicide and elucidation of the specific phytochemical compounds of the extracts were necessarily conducted.

Key words: *Aedes albopictus*, arboviruses vector, *Derris elliptica*, methanol extract, ethyl acetate extract, n-hexane extract, larvicidal activity

Citation: Sayono, S., R. Anwar, D. Sumanto, E. Nurmalasari and F.F. Abdullah, 2022. Susceptibility of *Aedes albopictus* larvae to the larvicidal activity of three types of *Derris elliptica* extract. Pak. J. Biol. Sci., 25: 263-269.

Corresponding Author: Sayono Sayono, Faculty of Public Health, Universitas Muhammadiyah Semarang, Semarang, Indonesia
Tel: +62-24-76740296-7 Fax: +62-24-76740291

Copyright: © 2022 Sayono Sayono *et al.* This is an open access article distributed under the terms of the creative commons attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

Competing Interest: The authors have declared that no competing interest exists.

Data Availability: All relevant data are within the paper and its supporting information files.

INTRODUCTION

The competence of *Ae. albopictus* mosquito-transmitted arboviruses such as dengue^{1,2}, Chikungunya³ and Zika⁴ has been reported in several countries and its ability in transmitting the other arboviral has been indicated⁵. The vectorial competence has triggered community attention in the arboviral impacted areas for implementing the control measures¹. Unfortunately, the preferred habitat for this species is placed with lush trees and far from human settlements such as cemeteries and beaches⁶. Globally, the area affected by dengue has expanded to 129 countries, mainly in Asia and the number of new cases has increased more than eightfold in 2 decades⁷. This has sparked community efforts to control its vectors, including *Ae. albopictus*.

The use of chemical methods in dengue vector control for decades has resulted in the emergence of *Aedes* mosquito strains that are resistant to several insecticide formulations, including the Temephos larvicide in Southeast Asia⁸. Several studies also proved that *Ae. albopictus* was resistant to the larvicide Temephos in Brazil⁹, Pakistan¹⁰, India, Malaysia, Sri Lanka, China and Central Africa¹¹. This phenomenon also occurs in Indonesia, including in Surabaya¹² and Bengkulu¹³ where *Ae. albopictus* is resistant to organophosphate insecticides (Temephos and Malathion). This resistance issue can hinder the success of arboviruses infection prevention efforts in affected areas. This is also exacerbated by other factors such as population mobility and high connectivity between rural and urban areas which can provide greater opportunities for arbovirus exposure by *Ae. albopictus* mosquitoes¹⁴. The emergence of Temephos resistant strain of *Ae. albopictus* can hinder efforts to control infectious diseases. This situation triggers researchers to develop alternative larvicides that are effective and environmentally friendly by exploring new active compounds¹⁵, including chemical compounds from natural ingredients.

Studies²² the larvicidal activity of various plant extracts have been carried out, especially for *Ae. aegypti* larvae. Previous studies resulted in ranking the effectiveness of plant extract larvicides, namely high, moderate, low and ineffective based on the lethal concentration 50% (LC₅₀) values <50, 50-100, 100-750 and higher than 750 mg L⁻¹¹⁶. *D. elliptica* is one of the local plants that has high larvicidal potential against *Ae. aegypti* larvae. Experiments with 3 types of plant extracts, namely methanol, ethyl acetate and n-hexane, showed a low effective concentration (LC₅₀), of 14.066, 21.063 and 4.086 mg L⁻¹, respectively¹⁷. In particular,

the results of the bioassay test for ethyl acetate of *Derris elliptica* extract also showed effective larvicidal activity even though it was exposed to larvae from *Ae. aegypti* mosquitoes that were resistant to cypermethrin 0.05%, with an LC₅₀ of 34.945 mg L⁻¹¹⁸. These results are interesting to apply to the 2nd vector of dengue, the *Ae. albopictus* mosquito.

This *in vitro* study was aimed to determine the larvicidal activity of methanol, ethyl acetate and n-hexane extracts of tubal roots against *Ae. albopictus* larvae.

19

MATERIALS AND METHODS

Study site: This study was carried out for 7 months from April-October, 2021 at 2 different laboratories, namely the Natural Chemical Laboratory of Sciences and Mathematics Faculty of Garut University, West Java Province for the extraction process and the Laboratory of Epidemiology and Tropical Diseases of Universitas Muhammadiyah Semarang, Central Java, Indonesia for mosquito collection, rearing and bioassay experiments.

Tuba root collection and processing: The origin, collection and processing of tubal roots were described as in the previous studies¹⁷. The extracts have been processed from April-June, 2021 delivered and stored in a refrigerator at 4-8 in the Laboratory of Epidemiology and Tropical Diseases of Universitas Muhammadiyah Semarang.

Mosquito collection and rearing: *Aedes albopictus* mosquito was obtained from larval surveys around the Muhammadiyah University Semarang campus, especially in breeding places far from human habitation, cemeteries and gardens with lush trees⁶. Mosquito larvae from the survey were reared into adult mosquitoes and subjected to morphological species identification. During rearing, the larvae are fed dog food. Breeding is continued until the 2nd generation of eggs is obtained. During breeding, mosquitoes were fed with a solution of 10% sugar and guinea pig blood and the environmental conditions were maintained at a temperature of 28±2 and humidity of 75±10%. The eggs of the 2nd offspring were bred into 3rd instar larvae and subjected to experiments as many as 20 larvae per treatment.

Experiments: The bioassay test was carried out in several stages. Preliminary tests were carried out with concentration ranges of 4, 25 and 40 mg L⁻¹ based on the previous study's LC₅₀ and LC₉₀¹⁷ and obtained larval mortality of 27, 87 and

98% respectively. Based on these results, a bioassay test was determined with a lower concentration range of 2, 4, 10 and 20 mg L⁻¹ and resulted in a larval mortality range of 24-97%. Lower concentration ranges were achieved in the 3rd stage of the bioassay test, namely 0.5, 1.0, 2.0, 4.0, 10.0 and 15.0 mg L⁻¹. Each concentration level was carried out in 5 replications. Experiments were compared with 2 control groups, namely Temephos 0.02 mg L⁻¹ as a positive control as well as a standard concentration, while the negative control was distilled water. The research subjects were 3rd instar *Ae. albopictus* larvae with active movement conditions. Twenty larvae were subject to each treatment and the larval mortality rate was observed at 24th and 48th hrs post-exposure.

Data analysis: Larval mortality data were analysed descriptively in the form of tables and graphs and analytically with probit and 2 way ANOVA tests to determine the effective concentration and significance of mortality based on the type and concentration of the extract. Data analysis using SPSS and Excel software.

Ethical approval: This study was obtained ethical approval from the Ethics Committee of Health Research of Public Health

Faculty of Universitas Muhammadiyah Semarang with registration number 231/KEPK-FKM/UNIMUS/2019.

RESULTS

Overall, the tuba root extracts showed high larvicidal activity against *Ae. albopictus* larvae. After 48 hrs of exposure, the results showed that the mortality rates based on the lowest to highest concentration, ranged from 43.75-100, 34-90 and 28-88%, respectively, for methanol, ethyl acetate and n-hexane extracts (Table 1).


Based on observation time of 24 hrs after exposure, only methanol extract showed high larvicidal potential with a mortality rate of 13.75-97.00%. However, the 3 types of extracts showed a high trend of increasing larvicidal activity based on observations 48 hrs after exposure (Fig. 1).

The results of statistical analysis showed significant differences in larval mortality based on extract type, concentration and interaction of extract type and concentration (Table 2). Based on the types of extracts, there were significant differences in larval mortality with the order of larvicidal activity from highest to lowest were methanol, ethyl acetate and n-hexane extracts (Table 3).

Table 1: Larval mortality of *Aedes albopictus* based on the types and concentrations of tuba root extract

Extract types	Concentration (ppm)	Post-exposure larval mortality (%)					
		24 hrs			48 hrs		
		Minimum	Maximum	Mean	Minimum	Maximum	Mean
Methanol	0.1	10.0	25.0	13.75	40.0	50.0	43.75
	0.5	15.0	25.0	16.25	65.0	70.0	66.25
	1.0	25.0	25.0	22.00	80.0	90.0	83.00
	2.0	25.0	30.0	24.00	85.0	90.0	86.00
	4.0	25.0	30.0	26.00	90.0	95.0	91.00
	10.0	40.2	86.0	83.00	85.0	100.0	98.00
Ethyl acetate	0.5	10.0	20.0	14.00	25.0	40.0	34.00
	1.0	15.0	20.0	15.00	30.0	55.0	40.00
	2.0	15.0	25.0	17.00	35.0	70.0	49.00
	4.0	15.0	30.0	21.00	55.0	85.0	66.00
	10.0	30.0	45.0	37.00	60.0	95.0	81.00
	15.0	35.0	50.0	44.00	85.0	95.0	90.00
n-hexane	0.5	10.0	15.0	12.00	25.0	35.0	28.00
	1.0	10.0	15.0	13.00	25.0	35.0	30.00
	2.0	15.0	15.0	15.00	35.0	60.0	47.00
	4.0	15.0	20.0	17.00	45.0	70.0	55.00
	10.0	25.0	50.0	33.00	50.0	90.0	78.00
	15.0	40.0	60.0	47.00	80.0	95.40	88.00
Temephos*	0.02	100	100	100	-	-	-
Aquadest†	0	0	0	0	0	0	0

*Positive control and †Negative control



1 **Fig. 1: Mortality rate of *Aedes albopictus* larvae after 24 and 48 hrs exposure to three extract types**
 Three extract types, namely MeOH: Methanol, EA: Ethyl acetate and Hex: n-hexane, MeOH extract type showed the rapid progress on mortality of research subject

Table 2: Effect of extract type, concentration and their interaction on larval mortality

Variables	F	p-value
Intercept	62.538	0.001
Extract types	37.662	0.000
Concentrations	31.564	0.000
Extract types and concentrations	2.360	0.018

Table 3: Multiple comparison of extract types on larval mortality

Extract types	Mean difference	p-value	Confidence interval (95%)
Methanol-ethyl acetate	23.83	0.000	19.10-28.57
Methanol-n-hexane	30.00	0.000	25.26-34.74
Ethyl acetate-n-hexane	6.17	0.011	1.43-10.90

6 **Table 4: Lethal concentration (LC₅₀ and LC₉₀) of the methanol, ethyl acetate and n-hexane extract types on mortality of *Aedes albopictus* larvae**

Extract types	Regression equations	Lethal concentration (ppm)		Chi-square	p-value
		LC ₅₀ (95% CI)	LC ₉₀ (95% CI)		
24 hrs exposure					
Methanol	Y = -1.915+0.655X	2.925 (2.641-3.200)	4.882 (4.487-5.423)	52.713	0.002
Ethyl acetate	Y = -1.108-0.068	16.184 (13.492-20.751)	34.899 (28.239-46.937)	12.013	0.999
n-hexane	Y = -1.246+0.079	15.789 (13.455-19.471)	32.022 (26.596-41.130)	10.400	0.999
48 hrs exposure					
Methanol	Y = -0.151+0.364X	0.414 (-11.872-1.863)	3.938 (2.291-40.334)	467.885	0.000
Ethyl acetate	Y = -0.351+0.121	2.900 (1.527-4.092)	13.473 (11.262-17.051)	49.079	0.011
n-hexane	Y = -0.540+0.123	4.380 (3.283-5.468)	14.767 (12.648-17.972)	40.834	0.071

4 **The final results of the bioassay test showed that the effective concentrations (LC₅₀ and LC₉₀) of each type of extract were 2.925 and 4.882, 16.184 and 34.899 and 15.789 and 32.022 mg L⁻¹ at 24 hrs observation and then 0.414 and 3.938, 2.900 and 13.473 and 4.380 and 14.767 mg L⁻¹ 48 hrs observation, respectively for methanol, ethyl acetate and n-hexane extracts (Table 4). The dead larvae were not found in the negative control and 100% mortality was found in**

the positive control groups. This finding indicated that the mortality of larvae in the treatment group was caused by the larvicidal activity of tuba root extracts.

DISCUSSION

In general, the results showed that the 3 types of tuba root extract indicated a high larvicidal activity, according to

the classification of larvicidal effectiveness of plant extracts that had been previously reported¹⁶. Larval mortality increased with increasing concentration¹⁵ and exposure time. The highest potency was shown by the methanol extract, both after 24 and 48 hrs of exposure, equivalent to the combined extract of petroleum-ether and methanol-chloroform¹⁹. From the aspect of cost, time and resources, this study is more efficient because it applies a sequential extraction which is carried out in a series of processes and with cheaper solvents, although both sequential and direct extraction have advantages and disadvantages²⁰.

The effective concentration in this study was lower than the exposure of the same extract to *Ae. aegypti* larvae in the previous study¹⁷. This indicates that the larvicidal activity or susceptibility of *Ae. albopictus* larvae are higher. *Ae. albopictus* larvae were more susceptible to the larvicidal activity of tuba root extract. This vulnerability can be attributed to the habitat preferences and flight ability of this species. The population of *Ae. albopictus* has a different habitat preference from *Ae. aegypti*, although co-occurrence often occurs²¹. The *Ae. albopictus* mosquito occupies habitats far from human settlements⁶ so it has a low chance of exposure to insecticides from dengue vector control programs, including the larvicide Temephos. The fact shows that there are fewer reports of monitoring *Ae. albopictus* resistance to insecticides from arboviruses control programs than *Ae. Aegypti*²², including in Indonesia^{12,13} which is only reported from a limited number of locations. This low history of exposure to insecticides causes the development of resistance mechanisms, both knockdown and lower metabolic rates⁸. Several studies have shown that this species is more dominant in rural and suburban areas than in urban areas²³ but this species is also dominant in urban and suburban environments in low-temperature areas²⁴, or in settlements where small breeding sites are found outdoors²¹. This low susceptibility is also supported by other factors, namely the flight distance of *Ae. albopictus* which reaches more than 200 m which allows it to avoid exposure to adulticides²⁵.

This study indicated that the methanol extract had a higher and faster larvicidal potential than the ethyl acetate and n-hexane extract types. Methanol is a solvent that can produce high extract products and phytochemical constituents, namely phenolics, alkaloids, flavonoids and terpenoids²⁶. Flavonoids are secondary metabolites that are widely found in tubal plants²⁷⁻²⁹. Flavonoids work by inhibiting the enzyme acetylcholinesterase by prolonging the effect of acetylcholine which increases nerve impulses at synapses³⁰, causing the larvae to spasm and die. Rotenone is one of the dominant flavonoids in *D. elliptica*. The activity of these

compounds affects electron transport or oxidative phosphorylation which inhibits cellular oxygen uptake so that energy production drops drastically. This situation triggers anaerobic cellular metabolism leading to increased lactic acid production and tissue acidosis and anoxia and death from heart and nervous system failure³¹.

The larvicidal activity of the n-hexane extract in this study was lower than previous findings¹⁷, with the order of highest to lowest larvicidal potency being n-hexane, methanol and ethyl acetate. In this study, the larvicidal activity of n-hexane extract was the lowest after methanol and ethyl acetate. This condition is thought to be due to the degradation of phytochemical compounds, especially polyphenols as reported by a study that proved that storage of fresh extracts for a period of 3-6 months caused the degradation of these compounds, except for methanol extracts which tended to be more stable³². Three types of tuba root extract used in this study were extracted 6 months ago and stored in a refrigerator at 4-8°C so that, this degradation factor is suspected to occur. Nevertheless, the larvicidal activity of this tubal root extract shows promising potential to be developed to the level of larvicidal prototype formulation and further exploration of the specific phytochemical compounds that play a role is also carried out.

CONCLUSION

Three types of tuba root extract, namely methanol, ethyl acetate and n-hexane, respectively have high larvicidal potential against the *Ae. albopictus* larvae. Further studies on the technical grade of larvicidal prototypes and elucidation of specific chemical compounds are necessary done.

SIGNIFICANCE STATEMENT

This study found high larvicidal activity in 3 types of a tuba root extract that can be beneficial for obtaining the specific chemical compounds as larvicide material for *Aedes* mosquitoes. This study will help the researchers to uncover critical areas of finding alternative methods for solving the resistance problems in mosquito vector control that many researchers are unable to explore. This finding reinforces that new theories on herbal chemical compounds can arrive in the near time.

ACKNOWLEDGMENT

We would like to thank the Deputy for Strengthening Research and Development, Ministry of Research and

Technology/National Research and Innovation Agency for funding this research with the grant contract number 1867/E4/AK.04/2021 and 067/E4.1/AK.04/PT2021.

REFERENCES

1. Kamgang, B., M. Vazeille, A.N. Tedjou, T.A. Wilson-Bahun and A.P. Yougang *et al.*, 2019. Risk of dengue in central Africa: Vector competence studies with *Aedes aegypti* and *Aedes albopictus* (Diptera: Culicidae) populations and dengue 2 virus. PLOS Neglected Trop. Dis., Vol. 13. 10.1371/journal.pntd.0007985.
2. Ferreira-de-Lima, V.H., P.D. Andrade, L.M. Thomazelli, M.T. Marrelli, P.R. Urbinatti, R.M.M. de Sá Almeida and T.N. Lima-Camara, 2020. Silent circulation of dengue virus in *Aedes albopictus* (Diptera: Culicidae) resulting from natural vertical transmission. Sci. Rep., Vol. 10. 10.1038/s41598-020-60870-1.
3. Vega-Rúa, A., M. Marconcini, Y. Madec, M. Manni and D. Carraretto *et al.*, 2020. Vector competence of *Aedes albopictus* populations for chikungunya virus is shaped by their demographic history. Commun. Biol., Vol. 3. 10.1038/s42003-020-1046-6.
4. McKenzie, B.A., A.E. Wilson and S. Zohdy, 2019. *Aedes albopictus* is a competent vector of zika virus: A meta-analysis. PLOS ONE, Vol. 14. 10.1371/journal.pone.0216794.
5. Pereira-dos-Santos, T., D. Roiz, R. Lourenço-de-Oliveira and C. Paupy, 2020. A systematic review: Is *Aedes albopictus* an efficient bridge vector for zoonotic arboviruses? Pathogens, Vol. 9. 10.3390/pathogens9040266.
6. Champion, S.R. and C.J. Vitek, 2014. *Aedes aegypti* and *Aedes albopictus* habitat preferences in south texas, USA. Environ. Health Insights, 8: 35-42. 10.4137/ehi.s16004.
7. Bhatt, S., P.W. Gething, O.J. Brady, J.P. Messina and A.W. Farlow *et al.*, 2013. The global distribution and burden of dengue. Nature, 496: 504-507.
8. Gan, S.J., Y.Q. Leong, M.F.H. bin Barhanuddin, S.T. Wong, S.F. Wong, J.W. Mak and R.B. Ahmad, 2021. Dengue fever and insecticide resistance in *Aedes* mosquitoes in southeast Asia: A review. Parasites Vectors, Vol. 14. 10.1186/s13071-021-04785-4.
9. Prophiro, J.S., O.S. Silva, J.E.D. Luna, C.F. Piccoli, L.A. Kanis and M.A.N. da Silva, 2011. *Aedes aegypti* and *Aedes albopictus* (Diptera: Culicidae): Coexistence and susceptibility to temephos, in municipalities with occurrence of dengue and differentiated characteristics of urbanization. Rev. Soc. Bras. Med. Trop., 44: 300-305.
10. Khan, H.A.A., 2020. Resistance to insecticides and synergism by enzyme inhibitors in *Aedes albopictus* from Punjab, Pakistan. Sci. Rep., Vol. 10. 10.1038/s41598-020-78226-0.
11. Bharati, M. and D. Saha, 2021. Insecticide resistance status and biochemical mechanisms involved in aedes mosquitoes: A scoping review. Asian Pac. J. Trop. Med., 14: 52-63.
12. Mulyatno, K.C, A. Yamanaka, Ngadino and E. Konishi, 2012. Resistance of *Aedes aegypti* (L.) larvae to temephos in Surabaya, Indonesia. Southeast Asian J. Trop. Med. Public Health, 43: 29-33.
13. Triana, D., S.R. Umniyati and B. Mulyaningsih, 2018. Resistance status of *Aedes albopictus* (skuse) on malathion in Bengkulu city. Unnes J. Public Health, 7: 113-119.
14. Kraemer, M.U., M.E. Sinka, K.A. Duda, A.Q. Mylne and F.M. Shearer *et al.*, 2015. The global distribution of the arbovirus vectors *Aedes aegypti* and *Ae. albopictus*. eLIFE, Vol. 4. 10.7554/eLife.08347.
15. Arnason, J.T., S.R. Sims and I.M. Scott, 2012. Natural products from plants as insecticides. In: Phytoc-hemistry and Pharmacognosy, Pezzuto, J.M. and M. Kato (Eds.). Oxford, UK.
16. Komalamisra, N., Y. Trongtokit, R. Rongsriyam and C. Apiwathnasorn, 2005. Screening for larvicidal activity in some Thai plants against four mosquito vector species. Southeast Asian J. Trop. Med. Public Health, 36: 1412-1422.
17. Sayono, S., R. Anwar and D. Sumanto, 2020. Evaluation of toxicity in four extract types of tuba root against dengue vector, *Aedes aegypti* (Diptera: Culicidae) larvae. Pak. J. Bio. Sci., 23: 1530-1538.
18. Sayono, S., R. Anwar and D. Sumanto, 2020. Larvicidal activity of ethyl acetate extract of *Derris elliptica* root against the third-instar larvae of cypermethrin-resistant *Aedes aegypti* offspring. J. Arthropod-Borne Dis., 14: 391-399.
19. Dohutia, C., D.R. Bhattacharyya, S.K. Sharma, P.K. Mohapatra and K. Bhattacharjee *et al.*, 2015. Larvicidal activity of few select indigenous plants of North East India against disease vector mosquitoes (Diptera: Culicidae). Trop. Biomed., 32: 17-23.
20. John, K.M.M., J. Harnly and D. Luthria, 2018. Influence of direct and sequential extraction methodology on metabolic profiling. J. Chromatogr. B, 1073: 34-42.
21. Djiappi-Tchamen, B., M.S. Nana-Ndjangwo, T. Tchuinkam, I. Makoudjou and E. Nchoutpouen *et al.*, 2021. Aedes mosquito distribution along a transect from rural to urban settings in yaoundé, Cameroon. Insects, Vol. 12. 10.3390/insects12090819.
22. Hassan, M.R., N.A. Azit, S.M. Fadzil, S.R.A. Ghani, N. Ahmad and A.M. Nawi, 2021. Insecticide resistance of dengue vectors in southeast Asia: A systematic review. Afr. Health Sci., 21: 1124-1140.
23. Lutinski, J.A., E.A. Bollis, A.B. Serena, D.M.M. Fritzen, M.A. Busato, W.A.R. Júnior and D.A. Simões, 2020. Aedes albopictus mosquito infestation in a municipality in the Western Region of the state of Santa Catarina. Saúde Pesqui., 13: 399-409.
24. Westby, K.M., S.A. Adalsteinsson, E.G. Biro, A.J. Beckermann and K.A. Medley, 2021. *Aedes albopictus* populations and larval habitat characteristics across the landscape: Significant differences exist between urban and rural land use types. Insects, Vol. 12. 10.3390/insects12030196.

25. Marini, F., B. Caputo, M. Pombi, M. Travaglio and F. Montarsi *et al.*, 2019. Estimating spatio-temporal dynamics of *Aedes albopictus* dispersal to guide control interventions in case of exotic arboviruses in temperate regions. *Sci. Rep.*, Vol. 9. 10.1038/s41598-019-46466-4.
26. Truong, D.H., D.H. Nguyen, N.T.A. Ta, A.V. Bui, T.H. Do and H.C. Nguyen, 2019. Evaluation of the use of different solvents for phytochemical constituents, antioxidants, and *in vitro* anti-inflammatory activities of *Severinia buxifolia*. *J. Food Qual.*, Vol. 2019. 10.1155/2019/8178294.
27. Wu, X., Z. Song, H. Xu, H. Zhang, W. Chen and H. Liu, 2012. Chemical constituents of the aerial part of *Derris elliptica*. *Fitoterapia*, 83: 723-736.
28. Lu, H.Y., L. Jing-Yu, Y. Ping and Q. Wei, 2008. Isoflavones and their derivatives from the root of *Derris elliptica* (Roxb.) Benth. *Chin. J. Nat. Med.*, 6: 265-267.
29. Russell, D.A., W.J.S. Fong, D.G. Twigg, H.F. Sore and D.R. Spring, 2017. Stereocontrolled semisyntheses of elliptone and 12 α -hydroxyelliptone. *J. Nat. Prod.*, 80: 2751-2755.
30. Ding, X., M.A. Ouyang, X. Liu and R.Z. Wang, 2013. Acetylcholinesterase inhibitory activities of flavonoids from the leaves of ginkgo biloba against brown planthopper. *J. Chem.*, Vol. 2013. 10.1155/2013/645086.
31. Rayner, T.S. and R.G. Creese, 2006. A review of rotenone use for the control of non indigenous fish in Australian fresh waters, and an attempted eradication of the noxious fish, *Phalloceros caudimaculatus*. *N. Z. J. Mar. Fresh water Res.*, 40: 477-486.
32. Materska, M., 2010. Evaluation of the lipophilicity and stability of phenolic compounds in herbal extract. *Acta Sci. Pol., Technol. Aliment.*, 9: 61-69.

Susceptibility of Aedes albopictus Larvae to the Larvicidal Activity of Three Types of Derris elliptica Extract

ORIGINALITY REPORT

13%

SIMILARITY INDEX

11%

INTERNET SOURCES

8%

PUBLICATIONS

3%

STUDENT PAPERS

PRIMARY SOURCES

1	publish.kne-publishing.com Internet Source	3%
2	Submitted to Universitas Riau Student Paper	2%
3	www.esciencecentral.org Internet Source	1%
4	smujo.id Internet Source	1%
5	Submitted to Surabaya University Student Paper	1%
6	www.ijpsonline.com Internet Source	1%
7	www.science.gov Internet Source	1%
8	pdfs.semanticscholar.org Internet Source	1%
9	jurnal.unimus.ac.id Internet Source	<1%

10

Whiting, D. C., L. E. Jamieson, and P. G. Connolly. "Effect of Sublethal Tebufenozide Applications on the Mortality Responses of *Epiphyas postvittana* (Lepidoptera: Tortricidae) Larvae Exposed to a High-Temperature Controlled Atmosphere", *Journal of Economic Entomology*, 1999.

Publication

<1 %

11

digilib.unimed.ac.id

Internet Source

<1 %

12

repository.istn.ac.id

Internet Source

<1 %

13

Aurelie P. Yougang, Basile Kamgang, Armel N. Tedjou, Theodel A. Wilson-Bahun, Flobert Njiokou, Charles S. Wondji. "Nationwide profiling of insecticide resistance in *Aedes albopictus* (Diptera: Culicidae) in Cameroon", *PLOS ONE*, 2020

Publication

<1 %

14

Abdul Khair, Noraida Noraida. "Young and Old Sugar Apple (*Annona squamosa* Linn) Leaf Extracts As an *Aedes aegypti* Larva Insecticide", *Medical Laboratory Technology Journal*, 2019

Publication

<1 %

15

D. Mathivanan, P. Rajiv Gandhi, R. Regina Mary, S.R. Suseem. "Larvicidal and acaricidal efficacy of different solvent extracts of *Andrographis echinoides* against blood-

<1 %

sucking parasites", Physiological and
Molecular Plant Pathology, 2018

Publication

16

Govindaraju Ramkumar, Sengodan Karthi,
Ranganathan Muthusamy, Ponnusamy
Suganya et al. "Mosquitocidal Effect of
Glycosmis pentaphylla Leaf Extracts against
Three Mosquito Species (Diptera:
Culicidae)", PLOS ONE, 2016

Publication

17

Nurul Jadid, Dewi Hidayati, Sylviana Rosyda
Hartanti, Byan Arasyi Arraniry, Rizka Yuanita
Rachman, Wiwi Wikanta. "Antioxidant
activities of different solvent extracts of
Piper retrofractum Vahl. using DPPH assay",
AIP Publishing, 2017

Publication

18

Sonia Ganassi, Marco Masi, Pasqualina
Grazioso, Antonio Evidente, Antonio De
Cristofaro. "Activity of Some Plant and
Fungal Metabolites Towards Aedes
Albopictus (Diptera, Culicidae)", Research
Square, 2020

Publication

19

journals.plos.org
Internet Source

<1 %

20

mail.scialert.net
Internet Source

<1 %

21

Abhisekh Subba, Minu Bharati, Priyanka Rai, Dhiraj Saha. "Phytochemical composition of *Heracleum nepalense* D. Don fruit extracts and its activity against the larvae of *Aedes albopictus* (Diptera: Culicidae)", *International Journal of Tropical Insect Science*, 2020

Publication

<1 %

22

Marimuthu Govindarajan. "Chapter 7 Green Synthesized Silver Nanoparticles: A Potential New Insecticide for Mosquito Control", Springer Science and Business Media LLC, 2016

Publication

<1 %

Exclude quotes Off

Exclude matches < 5 words

Exclude bibliography On

Susceptibility of Aedes albopictus Larvae to the Larvicidal Activity of Three Types of Derris elliptica Extract

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8
