

The Importance of Purification and Activity Analysis of the Purified Product of Thrombolytic Protease from Bacillus sp. HSFI-12– A Review

by Stalis Norma Ethica

Submission date: 03-Aug-2022 08:17AM (UTC-0400)

Submission ID: 1878413672

File name: 17._2022_Nurul_Islamiyah.pdf (987.99K)

Word count: 4384

Character count: 25871

The Importance of Purification and Activity Analysis of the Purified Product of Thrombolytic Protease from *Bacillus* sp. HSFI-12– A Review

Nurul Islamiyah¹ Stalis Norma Ethica^{1*} Muhammad Ardi Afriansyah² Ana Hidayati Mukaromah¹ Dewi Seswita Zilda³

¹Magister Study Program of Clinical Laboratory Science, Universitas Muhammadiyah Semarang, Jalan Kedungmundu Raya No. 18, Semarang, Central Java, Indonesia 50273

²Diploma Study Program of Medical Laboratory Technology, Faculty of Nursing and Health Sciences, Universitas Muhammadiyah Semarang, Jalan Kedungmundu Raya No. 18, Semarang, Central Java, Indonesia 50273

³Research Center for Marine and Fisheries Product Processing and Biotechnology, Jalan KS Tubun, Slipi, Jakarta, Indonesia 10260

*Corresponding author. Email: norma@unimus.ac.id

ABSTRACT

Mortality and morbidity of Cardio-vascular diseases (CVDs) have been the major issue in the group of non-communicable diseases worldwide. However, the existing antithrombotic drugs to combat CVD still have many shortcomings in terms of price and safety. *Bacillus* sp. HSFI-12 had been previously reported as a highly potential producer of thrombolytic protease, in which crude protease has competitive clot lysis or thrombolytic activity. However, the purity and other characteristics of the protease in its purified form have not been reported. The remained task is to ensure that *Bacillus* sp. HSFI-12, as a source of thrombolytic protease, can be used as an alternative of effective treatment with lower cost to combat thrombosis in cardiovascular diseases. This review aims to provide a highlight the most suitable purification methods to be applied to the protease dialysate of *Bacillus* sp. HSFI-12 and recommending appropriate assays following the recommended purification methods. Results of this review showed that both ion exchange and gel filtration chromatography (IEC and GFC) methods are suggested to be used in the purification step as a continuation of the dialysis process of HSFI-12 protease previously concentrated using ammonium sulfate. In addition, the results of the review found that the most common assays conducted on purified protease developed as antithrombotic agents are anticoagulation and antiplatelet assays. In conclusion, the purification and activity analysis of the purified product of thrombolytic protease from *Bacillus* sp. HSFI-12 using chromatographic methods is important to do. This review recommended that aside from anticoagulant and antiplatelet assays, thrombolytic activity tests on HSFI-12 protease dialysate are also important to do to support the protease's characteristics an antithrombotic agent.

Keywords: *Bacillus* sp. HSFI-12, Chromatography, Enzyme purification, Thrombosis, Thrombolytic protease.

1. INTRODUCTION

Cardiovascular disease (CVD) is among the primary causes of death worldwide triggered by thrombosis. Thrombosis is the formation of blood clots containing fibrin aggregates in the blood vessels causing block of blood flow. Such blockage could cause heart and brain

failures for not receiving an adequate oxygen supply [1,2]. The imbalance of the hemostasis system will result in pathological abnormalities that cause spontaneous bleeding because the blood cannot clot and the formation of thrombus due to excessive blockage. The presence of a thrombus will trigger vascular diseases including myocardial infarction, heart attack, brain embolism

(stroke), and various other vascular diseases. In addition, the thrombus also causes tears in the tissue and swelling of the arteries [3,4].

Therefore, treatment therapy is needed to overcome the abnormality using antithrombotic drugs such as anticoagulants and antiplatelets. Blood clots can be treated with antithrombotic drugs and can be eliminated through a fibrinolysis process involving hydrolysis by plasmin. However, the existing antithrombotic drugs still have many shortcomings in terms of high prices and serious side effects, so new drugs, which are more affordable, and safe are needed [5].

Due to the high rate of cardiovascular disease morbidity and mortality, special attention is required to develop more cost-effective therapies. *Bacillus* sp. has been widely recognized for its activity as a thrombolytic and antiplatelet agent and is a potential producer of fibrinolytic proteases. A member of the bacterial genus, namely *Bacillus* sp. HSFI-12 has been reported to be able to produce crude extracts of protease enzymes with the thrombolytic activity that competes with the antithrombotic drug Nattokinase. The isolate was originated from "Rusip" fermented product of the intestine organs of sea cucumber, *H. scabra*, collected from seawater of Kodek Bay, West Nusa Tenggara Island, Indonesia [6-7].

Bacillus sp. HSFI-12 was recognized in a previous study as a highly potential producer of thrombolysis protease. Crude protease of HSFI-12 isolated was found to have competitive clot lysis activity [6]. However, the purity of the protease has not been reported. The remained task is to ensure that *Bacillus* sp. HSFI-12, as a source of thrombolytic protease, can be used as an alternative of effective treatment with lower cost to combat thrombosis in cardiovascular diseases. However, the purity and other characteristics of the protease in its purified form have not been reported.

To develop the dialysate extract of the thrombolytic protease previously obtained from HSFI-12 strains, purification should be conducted and the product should be purified and further tested using *in vitro* and *in vivo* approaches. Without purification the antithrombotic activity of the enzyme could not be accurately determined. This article aims to provide a literature review on the importance of purifying the crude extract of the protease enzyme in *Bacillus* sp. HSFI-12 and recommending appropriate purification methods to further enhance its proteolytic activity.

2. METHODOLOGY

A literature search was first carried out from the literature databases such as PubMed, Science Direct, Proquest and Google Scholar. The screened articles were those published between 2011-2021 on the role of thrombolysis and its mechanisms in cardiovascular

disease. In particular, the search was advanced on bacterial enzymes developed as antithrombotic agents involving purification methods. Searching for subject titles in several combinations including "cardiovascular disease", "cardiovascular drugs and therapy", fibrinolytic proteases, "antithrombotic proteases", "bacterial protease", "fibrinolysis mechanism", "antithrombotic agent", "purification method", purified enzyme test", "antithrombotic activity" and their derivatives.

2.1. Criteria set for Relevant Studies

Study screening was based on inclusion criteria set as follows: [i] Subjects associated with the purification of thrombolytic protease enzymes in cardiovascular treatment; [ii] antithrombotic enzymes produced by bacteria [iii] enzyme purification studies and antithrombotic assays as a measure in determining pure enzyme characteristics; [iv] reported in Bahasa Indonesian or English; [v] observational studies as a search plan with the age of publication set to the last 10 years.

2.2. Study Selection

Following guidelines from Polanin et al. (2019), independent studies were conducted to identify studies, which met the inclusion criteria applied in this review. Both titles and abstracts from the records generated by the search were carefully identified and analyzed to determine which unsuitable sources should be excluded adhering to the exclusion criteria. Articles (full-text) of the remaining studies were also taken and evaluated whether or not they meet the inclusion criteria [8].

2.3. Bias Control Assessment

In this review, the following issues are included in the risk of tendency or quality assessment: completeness of reporting facts on the urgency of purification of thrombolytic bacterial proteases, choice of outcome measures (practical test of the capacity of thrombotic agents applied in cardiovascular treatment). When all of these criteria are fulfilled, the risk of possible bias is regarded low [9-10].

3. RESULT

This review summarized the purification methods and types and sources of bacterial proteases reported in the development of antithrombosis agents that have been reported in the last 5 years (during 2016-2020). The aim was to determine the most suitable purification methods to be applied to the dialysate product of protease originated from *Bacillus* sp. HSFI-12 to obtain the purified form of the enzyme with better antithrombotic characteristics supporting its role as a candidate of the antithrombotic agent.

Table 1. Chromatographic methods used to purify bacterial proteases developed as antithrombotic agents reported by various studies worldwide in the last 5 years

Bacterial species	Chromatography type of method	Country	References
<i>Bacillus tequilensis</i>	Ion Exchange Size Exclusion	China	Xin <i>et al.</i> , 2018 [11]
<i>Bacillus cereus</i>	DEAE-Cellulose Ion Exchange Gel Filtration	India	BKM <i>et al.</i> , 2018 [12]
<i>Bacillus</i> sp.	DEAE-Cellulose Ion Exchange Gel Filtration	Malaysia	Jimat <i>et al.</i> , 2017 [13]
<i>Triticum aestivum</i>	Ion Exchange Gel Filtration	Turkey	Altin <i>et al.</i> , 2017 [1]
<i>Bacillus cereus</i>	Ion Exchange	Indonesia	Junaidia <i>et al.</i> , 2017 [2]
<i>Bacillus</i> sp.	DEAE-Cellulose Ion Exchange	China	Yu <i>et al.</i> , 2019 [3]
<i>Geobacillus toebii</i>	DEAE-Cellulose Ion Exchange Gel Filtration	Tunisia	W <i>et al.</i> , 2016 [4]
<i>Bacillus</i> sp.	Column (unspecified)	India	Putatunda <i>et al.</i> , 2019 [18]
<i>Bacillus subtilis</i>	Ammonium Sulphate Precipitation DEAE-Cellulose Anion Exchange	Turkey	Nur <i>et al.</i> , 2019 [19]
<i>Bacillus subtilis</i>	Ammonium Sulphate Precipitation DEAE-Cellulose and sephadex G-100 Ion Exchange	India	Sujatha A <i>et al.</i> , 2018 [20]
<i>Geobacillus</i> sp.	Ammonium Sulphate Precipitation Ion Exchange	Indonesia	Iqbalyah <i>et al.</i> , 2019 [21]
<i>Bacillus subtilis</i>	Ammonium Sulphate Precipitation Ion Exchange Gel Filtration	Pakistan	M <i>et al.</i> , 2018 [5]
<i>Bacillus amyloliquefaciens</i>	DEAE-Cellulose Anion Exchange Gel Filtration	China	Hu <i>et al.</i> , 2020 [6]
<i>Bacillus subtilis</i>	Column Gel Filtration High Performance Liquid	China	Hue <i>et al.</i> , 2019 [24]

Table 1. Summarizes the chromatographic types of methods used to purify bacterial proteases developed as antithrombotic agents reported in various studies worldwide in the last 5 years (2016-2020). In Table 1 the trend of data displays that the chromatographic based purification methods widely used are dominated by Ion Exchange Chromatography (IEC) and Gel Filtration Chromatography (GFC). In Table 1, the trend of data also reveals that all of the identified studies reporting the purification process on antithrombotic bacterial proteases are dominated by Asian Countries Including China, Japan, India, and Indonesia. Those from Europe are only represented by one country (Turkey). Interestingly,

proteases subjected to the purification process are those mainly from *Bacillus* sp. [11-24].

Supporting the finding based on Table 1, there are several reasons why IEC is often selected in the purification process of antithrombotic proteases (usually done after ammonium deposition and dialysis process). First, IEC could separate ionizable molecules based on charge property differences. Its strong resolving capability, wide applicability (especially for proteins and enzymes), large sample handling capacity, moderate cost, and ease of scaling and automation have made it among the most widely used and versatile liquid chromatography methods found today [25-27]. On the

Table 2. Studies in the last 5 years reporting *in vitro* antithrombotic assays and associated instrumentation on bacterial enzymes following purification process using chromatographic methods

No	Antithrombotic assays	Instruments	References
1	<i>In vitro</i> Anticoagulant Activity (PT, aPTT, TT)	Automated Coagulation Analyzer Diagnostica Stago STart Hemostasis Analyzer	²³ Frías <i>et al.</i> , 2021 [7]
2	<i>In vitro</i> Anticoagulant Activity (PT, aPTT)	Automated Coagulation Analyzer Sysmex	Sukati <i>et al.</i> , 2021 [8]
3	Antiplatelet Activity (Platelet Adhesion & Platelet Aggregation)	Collagen Type IV Coated Strip	¹³ Sukati <i>et al.</i> , 2021 [8]
4	<i>In vitro</i> Anticoagulant Activity (PT, aPTT, TT)	Automated Coagulation Analyzer Diagnostica Stago STart Hemostasis Analyzer	Bougetef <i>et al.</i> , 2020 [9]
5	Thrombin Clotting Time (TCT)	ACL 200 Coagulation Analyzer	Pepe <i>et al.</i> , 2016 [10]
6	<i>In vitro</i> Antiplatelet Activity	Spektrofotometer UV Vis	¹⁷ Rohmah <i>et al.</i> , 2020 [11]
7	<i>In vitro</i> Anticoagulant Activity (CT, PT, aPTT)	Lee-White	Rohmah <i>et al.</i> , 2020 [11]
8	<i>In vitro</i> Anticoagulant Activity, (PT, aPTT)	Siemens Fibrintimer II Optical Coagulometer and Thromborel Kits	Bijak <i>et al.</i> , 2019 [12]
9	<i>In vitro</i> Anticoagulant Activity (PT, aPTT)	Commercial Kits	Gogoi <i>et al.</i> , 2019 [13]
10	<i>In vitro</i> Anticoagulant Activity (PT, aPTT, TT) and FB	Automated Coagulometer Sysmex CA-1500	¹⁹ Choi <i>et al.</i> , 2013 [14]
11	<i>In vitro</i> Anticoagulant Activity (aPTT dan PT)	Sysmex CA-50 Semi-automated Coagulation	Wang <i>et al.</i> , 2018 [15]
12	<i>In vitro</i> Anticoagulant Activity (PT, aPTT, TT)	Diagnostoc Stago	Krichen <i>et al.</i> , 2018 [16]
13	<i>In vitro</i> Anticoagulant Activity (PT, aPTT, TT)	Humacdot duo coagulometer and human reagent kits	¹² Rehman <i>et al.</i> , 2019 [17]
14	Platelet Aggregation	Optical lumi-aggregation system	Sachin <i>et al.</i> , 2021 [18]
15	<i>In vitro</i> Anticoagulant Activity	Lee-White and Blood Smear Eustrek	²² Fuad <i>et al.</i> , 2021 [42]
16	<i>In vitro</i> Anticoagulant Activity	Lee -White	Krishnamurthy <i>et al.</i> , 2017 [43]
17	<i>In vitro</i> Anticoagulant Activity	Lee-White and Blood Smear Eustrek	Rusyiana <i>et al.</i> , 2021 [44]
18	<i>In vitro</i> Anticoagulant Activity	Lee-White	Tian <i>et al.</i> , 2015 [22]
19	<i>In vitro</i> Anticoagulant Activity	Lee-White	Mahajan <i>et al.</i> , 2012 [23]

other hand, GFC is also preferred because it can perform the separation of different molecules with high yields, and is carried out with a special design to keep the activity and stability of the desired molecule without reducing its resolution [28]. Both IEC and GFC are not only the most widely reported in the literature, yet most suitable to be used to increase the activity of a proteolytic enzyme in general [16-19].

Table 2. Summarizes antithrombotic assays and used instrumentation on bacterial enzymes following purification using chromatography methods process [29-39]. Data in table 2 interestingly shows that regardless of the widely varied instrumentation used, types of antithrombotic assays are grouped into 3 main categories: *In vitro* anticoagulant activity and antiplatelet activity assays. *In vitro* Anticoagulant Activity assays include PT




Figure 1 Summarized factors contributing to the importance of purification and activity analysis of the purified product of thrombolytic protease from *Bacillus* sp. HSFI-12

7
(Prothrombin Time), aPTT (Activated Partial Thromboplastin Time), TT (Thrombin Time). Meanwhile Platelet Activities assays encompasses Platelet Adhesion and Platelet Aggregation. These *in vitro* aPTT, PT, and TT tests calculate the time elapsed from activation of the coagulation cascade [40].

It is known that antithrombotic characteristics encompass anticoagulant, antiplatelet and thrombolysis activities [41]. So, to develop an antithrombotic agent from proteases, it is also important to conduct thrombolytic assays on the purified product of the enzyme aside from anticoagulation and antiplatelet activities. This should also be applied to thrombolytic protease produced by isolate HSFI-12 if the aim is to develop it as an antithrombotic agent.

The mortality and morbidity of CVD as well as the death risk of thrombosis in CVD has motivated many researchers to search for better antithrombotic agents in terms of price, availability and side effects. Figure 1 summarizes factors contributing to the importance of purification and activity analysis of the purified product of thrombolytic protease from *Bacillus* sp. HSFI-12.

As schemed in Figure 1, the development of an antithrombotic agent using protease from *Bacillus* sp HSFI-12 offers a possible novelty in cardiovascular medicine therapy. The purification and activity analysis of the purified product of thrombolytic protease from *Bacillus* sp. HSFI-12 using IEC and GFC has not been reported, thus they are suggested to do to enhance the enzyme's activities. Anticoagulant and antiplatelet

activity assays on the same product have also not been reported, yet they are mostly reported by successful studies worldwide in the last 5 years. Thus, they are also required to do. In addition to this, however, to support the protease's characteristics as an antithrombotic agent, thrombolytic activity tests on the purified HSFI-12 protease dialysate should be done.

Both ion exchange and gel filtration chromatography (IEC and GFC) methods are suggested to be used in the purification step as a continuation of the concentration process of HSFI-12 protease dialysate as obtained in the previous study. In addition, this study suggested the importance of thrombolytic assays aside from anticoagulant and antiplatelet activity assays, which are often reported, to fully support the characteristics of HSFI-12 protease dialysate if developed as an antithrombotic agent.

AUTHORS' CONTRIBUTIONS

NI and SNE conceived the original idea, NI and MAA screened and summarized all obtained literatures. AHM evaluated the generation of tables and schemes, as well as analysed the bias of the study. The main text was written by NI and SNE. The manuscript was initially written by NI, and the improved and revised by SNE.

ACKNOWLEDGMENTS

The authors thanked Magister of Clinical Laboratory Science, Postgraduate Program, Universitas

Muhammadiyah Semarang for facilitating and supporting this study.

REFERENCES

- [1] C. Sharma, G. Elsayed, M. Salem, N. Sharma, P. Gautam, R. Singh, Thrombolytic Potential of Novel Thiol-Dependent Fibrinolytic Protease from *Bacillus cereus* RSA1, *Biomolecules*, 2020, pp. 3. DOI:10.3390/biom10010003
- [2] F. Altaf, S. Wu, V. Kasim, Role of Fibrinolytic Enzymes in Anti-Thrombosis Therapy, *Front Mol Biosci.*, 2021, pp. 476. DOI: 10.3389/FMOLB.2021.680397
- [3] H. Bougategf, Z. Ghilissi, R. Kallel, *et al.*, Chondroitin/dermatan sulfate purified from corb (*Sciaena umbra*) skin and bone: *In vivo* assessment of anticoagulant activity, *Int J Biol Macromol*, 2020, pp. 131-139. DOI: 10.1016/J.IJBIOMAC.2020.07.09
- [4] R. Setyowati, Uji Activity Antiplatelet dan Trombolitik Ekstrak Etanol Kulit Buah Jeruk Purut (*Citrus hystrix* D.C.) *In Vitro* [Antiplatelet and Thrombolytic Activity Test of Ethanol Extract of Kaffir Citrus Fruit Peel (*Citrus hystrix* D.C.) *In Vitro* [internet], 2021. Available from: <http://repository.unej.ac.id/handle/123456789/67640>
- [5] D. Dhamodharan, Novel fibrinolytic protease producing *Streptomyces radiopugnans* VITSD8 from marine sponges, *Marine drugs*, 2019, pp.164. DOI: <https://doi.org/10.3390/md17030164>
- [6] N. Hidayati, H. Fuad, H. Munandar, D.S. Zilda, N. Nurrahman, M. Fattah, O. Oedjijono, A. Samiasih, S.N. Ethica, Proteolytic and Clot Lysis Activity Screening of Crude Proteases Extracted from Tissues and Bacterial Isolates of *Holothuria scabra*. In IOP Conference Series: Earth and Environmental Science, 2021, pp. 012016. DOI: <http://dx.doi.org/https://doi.org/10.1088/1755-1315/755/1/012016>
- [7] N. Hidayati, N. Nurrahman, H. Fuad, H. Munandar, D.S. Zilda, A.R. Ernanto, A. Samiasih, O. Oedjijono, S.N. Ethica, *Bacillus tequilensis* Isolated from Fermented Intestine of *Holothuria scabra* Produces Fibrinolytic Protease with Thrombolysis Activity, In IOP Conference Series: Earth and Environmental Science, 2021, pp. 012008. <https://doi.org/10.1088/1755-1315/707/1/012008>
- [8] J.R. Polanin, T.D. Pigott, D.L. Espelage, J.K. Grotpeter, Best practice guidelines for Abstract screening large-evidence systematic reviews and meta-analyses, *Research Synthesis Methods*, 2019, pp. 330-42. DOI: <https://doi.org/10.1002/jrsm.1354>
- [9] S.N. Ethica, D.S. Zilda, O. Oedjijono, W. Nurgayah, M. Muhtadi, Bioprospection of alginate lyase from bacteria associated with brown algae *Hydroclathrus* sp. as antibiofilm agent: a review, *Aquaculture, Aquarium, Conservation & Legislation*, 2021, pp. 1974-89. DOI: <http://www.bioflux.com.ro/docs/2021.1974-1989.pdf>
- [10] S.N. Ethica, R. Saptaningtyas, S.I. Muchlissin, A. Sabdono, The development method of bioremediation of hospital biomedical waste using hydrolytic bacteria, *Health and Technology*, 2018, pp. 239-54. DOI: <https://doi.org/10.1007/s12553-018-0232-8>
- [11] X. Xin, R.R. Ambati, Z. Cai, B. Lei, Purification and characterization of fibrinolytic enzyme from a bacterium isolated from soil, *Biotech*, vol. 3, 2018. DOI:10.1007/S13205-018-1115-4
- [12] B.K. Lakshmi, D.M. Kumar, K.P. Hemalatha, Purification and characterization of alkaline protease with novel properties from *Bacillus cereus* strain S8, *Journal of Genetic Engineering and Biotechnology*, 2018, pp. 295-304. DOI: <https://doi.org/10.1016/j.jgeb.2018.05.009>
- [13] D.N. Jimat, I.B.F. Mohamed, A.S. Azmi, P. Jamal, Purification and Partial Characterization Of L-Asparaginase Enzyme Produced By Newly Isolate *Bacillus* sp. *Iium Eng J.*, 2017, pp. 1-10. DOI:10.31436/IJUMJE.V18I2.654
- [14] S. Altun, H. Tohma, İ. Gülçin, E. Köksal, Purification, characterization, and inhibition sensitivity of peroxidase from wheat (*Triticum aestivum* ssp. vulgare), *International Journal of Food Properties*, 2017, pp. 1949-59. DOI:10.1080/10942912.2016.1225308
- [15] Y. Junaidia, A. Pertiwinigruma, Y. Erwanto, N.A. Fitriyanto, Semi Purification and Identifications Molecule Protein Weigh of Alkaline Protease Enzyme from *Bacillus cereus* LS2B, *Int J Bio-Science Bio-Technology*, 2017, pp. 89-100. DOI:10.14257/ibsbt.2017.9.3.08
- [16] P. Yu, X. Huang, Q. Ren, X. Wang, Purification and characterization of a H₂O₂-tolerant alkaline protease from *Bacillus* sp. ZJ1502, a newly isolated strain from fermented bean curd, *Food Chem.*, 2019, pp. 510-517. DOI: 10.1016/J.FoodChem.2018.09.013
- [17] W. Thebti, Y. Riahi, O. Belhadj, Purification and characterization of a new thermostable,

- haloalkaline, solvent stable, and detergent compatible serine protease from *Geobacillus toebii* strain LBT 77, BioMed Research International, 2016. DOI: <https://doi.org/10.1155/2016/9178962>
- [18] C. Putatunda, B.S. Kundu, R. Bhatia, Purification and Characterization of Alkaline Protease from *Bacillus* sp. HD292, Proc Natl Acad Sci India Sect B - Biol Sci., 2019, pp. 957-965. DOI:10.1007/S40011-018-1011-Z
- [19] M. Nur, S. Karaboga, E. Logoglu, Purification of Alkaline Serine Protease From Local *Bacillus subtilis* M33 by Two Steps: a Novel Organic Solvent and Detergent Tolerant Enzyme [internet], 2019, pp. 116-129. Available from: <http://dergipark.gov.tr/gujs>
- [20] A. Sujatha, A. Subash, Isolation and Purification Of An Extracellular Protease From *Bacillus subtilis* Asasbt Isolated from Termite Soil, Int J Pharm Sci Res., 2018, pp. 4812-4819. DOI:10.13040/IJPSR.0975-8232.9
- [21] T.M. Iqbalsyah, Malahayati, Atikah, Febriani, Purification and partial characterization of a thermo-halostable protease produced by *Geobacillus* sp. strain PLS A isolated from undersea fumaroles, 2019. DOI:<https://doi.org/10.1080/16583655.2019.1650489>
- [22] M. Iqbal, M. Asgher, F. Bashir, Purification and kinetic characterization of alkaline protease for UV-90 mutant of *Bacillus subtilis*, J. Biochem. Anal. Stud., 2018. DOI: <http://dx.doi.org/10.16966/2576-5833.112>
- [23] H. Yang, L. Yang, X. Li, H. Li, Z. Tu, X. Wang, Genome sequencing, purification, and biochemical characterization of a strongly fibrinolytic enzyme from *Bacillus amyloliquefaciens* Jxnuwx-1 isolated from Chinese traditional Douchi, J. Gen Appl Microbiol., 2020, pp. 153-162. DOI: 10.2323/JGAM.2019.04.005
- [24] Y. Hu, D. Yu, Z. Wang, *et al.*, Purification and characterization of a novel, highly potent fibrinolytic enzyme from *Bacillus subtilis* DC27 screened from Douchi, a traditional Chinese fermented soybean food, Sci Reports, 2019, pp. 1-10. DOI:10.1038/s41598-019-45686-y
- [25] S. Mukherjee, Isolation and purification of industrial enzymes: advances in enzyme technology, In Advances in Enzyme Technology, 2019, pp. 41-70. DOI: <https://doi.org/10.1016/B978-0-444-64114-4.00002-9>
- [26] F.J. Contesini, R.R. Melo, H.H. Sato, An overview of *Bacillus proteases*: from production to application, 2017, pp. 321-334. DOI: <https://doi.org/10.1080/073885512017135435>.
- [27] Z. Che, X. Cao, G. Chen, Z. Liang, An effective combination of codon optimization, gene dosage, and process optimization for high-level production of fibrinolytic enzyme in *Komagataella phaffii* (*Pichia pastoris*), BMC Biotechnol., 2020, pp 1-13. DOI:10.1186/S12896-020-00654-7
- [28] C. Ó'Fágáin, P.M. Cummins, B.F. O'Connor, Gel-Filtration Chromatography. Protein Chromatogr., 2017, pp. 15. DOI:10.1007/978-1-4939-6412-3_2
- [29] J. Frias, D. Toubarro, A. Fraga, *et al.*, Purification and Characterization of a Thrombolytic Enzyme Produced by a New Strain of *Bacillus subtilis*, J Microbiol Biotechnol., 2021, pp. 327-337. DOI:10.4014/JMB.2008.08010
- [30] S. Sukati, W. Khobjai, *In vitro* antiplatelet and anticoagulant activity of indigenous vegetables from Southern Thailand, Int J Appl Pharm., 2021, pp. 38-42. DOI:10.22159/ijap.2021.v13s1.Y0100
- [31] H. Bougatef, Z. Ghilissi, R. Kallel, *et al.*, Chondroitin/dermatan sulfate purified from corb (*Sciaena umbra*) skin and bone: In vivo assessment of anticoagulant activity, Int J Biol Macromol, 2020, pp. 131-139. DOI: 10.1016/J.IJBIOMAC.2020.07.096
- [32] A. Pepe, E. Frey, F. Mu ~ Noz, *et al.*, Fibrin(ogen)olytic and antiplatelet activities of a subtilisin-like protease from *Solanum tuberosum* (StSBTc-3), 2016, pp. 165-170. DOI:10.1016/j.biochi.2016.03.015
- [33] M. Bijak, A. Sut, A. Kosiorek, J. Saluk-Bijak, J. Golanski, Dual Anticoagulant/Antiplatelet Activity of Polyphenolic Grape Seeds Extract, Nutr, 2019, pp. 93. DOI:10.3390/NU11010093
- [34] D. Gogoi, S. Ramani, S. Bhartari, P. Chattopadhyay, A.K. Mukherjee, Characterization of active anticoagulant fraction and a fibrin(ogen)olytic serine protease from leaves of *Clerodendrum colebrookianum*, a traditional ethno-medicinal plant used to reduce hypertension, J. Ethnopharmacol, 2019, pp. 112099. DOI:10.1016/J.JEP.2019.112099
- [35] J.H. Choi, K. Sapkota, S.E. Park, S. Kim, S.J. Kim, Thrombolytic, anticoagulant and antiplatelet activities of codiase, a bi-functional fibrinolytic enzyme from *Codium fragile*, Biochimie., 2013, pp. 1266-1277. DOI:10.1016/J.BIOCHI.2013.01.023
- [36] K.H. Wang, S.F. Li, Y. Zhao, H.X. Li, L.W. Zhang, *In Vitro* Anticoagulant Activity and Active

- Components of Safflower Injection, *Mol.*, 2018, pp. 170. DOI:10.3390/Molecules23010170
- [37] F. Krichen, H. Bougateg, F. Capitani F, *et al.*, Purification and structural elucidation of chondroitin sulfate/ dermatan sulfate from Atlantic bluefin tuna (*Thunnus thynnus*) skins and their anticoagulant and ACE inhibitory activities, *RSC Adv.*, 2018, pp. 37965-37975. DOI: 10.1039/C8RA06704J
- [38] A.A. Rehman, A. Riaz, M.A. Asghar, M.L. Raza, S. Ahmed, K. Khan, In vivo assessment of anticoagulant and antiplatelet effects of *Syzygium cumini* leaves extract in rabbits, *BMC Complement Altern Med.*, 2019, pp. 236. DOI:10.1186/S12906-019-2661-Y
- [39] H.R. Sachin, M.N. Sharath Kumar, S. Devaraja, A.H. Sneharani, Anticoagulant and antiplatelet activities of novel serine protease purified from seeds of *Cucumis maderaspatensis*, *Biotech.*, vol. 3, 2021, pp. 1-11. DOI:10.1007/s13205-020-02565-y
- [40] M.N. Raber, Coagulation Tests. In: Walker HK, Hall WD, Hurst JW, editors. *Clinical Methods: The History, Physical, and Laboratory Examinations*. 3rd edition, Boston, Butterworths, 1990, Chapter 157, Available from: <https://www.ncbi.nlm.nih.gov/books/NBK265/>
- [41] K. Manna, M. Das, Briefing Therapeutic Approaches in Anticoagulant, Thrombolytic, and Antiplatelet Therapy, In *Medicinal Chemistry with Pharmaceutical Product Development*, 2019, pp. 253-292.
- [42] H. Fuad, N. Hidayati, S. Darmawati, H. Munandar, A.R. Sulistyanningtyas, A.R. Ernanto, S.I. Muchlissin, D.S. Zilda, N. Nurrahman, S.N. Ethica, Exploration of bacteria isolated from "rusip" fermented tissue of sand sea cucumber *Holothuria scabra* with fibrinolytic, anticoagulant and antiplatelet activities, *AACL Bioflux*, 2021, pp. 1242-58. Available from: <http://www.bioflux.com.ro/docs/2021.1242-1258.pdf>
- [43] A. Krishnamurthy, P.D. Belur, P. Rai, P.D. Rekha, Production of Fibrinolytic Enzyme by the Marine Isolate *Serratia marcescens* subsp. *sakuensis* and its *In-vitro* Anticoagulant and Thrombolytic Potential, *Pure Appl. Microbiol.*, 2017, pp. 1987-98. DOI: 10.22207/JPAM.114.41.
- [44] R. Rusyiana, I.A. Lestari, C.D. Hamdin, H. Muliasari, Anticoagulant Activity of Mangrove (*Avicennia alba*) Leaves Extract *in Vitro*, *Indonesian Journal of Marine Sciences/Ilmu Kelautan*, 2021. Available from: <https://ejournal.undip.ac.id/index.php/ijms/article/view/35990/pdf>
- [45] Z. Tian, B. Li, L. Guo, M. Wu, T. Fu, H. Cheng, H. Zhu, Purification and biochemical characterization of a novel fibrinolytic enzyme, PSLTro01, from a medicinal animal *Porcellio scaber* Latreille, *International Journal of Biological Macromolecules*, 2015, pp. 53646. DOI:10.1016/j.ijbiomac.2015.06.046.
- [46] P.M. Mahajan, S. Nayak, S.S. Lele, Fibrinolytic enzyme from newly isolated marine bacterium *Bacillus subtilis* ICTF-1: Media optimization, purification and characterization, *Journal of bioscience and bioengineering*, 2012, pp. 307-14. DOI:10.1016/j.jbiosc.2011.10.023

The Importance of Purification and Activity Analysis of the Purified Product of Thrombolytic Protease from Bacillus sp. HSFI-12– A Review

ORIGINALITY REPORT

11%

SIMILARITY INDEX

8%

INTERNET SOURCES

9%

PUBLICATIONS

2%

STUDENT PAPERS

PRIMARY SOURCES

- | | | |
|---|--|----|
| 1 | www.atlantispress.com
Internet Source | 1% |
| 2 | Eka Sakti Wahyuningtyas, Arya Iswara, Yunita Sari, Sodik Kamal et al. "Comparative study on Manuka and Indonesian honeys to support the application of plasma jet during proliferative phase on wound healing", Clinical Plasma Medicine, 2018
Publication | 1% |
| 3 | pesquisa.bvsalud.org
Internet Source | 1% |
| 4 | repository.unimus.ac.id
Internet Source | 1% |
| 5 | Ping Yu, Xingxing Huang, Qian Ren, Xinxin Wang. "Purification and characterization of a H ₂ O ₂ -tolerant alkaline protease from Bacillus sp. ZJ1502, a newly isolated strain from fermented bean curd", Food Chemistry, 2019
Publication | 1% |

6	ipi.portalgaruda.org Internet Source	1 %
7	Mitchell Miglis, Donna Wilder, Thomas Reid, Irina Bakaltcheva. "Effect of taurine on platelets and the plasma coagulation system", <i>Platelets</i> , 2009 Publication	<1 %
8	paahjournal.com Internet Source	<1 %
9	Heni Setyowati Esti Rahayu, N. Nasruddin, Laela Hayu Nurani, Sri Darmawati et al. "Ethanol extract of the natural product of Daun sirih (Piper betle) leaves may impede the effectiveness of the plasma jet contact style for acute wounds", <i>Clinical Plasma Medicine</i> , 2019 Publication	<1 %
10	www.mdpi.com Internet Source	<1 %
11	Lesley Lee. "Culture-independent analysis of bacterial diversity in a child-care facility", <i>BMC Microbiology</i> , 2007 Publication	<1 %
12	P.R. Yaashikaa, M. Keerthana Devi, P. Senthil Kumar. "Engineering microbes for enhancing the degradation of environmental pollutants:	<1 %

A detailed review on synthetic biology",
Environmental Research, 2022

Publication

13

Jayarajapazham Rajaselvam, Natarajan Benit, Sager S. Alotaibi, M.A. Rathi et al. "In vitro fibrinolytic activity of an enzyme purified from Bacillus amyloliquefaciens strain KJ10 isolated from soybean paste", Saudi Journal of Biological Sciences, 2021

Publication

<1 %

14

"Chapter 42 4-Hydroxyphenylacetate 3-monooxygenase", Springer Science and Business Media LLC, 2006

Publication

<1 %

15

journals.ametsoc.org

Internet Source

<1 %

16

www.ncbi.nlm.nih.gov

Internet Source

<1 %

17

Ashis K. Mukherjee, Dhruba J. Chattopadhyay. " Potential clinical applications of phytopharmaceuticals for the in - patient management of coagulopathies in - 19 ", Phytotherapy Research, 2022

Publication

<1 %

18

Hb25_Springer Handbook of Marine Biotechnology, 2015.

Publication

<1 %

19 assets.publishing.service.gov.uk <1 %
Internet Source

20 mdpi-res.com <1 %
Internet Source

21 orbi.uliege.be <1 %
Internet Source

22 Sium Ahmed, Abdullah Mohammad Shohael, Kee Yoeup Paek. " Evaluation of growth and some unexplored bioactivities of bioreactor grown adventitious root culture of ginseng (C.A. Meyer) ", Biotechnology and Applied Biochemistry, 2021 <1 %
Publication

23 Nguyen Hoang Minh, Huynh Thi Quynh Trang, Tran Bao Van, Nguyen Hoang Loc. " Production and purification of nattokinase from ", Food Biotechnology, 2022 <1 %
Publication

Exclude quotes On

Exclude matches Off

Exclude bibliography On