

The leucocytes, neutrophils and superoxydes dismutase (SOD) level after consuming guava juice (*Psidium guajava* L) during aerobic exercise for beginners

by Yuliana Ulvie

Submission date: 17-Nov-2022 05:42AM (UTC+0700)

Submission ID: 1956188470

File name: BIK_ana.pdf (793.22K)

Word count: 3338

Character count: 18406

The leucocytes, neutrophils and superoxydes dismutase (SOD) level after consuming guava juice (*Psidium guajava* L) during aerobic exercise for beginners

Yuliana Noor Setiawati Ulvie^{1*}, Wiryatun Lestariana², Zaenal Muttaqien Sofro³

¹Nutritional Program, Faculty of Health Sciences and Nurshing, Universitas Muhammadiyah, Semarang, ²Department of Biochemistry, ³Department of Physiology, Faculty of Medicine, Universitas Gadjah Mada, Yogyakarta

ABSTRACT

Physical exercise is important in preventing and adjunctive therapy certain diseases. However, over physical exercise for beginners may cause oxidative stress and muscle injury leads to generate chemoattractants which enables to attract neutrophil and monocyte towards the injury and stimulates leucocytes activation. The use of antioxidant-rich fruits to mitigate exercise induced oxidative stress has been applied. This study was conducted to evaluate the effect of guava juice (*Psidium guajava* L) consumption during aerobic exercise on leucocytes, neutrophils and superoxydes dismutase (SOD) level of beginners. Sixteen students of National Land College (*Sekolah Tinggi Pertanian Nasional/STPN*), Yogyakarta who met the inclusion and exclusion criteria were involved in this study. Subjects were divided into two groups i.e. a group given mineral water as control and another group given 240 mL guava juice 0.67 g/mL daily for 27 days as treatment group. Both groups then underwent aerobic exercise for 30 minutes every day for 27 days. Blood samples were taken before and after exercise on day 1, 7 and 14 for leucocyte and neutrophil count as well as SOD analysis. The leucocyte count and SOD level before and after exercise in both Guava juice and Mineral water groups were not significantly different ($p > 0.05$). However, the percentage of neutrophil on day 1 and 7 were significantly higher than that in Mineral water group ($p < 0.05$). Moreover, during the exercise, the percentage of neutrophils in Guava juice group significantly increased ($p < 0.05$), whereas in Mineral water group was not significantly different ($p > 0.05$). It can be concluded that consuming guava juice during exercise for the beginners does not influence leucocyte count and SOD levels. However, it can increase percentage of neutrophil.

ABSTRAK

Latihan fisik penting dalam mencegah dan terapi pendamping untuk penyakit tertentu. Namun demikian, latihan fisik berlebihan bagi pemula kemungkinan bisa menyebabkan stres oksidatif dan luka otot yang dapat menghasilkan *chemoattractants* untuk menarik neutrofil, dan monosit ke daerah luka dan juga mengaktifasi leukosit. Penggunaan buah yang kaya antioksidan untuk mengurangi stres oksidatif akibat latihan fisik telah banyak dilakukan. Penelitian ini dilakukan untuk mengkaji pengaruh pemberian jus jambu merah pada latihan aerobik pemula terhadap leukosit, neutrofil dan kadar superoxydes dismutase (SOD) darah. Enam belas mahasiswa dari Sekolah Tinggi Pertanian (STPN), Yogyakarta yang memenuhi kriteria inklusi dan eksklusi dilibatkan dalam penelitian. Subjek dibagi menjadi dua kelompok yaitu kelompok kontrol yang diberi air mineral dan kelompok perlakuan yang diberi jus jambu merah 0,67 g/mL sebanyak 240 mL setiap hari selama 27 hari. Kedua kelompok selanjutnya melakukan aerobik selama 30 menit setiap dua hari sekali selama 27 hari. Sampel darah diambil sebelum dan setelah aerobik pada hari ke 1, 7 dan 14

* corresponding author: ulvieanna@gmail.com

untuk pemeriksaan leukosit, neutrofil dan kadar SOD darah. Jumlah leukosit dan kadar SOD darah sebelum dan setelah aerobik pada kelompok yang diberi jus jambu merah dan air mineral tidak berbeda nyata ($p > 0.05$). Namun demikian, persentase neutrofil pada hari ke 1 dan 7 aerobik lebih tinggi nyata dibandingkan dengan air mineral ($p < 0.05$). Selin itu, selam aerobik persentase neutrofil pada kelompok yang diberi jus jambu merah naik secara nyata ($p < 0.05$), sedangkan pada kelompok yang diberi air mineral tidak berbeda nyata ($p > 0.05$). Dapat disimpulkan bahwa konsumsi jus buah merah selama latihan fisik bagi pemula tidak mempengaruhi leukosit dan kadar SOD secara nyata, tetapi dapat meningkatkan persentase neutrofil.

Keywords: aerobic exercise - guava juice - leucocytes - neutrophils - SOD level

INTRODUCTION

Numerous studies have revealed a relationship between regular exercise and improvements in physical and mental healths. Physical exercise is important in preventing and adjunctive therapy certain diseases. The beneficial effects of exercise are associated with metabolic improvement and neutralization of some risk factors connected with diseases.^{1,2} However, it should be considered that over physical exercise may cause muscle damage called delayed onset muscle soreness (DOMS) due to mechanical trauma followed by oxidative stress.³

The DOMS is often experienced by the beginners. Over contraction and unusual use of muscle on the beginners during physical exercise need additional unit motor recruitment. Bone muscle contraction is a mechanical, chemical, and electrical process consisting of six stages called cross bridge cycle. For the beginners or untrained people, power stroke, sliding filament, and disconnecting in cross bridge cycle are mechanical trauma which may cause immune dysfunction, inflammation, oxidative stress, muscle soreness and muscle injury.^{4,6}

The muscle injury can generate chemo-attractants which enables to attract neutrophil and monocyte towards the injury site followed by reactive oxygen species (ROS) production.^{6,7} Moreover, oxidative stress due to over exercise causes the increase of body temperature and stimulates leucocytes production and activation. In physiological conditions, the ROS production

will be neutralized by endogenous antioxidant such as superoxydes dismutase (SOD), glutathione peroxides (GPx), and catalase. Oxidative stress will occur when the endogenous antioxidant is not enough to neutralize ROS production.^{8,9}

Recently, there is growing interest in the use of antioxidant-rich fruits or vegetables to mitigate exercise induced oxidative stress. Several fruits or vegetables have been studied for their benefits as dietary supplements and sport nutrition foods in countering oxidative stress during physical exercise.¹⁰ *Psidium guava* L. is a fruit tree commonly known as guava or *jambu biji* in Indonesai, which belongs to the family Myrtaceae. Guava fruit is rich antioxidant such as vitamin C, lycopene and flavonoids. Guava fruit is also a good source of pectin, sugar and minerals.¹¹⁻¹³ Guava fruit has been reported to have antidiarrheal, antibacterial and antioxidant activities.^{13,14} Commercially, the fruit is consumed raw or used in making jams, jellies, pastes and juice.¹¹

In this study, we reported the effect of guava juice consumption during aerobic exercise on leucocytes, neutrophils and superoxydes dismutase (SOD) level of beginners.

MATERIALS AND METHODS

Subjects

This was a quasi experimental study with nonrandomized pre-post test control group design involving students from National Land College (*Sekolah Tinggi Pertanahan Nasional/STPN*),

Yogyakarta who met inclusion and exclusion criteria. The inclusion criteria were students aged 17-25 years who stay in dormitory, normal nutritional status, normal healthy status based on clinical and hemoglobin examinations, non athletes, no suffered from chronic diseases such as asthma, hipertension, cardiovascular diseases, renal disease, and gartritis, no smoking, and never followed the aerobic exercise program based on frequency, intensity, time, and type (FITT) guidelines minimally in the last 6 years. The exclusive criteria include consuming antioxidant and or vitamin and not willing to be research subject. The study has been approved by the Medical and Health Research Ethics Committee, Faculty of Medicine, Universitas Gadjah Mada, Yogyakarta.

Procedure of study

On the day that has been agreed, the students were gathered to be selected. An explanation about objectives, benefits and risk of the study was given. Students were then given opportunities to ask everything related to the study. Sixteen students who fulfilled the inclusion and exclusion criteria were recruited in this study and given an informed consent to be signed. All subjects underwent physical and clinical as well as laboratory examinations. Characteristics of subjects including body weight (BW), body height (BH), systolic and diastolic blood pressure (SBP and DBP), hear rate (HR), respiratory rate, body temperature, haemoglobin level, leucocyte and neutrophils counts were recorded. Subjects were divided into two groups with 8 students in each group. Group I as control was given mineral water and Group II

as treatment group given was given 240 mL guava juice 0.67 g/mL daily for 27 days. The guava juice was given dailay at 05.00 AM. Subjects then underwent aerobic exercise every two days for 27 days consisting 30 minute for exercise, five minutes for warming up and five minutes for recovery. Body weight, blood pressure, HR, temperature of subjects before and after aerobic exercise were monitored and recorded. Food intake of subjects were calculated based on 24-hour food recall for 27 days using nutrisurvey program. Blood samples of subjects after exercise on day 1, 13 and 27 were collected for leucocyte and neutrophil count as well as SOD examination. Leucocyte and neutrophil counts were measured using Hematology Analyzer Sysmex XT-2000i, whereas blood SOD level was measured using cayman kit.

4 Statistical analysis

Data were presented as mean \pm standard deviation (SD). The different of leucocyte and neutrophil counts as well as blood SOD level inter group were analyzed using independent t-test. The different of measurement of each group was analyzed using repeated Anova. A p value $p < 0.05$ was considered as significant.

RESULTS

The characteristics of subjects of Guava juice and Mineral water groups before exercise are presented in TABLE 1. No significant difference was observed between Guava juice group and Mineral water group before exercise indicating that both of groups were similar.

TABLE 1. Characteristics of subjects of Guava juice and Mineral water groups

Variable	Guava juice (mean ± SD)	Mineral water (mean ± SD)
Body weight (kg)	58.50 ± 2.75	59.50 ± 3.72
Body height (cm)	171.75 ± 4.17	167.00 ± 4.22
Body Mass Index	19.84 ± 0.86	21.38 ± 1.80
Heart rate (pulse/minute)	82.00 ± 6.05	80.00 ± 6.84
Respiratory rate (x/minute)	19.00 ± 1.51	19.50 ± 2.07
Body temperature (°C)	36.11 ± 0.49	35.99 ± 0.59
Hemoglobin (g/dL)	15.86 ± 0.67	15.16 ± 0.53
Hematocrit (%)	47.45 ± 2.25	45.84 ± 1.84
Leucocyte (μL ⁻¹)	8217 ± 1862	7478 ± 1716
Neutrophil (%)	54.12 ± 6.13	48.12 ± 7.51

The mean of BW, HR and body temperature of subjects before and after exercise in Guava juice and Mineral water groups are presented in TABLE 2. Change in BW was observed before and after exercise in both groups (p<0.05). Heart rate of both groups before exercise was significantly

different (p<0.05), however it was not significantly different after warming up, exercise and recovery (p>0.05). The body temperature between Guava juice and Water mineral before exercise was significantly different (p<0.05), however after exercise it was not significantly different (p>0.05).

TABLE 2. Mean of BW, HR and body temperature of subjects before and after exercise in Guava juice and Mineral water groups

Variable	Guava juice	Mineral water	p
BW before exercise (kg)	58.07 ± 0.21	57.86 ± 0.20	0.000
BW after exercise (kg)	57.86 ± 0.20	59.10 ± 0.35	0.000
HR before exercise (x/minute)	86.14 ± 3.58	82.53 ± 2.89	0.007
HR after warming up (x/minute)	125.57 ± 12.68	123.87 ± 13.45	0.733
HR after exercise (x/minute)	157.67 ± 0.71	158.01 ± 0.76	0.236
HR after recovery (x/minute)	121.01 ± 4.22	117.92 ± 5.73	0.116
Temperature before exercise (°C)	35.78 ± 0.15	35.58 ± 0.17	0.003
Temperature after exercise (°C)	35.95 ± 0.21	35.87 ± 0.27	0.348

The nutrient intake of subjects in Guava juice and Mineral water groups during study are presented in TABLE 3. A significantly higher intake in energy, fat, protein, vitamin C, vitamin E, cuprum, zinc and mangan was observed in Guava

juice group compare to Mineral water group (p<0.05). However, the carbohydrate intake was not significantly different between two groups (p>0.05).

TABLE 3. Nutrient intake of subjects in Guava juice and Mineral water groups

Nutrient intake	Guava juice	Mineral water	p
Energy (kcal)	2807.83 ± 235.85	2405.67 ± 263.81	0.006
Fat (g/day)	99.40 ± 6.71	85.07 ± 11.95	0.033
Protein (g/day)	95.36 ± 5.98	84.68 ± 11.26	0.010
Carbohydrate (g/day)	349.50 ± 18.90	318.99 ± 38.49	0.064
Vitamin C (mg/day)	225.98 ± 23.70	45.75 ± 13.17	0.000
Vitamine E (mg/day)	7.52 ± 0.95	5.57 ± 1.00	0.001
Iron (mg/day)	24.17 ± 10.29	14.97 ± 3.47	0.031
Cuprum (mg/day)	2.62 ± 0.36	2.02 ± 0.29	0.003
Zinc (mg/day)	10.74 ± 1.01	9.43 ± 1.07	0.025
Mangan (mg/day)	31.26 ± 7.52	24.09 ± 5.42	0.046

The leucocytes count before and after exercise in Guava juice and Mineral water groups are presented in TABLE 4. No significantly difference

in the leucocytes count before and after exercise in both groups was observed in this study ($p > 0.05$).

TABLE 4. Leucocytes count (μL^{-1}) before and after exercise in Guava juice and Mineral water groups

Exercise	Guava juice	Mineral water	p*	p**	p***
Before exercise	7770 ± 1545	6652 ± 1154	0.123		
After exercise 1	7687 ± 1939	6686 ± 1035	0.219	0.678	0.092
After exercise 7	8076 ± 1440	7564 ± 1153	0.445		
After exercise 14	7340 ± 1735	7108 ± 1734	0.794		

Note: * t-test; ** Anova on Guava juice group; *** Anova on Mineral water group

The percentage of neutrophils before and after exercise in Guava juice and Mineral water groups are presented in TABLE 5. No significantly difference in the percentage of neutrophils before exercise and after exercise on day 14 between two groups was observed ($p > 0.05$). However, after exercise on day 1 and 7 the percentage of

neutrophils in Guava juice group were significantly higher than that in Mineral water group ($p < 0.05$). During the exercise, the percentage of neutrophils in Guava juice group significantly increased ($p < 0.05$), whereas in Mineral water group was not significantly different ($p > 0.05$).

TABLE 5. Percentage of neutrophil (%) before and after exercise in Guava juice and Mineral water groups

Exercise	Guava juice	Mineral water	p*	p**	p***
Before exercise	50.62±5.12	45.75±4.03	0.053		
After exercise 1	62.25±10.09	50.62±8.24	0.024	0.004	0.160
After exercise 7	52.75±5.15	43.87±6.40	0.009		
After exercise 14	51.25±4.30	48.37±8.60	0.412		

Note: * t -test; ** Anova on Guava juice group ; *** Anova on Mineral water group

The blood SOD level before and after exercise in Guava juice and Mineral water groups are presented in TABLE 6. Although the blood SOD level tended to be higher after exercise, however

no significantly difference in the blood SOD level before and after exercise in both groups was observed in this study ($p > 0.05$).

TABLE 6. Level of blood SOD (U/mL) before and after exercise in Guava juice and Mineral water groups

Exercise	Guava juice	Mineral water	p*	p**	p***
Before exercise	2.31 ± 0.94	2.27 ± 0.98	0.924		
After exercise 7	2.34 ± 1.88	2.49 ± 1.02	0.837	0.247	0.190
After exercise 14	3.06 ± 0.66	3.24 ± 1.46	0.759		

Note: * t -test; ** Anova on Guava juice group ; *** Anova on Mineral water group

DISCUSSION

Regular physical exercise has many health benefits including a reduced risk of certain diseases such as cardiovascular disease, cancer, and diabetes.^{14,15} Otherwise, it is also widely accepted that contracting skeletal muscles generate free radicals and that prolonged and intense exercise can result in oxidative stress and muscle injury.^{16,7,16} Furthermore, the oxidative stress has substantial effect on leucocytes life span. The amounts of several hormones, cytokines and other factors which might influence cellular survival are increased or decreased in organs, tissues and peripheral blood during exercise.^{17,18}

This study showed that in Mineral water group the amount of leucocytes and percentage of neutrophil tended to increase after exercise on day 1 and 7, although it was not significantly different. It was indicated that actually aerobic exercise for beginner induce oxidative stress that lead to stimulate leucocytes production due to additional unit motor recruitment and muscle injury. However, ROS production due to the oxidative stress might be neutralized by endogenous antioxidant SOD that found also tended to increase in this study. Although it was also not significantly different. Result of this study is in accordance with study conducted by Shojaei *et al.*⁹ that reported that leucocyte increased immediately after 45

minutes cycling with intensity 50% of VO_2 max for untreated human.

Furthermore, in this study also found that the amount of leucocytes and percentage of neutrophil tended to decrease after exercise on day 14, although it was not significantly different. These decrease might be caused by leucocytes apoptosis after the prolongation of aerobic exercise. Exercise-induced leucocyte apoptosis has been reported previously. Syu *et al.*¹⁹ demonstrated that acute severe exercise induced an oxidative stress which resulted in acceleration of spontaneous neutrophil apoptosis. Moreover, repeated moderate exercise (30 min a day, 5 days a week at 60% of maximal workload) delayed neutrophil apoptosis.²⁰ A recent study reported that a significant delay of neutrophil apoptosis after marathon run, intensive endurance and downhill running as well as intensive resistance exercise was observed.²¹

Similar results were obtained on Guava juice group for leucocytes which tended increase after exercise on day 1 and 7 and decrease on day 14, whereas SOD level tended to increase on day 7 and 14. Although, these were not significantly different. However, the percentage of neutrophil significantly increased during aerobic exercise. It was indicated that guava juice can protect exercise-induced neutrophil apoptosis.

The effect of guava fruit to protect exercise-induced neutrophil apoptosis might be caused its antioxidant activity. Guava fruit is rich antioxidant such as vitamin C, vitamin E, lycopene and flavonoids. Guava fruit is also a good source of minerals such as copper, iron, zinc, magnesium and selenium.¹¹⁻¹³ Moreover, the antioxidant activity of guava fruit has been proven by some authors.^{13,22,23} Antioxidant supplements may benefit exercise performance directly through the reduction of muscle fatigue at the level of contractile function and indirectly through reduction of physiological stressors that negatively impact on training or improvement in the ability to recover from training.^{6,24}

CONCLUSION

It can be concluded that consuming guava juice during exercise for the beginners does not significantly influence leucocyte count and SOD levels. However, the consuming guava juice can significantly increase percentage of neutrophil.

ACKNOWLEDGEMENTS

We would like to thank all volunteers who participated in this study. We also grateful to Director of STPN, Yogyakarta Prof. Endriatmo Soetarto who has given his permission to conduct this study. This study can not be conducted without valuable assistances from technicians of Biochemistry Laboratory, Faculty of Medicine, Universitas Gadjah Mada.

REFERENCES

1. Penedo FJ and Dahn JR. Exercise and well-being: a review of mental and physical health benefits associated with physical activity. *Curr Opin Psychiatry* 2005; 18(2): 189–93. Doi:10.1097/00001504-200503000-00013
2. Puetz TW, O'Connor PJ, Dishman RK. Effects of chronic exercise on feelings of energy and fatigue: a quantitative synthesis. *Psychol Bull* 2006; 132(6): 866–76. Doi:10.1037/0033-2909.132.6.866
3. Foss LM. *Physiological basis for exercise and sport*. New York: McGraw Hill Book Company, 1998.
4. Len J, Davies CT, Young K. Changes in indicators of inflammation after eccentric exercise of the elbow flexors. *Med Sci Sports Med* 2002; 25:236-9.
5. Nieman DC. Immune function responses to ultramarathon race competition. *Med Sportiva* 2009; 13: 189–96.
6. Powers SK, Jackson MJ. Exercise-induced oxidative stress: cellular mechanisms and impact on muscle force production. *Physiol Rev* 2008; 88: 1243–76.
7. Barbieri E and Sestili P. Reactive oxygen species in skeletal muscle signaling. *J Signal Transduct* 2012;2012:982794. Doi: 10.1155/2012/982794.
8. Halliwell B, Gutteridge JMC. *Free radical in biology and medicine*. 3rd ed. New York: Oxford University Press Inc, 1999.

9. Shojaei EA, Jafari A, Farajov A. Effect of acute moderate aerobic cycling on systemic inflammatory responses in young untrained men. *Sci Sports* 2010; 1-5.
10. Nieman DC, Stear SJ, Castell LM, Burke LM. A-Z of nutritional supplements: dietary supplements, sports nutrition foods and ergogenic aids for health and performance: part 15. *Br J Sports Med* 2010; 44: 1202-5.
11. Kamath J V, Rahul N, Ashok Kumar CK, Lakshmi S M. *Psidium guajava* L: A review. *Int J Green Pharm* 2008;2(1):9-12
12. Sunttornusk L. Quantitation of vitamin C content in herbal juice using direct titration. *J Pham Biomed Anal* 2002; 28(5):849-55.
13. Hassimotto NM, Genovese MI, Lajolo FM. Antioxidant activity of dietary fruits, vegetables, and commercial frozen fruit pulps. *J Agric Food Chem* 2005; 53: 2928-35.
14. Blair SN, Cheng Y, Holder JS. Is physical activity or physical fitness more important in defining health benefits? *Med Sci Sports Exerc* 2001;33:S379-99.
15. Crespo CJ, Palmieri MR, Perdomo RP, McGee DL, Smit E, Sempos CT, *et al.* The relationship of physical activity and body weight with all-cause mortality: results from the Puerto Rico Heart Health Program. *Ann Epidemiol* 2002; 12:543-52.
16. Fisher-Wellman K and Bloomer RJ. Acute exercise and oxidative stress: a 30 year history. *Dynamic Med* 2009; 8(1): doi:10.1186/1476-5918-8-1
17. Krammer PH, Arnold R, Lavrik IN. Life and death in peripheral T cells. *Nat Rev Immunol* 2007; 7(7):532-42.
18. Malm C, Nyberg P, Engstrom M, Sjodin B, Lenkei R, Ekblom B, *et al.* Immunological changes in human skeletal muscle and blood after eccentric exercise and multiple biopsies. *J Physiol* 2000;529(Pt 1):243-62.
19. Syu GD, Chen HI, Jen CJ. Acute severe exercise facilitates neutrophil extracellular trap formation in sedentary but not active subjects. *Med Sci Sports Exerc* 2013; 45(2):238-44.
20. Su SH, Jen CJ, Chen HI. NO signaling in exercise training-induced antiapoptotic effects in human neutrophils. *Biochem Biophys Res Commun* 2011;405(1):58-63. Doi: 10.1016/j.bbrc.2010.12.123.
21. Mooren FC, Volker K, Klocke R, Nikol S, Waltenberger J, Kruger K. Exercise delays neutrophil apoptosis by a G-CSF-dependent mechanism. *J Appl Physiol* 2012; 113(7):1082-90.
22. Jiménez-Escrig A, Rincón M, Pulido R, Saura-Calixto. Guava fruit (*Psidium guajava* L.) as a new source of antioxidant dietary fiber. *J Agric Food Chem* 2001; 49:5489-93.
23. Gull J, Sultana B, Anwar F, Naseer R, Ashraf M, Ashrafuzzaman. Variation in antioxidant attributes at three ripening stages of guava (*Psidium guajava* L.) fruit from different geographical regions of Pakistan. *Molecules* 2012; 173165-80. Doi:10.3390/molecules17033165
24. Reid MB. Invited review: redox modulation of skeletal muscle contraction: what we know and what we don't. *J Appl Physiol* 2001; 90(2):724-31.

The leucocytes, neutrophils and superoxydes dismutase (SOD) level after consuming guava juice (*Psidium guajava* L) during aerobic exercise for beginners

ORIGINALITY REPORT

14%

SIMILARITY INDEX

16%

INTERNET SOURCES

8%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

1	ojs.unpkediri.ac.id Internet Source	4%
2	moam.info Internet Source	3%
3	img1.wsimg.com Internet Source	3%
4	repository.stikim.ac.id Internet Source	2%
5	Guntur Muhammad Taqwin, I Gusti Ngurah Artika, Sri Rahardjo. "Optimal time of administration of fentanyl in reducing hemodynamic response in endotracheal intubation", <i>Journal of the Medical Sciences (Berkala Ilmu Kedokteran)</i> , 2014 Publication	2%

Exclude bibliography On