

Turnitin Artikel Mutiara Medika

dr. Ika

by Dr, Ika

Submission date: 30-Jan-2023 09:44AM (UTC+0700)

Submission ID: 2002030296

File name: 4._15599_21-27.pdf (598.89K)

Word count: 3772

Character count: 20833

Antimicrobial Activity of Pecut Kuda Leaf Extract (*Stachytarpheta jamaicensis* (L. Vahl) against *Mycobacterium smegmatis*

Putri Madia Ramadhani¹, Ika Dyah Kurniati², Maya Dian Rakhmawati^{2*}

¹Faculty of Medicine, Universitas Muhammadiyah Semarang, Semarang, Indonesia

²Department of Biomedical Sciences, Faculty of Medicine, Universitas Muhammadiyah Semarang, Semarang, Indonesia

DATE OF ARTICLE:

Received: 25 July 2022

Reviewed: 20 Dec 2022

Revised: 03 Jan 2023

Accepted: 07 Jan 2023

*CORRESPONDENCE:

mayadianr@unimus.ac.id

DOI:

10.18196/mmjkk.v22i2.15599

TYPE OF ARTICLE:

Research

Abstract: *Mycobacterium smegmatis* infrequently causes infection, but it is easy to be pathogenic in immunosuppressed patients. Many reported that *M. smegmatis* resistance to several antibiotics became an impetus for searching for new antimicrobials. Therefore, this study aims to prove the effect of Pecut Kuda leaf extract (*Stachytarpheta jamaicensis* (L.) Vahl) on the growth of *M. smegmatis* mc2 155. This research is an experimental study with a post-test control group design. The susceptibility test was carried out using the two-fold microdilution method and resazurin staining. The concentration of Pecut Kuda leaf ethanol extract was prepared in the concentration range of 10000.0 – 625.0 µg/ml. Phytochemical analysis of the content of saponins, tannins, flavonoids, and alkaloids was also carried out on Pecut Kuda leaf ethanol extract. Pecut Kuda leaf ethanol extract can inhibit the growth of *M. smegmatis* with a minimum inhibitory concentration (MIC) of 5000 µg/ml (very weak activity) because, at the highest concentration of 10000 µg/ml, *M. smegmatis* still cannot be killed. Furthermore, Pecut Kuda leaf ethanol extract contains saponins, tannins, flavonoids, and alkaloids which are known to have antibacterial activity. However, further evaluation is needed to maximize the antibacterial activity of Pecut Kuda leaf extract, for example, by fractionating the extract.

Keywords: antibacterial agent; *Mycobacterium smegmatis*; phytochemical analysis; *Stachytarpheta jamaicensis* (L.) Vahl

INTRODUCTION

Mycobacterium smegmatis is an acid-fast bacillus classified as a rapidly growing (RGM) Non-Tuberculosis Mycobacteria.¹ These bacteria infrequently cause infection but can easily infect immunosuppressed patients. Infections due to *M. smegmatis* are non-tuberculous infections and have broad clinical manifestations such as hypersensitivity, pneumonitis, asthma and bronchitis, skin infection, wounds, and glands.² The prevalence of non-tuberculous infections in Indonesia has increased since 2000. The incidence of these infections in Soetomo Hospital from 2014 to 2015 reached 5.78%. These infections increased in 2017, reaching 25% of patients.³ The most common clinical manifestation caused by *M. smegmatis* is pulmonary infection.⁴

Treatment of non-tuberculous infection consists of prophylaxis and medication.⁵ In terms of drug therapy, antibiotics such as erythromycin, clarithromycin, rifampin, rifabutin, and ethambutol can be used to treat this infection.⁶ However, there have been many reports of resistance of *M. smegmatis* to several antibiotics such as ampicillin, amoxicillin, linezolid, erythromycin, streptomycin, and tetracycline.⁷ It has become an impetus for searching for new antimicrobials, especially those from natural products. One source of natural products that can be explored is medicinal plants. Pecut Kuda leaves (*Stachytarpheta jamaicensis* (L.) Vahl) is a plant that has the potential as an antibacterial. The antibacterial activity of Pecut Kuda extract has been tested against several bacteria. Pecut Kuda leaf ethanol extract with a concentration of 12.5% is known to inhibit the growth of *Escherichia coli*. In addition, Pecut Kuda leaf extract with a concentration of

80% can also inhibit the growth of Gram-positive bacteria *Streptococcus pyogenes*, with an average inhibition zone diameter of 29.2 mm. The extract is known to contain compounds that have the potential as antimicrobials, including flavonoids, tannins, saponins, and alkaloids.^{8,9}

Research on the effect of Pecut Kuda leaf (*S. jamaicensis* (L.) Vahl) extract on *M. smegmatis* has not been carried out until now. Therefore, this research was conducted to identify the potential of Pecut Kuda leaf extract to inhibit the growth of *M. smegmatis*. In addition, phytochemical analysis was carried out to qualitatively identify the content of chemical compounds in the Pecut Kuda leaves.

MATERIAL AND METHOD

The test bacteria used in this study was *M. smegmatis* mc2 155 (ATCC 700084), obtained from the Microbiology Laboratory, Faculty of Medicine, Universitas Muhammadiyah Semarang. This research has been approved by the Health Research Ethics Commission of the Medical Faculty of Universitas Muhammadiyah Semarang under the number 160/EC/FK/2021.

Secondary Metabolites Extraction of Pecut Kuda Leaves (*Stachytarpheta jamaicensis* (L.) Vahl)

First, Pecut Kuda leaves were dried under the sun with a covered cloth for approximately three days. Furthermore, 1 kg of Pecut Kuda leaves was prepared in the form of dried simplicia. The simplicia powder of Pecut Kuda leaves was soaked with 96% ethanol solvent for five days and stirred occasionally. After five days, the final result of the maceration process of dried Pecut Kuda Simplicia was filtered, and the supernatant was concentrated using a rotary evaporator (Ika® RV 10) at a temperature of 70 °C and a speed of 100 rpm.

Phytochemical Analysis of Pecut Kuda (*Stachytarpheta jamaicensis* (L.) Vahl) Leaves Ethanol Extract

Phytochemical analysis was carried out to identify the content of chemical compounds in Pecut Kuda leaf ethanol extract. The test was conducted to qualitatively identify the presence of tannins, saponins, flavonoids, and alkaloids. In the tannin test, a total of 0.5 g of Pecut Kuda leaf extract was put into a test tube. 20 ml of distilled water was then added to the test tube and heated to boiling. The boiling solution was filtered and added with 1% FeCl₃. The positive presence of tannins is indicated by a change in color to green or blue-black ink.¹⁰⁻¹²

Furthermore, in the saponins test, 100 mg of extract was mixed with 5 ml of water and shaken vigorously. If the solution forms a stable foam, it indicates the presence of saponins. To determine the presence of flavonoids, 0.5 ml of the extract was added with 5 ml of dilute ammonia and 2 ml of concentrated sulfuric acid. Flavonoid compounds can be seen from the appearance of a brown color in the solution. Meanwhile, to identify alkaloid compounds, as much as 1 ml of Pecut Kuda leaf extract was added with 1% HCl in a test tube. The test tube was heated slowly, and the solution was filtered. 3 drops of Mayer and Wagner reagent can be added to the filtered supernatant. If a precipitate is formed in the solution, it is confirmed that the presence of alkaloids in the Pecut Kuda leaf extract is confirmed.¹⁰⁻¹²

Antibacterial Activity Test of Pecut Kuda Leaves (*Stachytarpheta jamaicensis* (L.) Vahl) Ethanol Extract against *Mycobacterium smegmatis*

To determine the minimum inhibitor concentration (MIC) of Pecut Kuda leaf ethanol extract, the susceptibility test was carried out using the two-fold microdilution method. The concentration of Pecut Kuda leaf extract was prepared in the concentration range of 10000.0 – 625.0 µg/ml. The concentration of the *M. smegmatis* suspension in each well of the microplate was 1×10^5 CFU/mL. Furthermore, to determine the activity of Pecut Kuda leaf extract in inhibiting *M. smegmatis*, the microplate was incubated at 37 °C for 24 hours. After incubation, 0.0025% resazurin (Sigma Aldrich), as much as 20 µl, was added to each test well. The color change of resazurin from blue to pink became a parameter for the growth of *M. smegmatis*.¹³

As an antibiotic control, Rifampicin (Phapros Tbk) was used to determine the resistance pattern of *M. smegmatis*. The susceptibility test method of rifampicin against *M. smegmatis* was done using the same method to test the activity of Pecut Kuda leaf extract, but the concentration of rifampicin was prepared in the range of 2.0 – 0.0625 µg/ml. After obtaining the MIC value, the research continued to determine the minimum bactericidal concentration (MBC). The test suspension in each well of various concentrations of

Pecut Kuda leaf extract/rifampicin was taken as much as 10 μ l. Furthermore, the suspension was grown in Mueller Hinton Agar (Merck Millipore) using the streak method. The Petri dishes were then incubated at 37 $^{\circ}$ C for 24 hours. The MBC value was determined from the smallest Pecut Kuda leaf extract/rifampicin concentration that caused the absence of *M. smegmatis* growth.

RESULT

The MIC value was determined from the smallest concentration of the microplate wells that did not change their color to pink. Based on the result of the susceptibility test, the MIC value of Pecut Kuda leaf (*Stachytarpheta jamaicensis* (L.) Vahl) extract in inhibiting the growth of *M. smegmatis* was 5000 μ g/ml (Figure 1).




Figure 1. The result of the minimum inhibitory concentration of Pecut Kuda Leaf (*Stachytarpheta jamaicensis* (L.) Vahl) extract against *M. smegmatis* carried out with two-fold microdilution and resazurin staining

Note: B-E2 until B-E6 is a well containing Pecut Kuda leaf extract with a concentration range of 10000 - 625 μ g/ml; B-E7 are well of Pecut Kuda leaf extract sterility control; B-E10 are *M. smegmatis* growth control; and B-E11 are media sterility control.

Furthermore, for the results of the susceptibility testing of rifampicin against *M. smegmatis*, the MIC value was 0.0625 μ g/ml (Figure 2). Based on the MIC value, it can be concluded that *M. smegmatis* is still sensitive to rifampicin.¹⁴




Figure 2. The result of the minimum inhibitory concentration of rifampicin against *M. smegmatis* carried out with two-fold microdilution and resazurin staining

Note: B-E2 until B-E7 is a well-containing rifampicin with a concentration range of 2 - 0.0625 μ g/ml; B-E8 are well of rifampicin sterility control; B-E11 are *M. smegmatis* growth control.

Both Pecut Kuda leaf (*Stachytarpheta jamaicensis* (L.) Vahl) extract and rifampicin were still able to inhibit the growth of *M. smegmatis*. Therefore, the test was continued to determine the MBC value. Based on the results of *M. smegmatis* growth on agar media, it can be concluded that Pecut Kuda leaf extract was still unable to kill *M. smegmatis*. Meanwhile, at the smallest concentration used for the test, rifampicin had no *M. smegmatis* growth. Therefore, it can be concluded that the MBC value of rifampicin against *M. smegmatis* is 0.0625 µg/mL.

Because Pecut Kuda leaf extract had the potential to be developed as an antibacterial, a phytochemical analysis was carried out to determine the compound content. According to the results of the phytochemical analysis, it can be concluded that the Pecut Kuda leaf extract contained saponins, tannins, flavonoids, and alkaloids (Table I).

Table I. Result of Phytochemical Analysis of Pecut Kuda (*Stachytarpheta jamaicensis* (L.) Vahl) Leaf Extract

Compound	Phytochemical Test Observation	Result
Saponins	Stable foam	+
Tannins	Dark blue	+
Flavonoids	Maroon	+
Alkaloids	Dissolving precipitation	+

DISCUSSION

Determination of the MIC value of Pecut Kuda (*Stachytarpheta jamaicensis* (L.) Vahl) leaf extract and rifampicin was carried out using the Resazurin Microplate Assay (REMA) method. This method is known to be utilized for the rapid screening of natural product compounds' antimycobacterial activity compared with the Crystal Violet Decolorization Assay (CVDA) method. In addition, the change in resazurin color from blue to pink due to *M. smegmatis* growth can simplify the interpretation of the test results.

Based on the results of this study, Pecut Kuda leaf extract was able to inhibit *M. smegmatis* growth with a MIC value of 5000 µg/mL. However, Pecut Kuda leaf extract could still not kill the *M. smegmatis* at the highest test concentration of 10000 µg/mL. There has never been a study reporting the activity of Pecut Kuda leaf extract against *M. smegmatis*. However, Pecut Kuda leaf extract has been known to have antibacterial activity against other Gram-positive bacteria. Pecut Kuda leaf extract was able to inhibit the growth of *S. pyogenes*, and it has been developed into an antiseptic candy dosage form.^{9,15} In addition, Pecut Kuda leaf extract can inhibit the growth of *Staphylococcus aureus*¹⁶ and several other pathogenic bacteria such as *E. coli*, *Klebsiella pneumoniae*, *Pseudomonas aeruginosa*, and *Salmonella typhimurium*.¹⁷ The ability of Pecut Kuda leaves extract to inhibit the growth of *M. smegmatis* can be categorized as very weak compared to its activity against other Gram-negative and positive bacteria. The MIC value of 5000 µg/mL was also very high compared to the MIC value of the control drug, rifampicin. Furthermore, according to another study, an extract can be developed as an antibacterial if the MIC value is less than 100 µg/mL.¹⁸

Mycobacterium smegmatis mc2 155 used in this study was a wild-type isolate, and it was proven that rifampicin could inhibit the growth and kill *M. smegmatis* at a concentration of 0.0625 µg/mL. Although the majority of *M. smegmatis* strains are not susceptible to tuberculosis drugs such as rifampicin, isoniazid, and ethambutol, the result is relevant to other research, which states that *M. smegmatis* strain ATCC 607 was reported to be susceptible to rifampicin with a MIC value of 0.97 - 1.6 µg/mL.¹⁹ *Mycobacterium smegmatis* resistance pattern to rifampicin has been widely reported. The MIC value of rifampicin can reach 20 µg/mL in *M. smegmatis* resistant. *Mycobacterium smegmatis* itself can become resistant to rifampicin due to the mechanism of cellular permeability, enzymatic modification, and efflux system.²⁰

Furthermore, according to the results of phytochemical tests, Pecut Kuda leaf extract contained saponins, tannins, flavonoids, and alkaloids. All of these compounds are known to have the potential as antibacterial. Flavonoids have antibacterial activity, as most phenols can inhibit protein synthesis and cause protein denaturation in cell walls. This activity is then followed by a disruption in the permeability of the bacterial cell wall.²¹ In particular, flavonoids' effect on mycobacteria inhibits mycolic acid synthesis. Molecular docking tests of several flavonoids (quercetin and taxolin) showed their activity to inhibit mycobacterium DNA gyrase, preventing bacterial cells from dividing.²² The effect of flavonoids on mycobacteria cell walls was also demonstrated by inhibiting the glutamate racemase enzyme, which inhibited peptidoglycan synthesis.²³ Meanwhile, saponins can have antibacterial activity by inhibiting cell membrane function, ultimately changing cell membrane permeability and damaging cell walls.²⁴ Until now, the mechanism of saponins against mycobacteria has never been reported. Furthermore, Alkaloids have a mechanism to inhibit cell wall

synthesis and interfere with peptidoglycan components, so bacterial cell walls are not formed properly.²⁵ Alkaloid activity in mycobacterium mainly depends on the ease of transporting substances through the cell membrane. Alkaloids are lipophilic compounds that can easily penetrate the walls of mycolic acid. The berberine derivative is an example of an alkaloid with antimycobacterial activity.^{26,27} The last compound, tannin, has antibacterial activity related to its ability to activate microbial cell adhesion, inactivate enzymes, and interfere with protein transport in the inner layer of cells.²⁸

When viewed from the mechanism of action of flavonoids, saponins, and alkaloids, *Pecut Kuda* leaf extract has the potential to be bactericidal. However, this study did not isolate compounds, so the type and amount of flavonoids, saponins, and alkaloids contained in *Pecut Kuda* leaf extract are unknown. Therefore, it cannot be ascertained why *Pecut Kuda* leaf extract identified in this study could not kill the *M. smegmatis*. Another possible reason for the lack of bactericidal effect of *Pecut Kuda* leaf extract is that *M. smegmatis* has N-acetylmuramic acid (MurNAc) and N-glycolylmuramic acid (MurNGlyc), which can increase drug efflux.²⁹ In addition, the cell wall of mycobacteria is different from that of Gram-positive and negative bacteria in general. Mycobacterial cell walls may consist of the mycoyl-arabinogalactan-PG (mAGP) complex. Peptidoglycan is bound covalently to arabinogalactan, which is esterified with mycolic acid and becomes a lipid barrier in mycobacteria. It is very useful for controlling the cell wall's osmotic pressure and reduces the effectiveness of *Pecut Kuda* leaf extract as an antibacterial.³⁰ Despite all that, *Pecut Kuda* leaf (*Stachytarpheta jamaicensis* (L.) Vahl) still has the potential to be developed into antibacterial dosage forms. However, prior evaluation is needed to maximize the activity of the *Pecut Kuda* leaf extract, such as by fractionating the extracts to obtain purer active compounds.

CONCLUSION

Pecut Kuda leaf (*Stachytarpheta jamaicensis* (L.) extract could inhibit the growth of *Mycobacterium smegmatis* with minimum inhibitory concentration (MIC) of 5000 µg/ml (very weak activity), and at the highest concentration of 10000 µg/ml, they were still not able to kill the *M. smegmatis*. Furthermore, *Pecut Kuda* leaf extract contained saponins, tannins, flavonoids, and alkaloids which were known to have antibacterial activity.

CONFLICT OF INTEREST

None of the Conflicts of Interest

REFERENCES

1. Alqurashi MM, Alsaileek A, Aljizeeri A, Bamefleh HS, Alenazi TH. *Mycobacterium smegmatis* causing a granulomatous cardiomedastinal mass. IDCases. 2019 Jan;18:e00608. <https://doi.org/10.1016/j.idcr.2019.e00608>
2. Moghim S, Sarikhani E, Esfahani BN, Faghri J. Identification of nontuberculous mycobacteria species isolated from water samples using phenotypic and molecular methods and determination of their antibiotic resistance patterns by E- test method, in Isfahan, Iran. Iran J Basic Med Sci. 2012;15(5):1076-82. <https://pubmed.ncbi.nlm.nih.gov/23493797>
3. Mertaniasih N, Kusumaningrum D, Koendhori E, Soedarsono, Kusmiati T, Dewi DSS. Nontuberculous mycobacterial species and *Mycobacterium tuberculosis* complex coinfection in patients with pulmonary tuberculosis in dr. Soetomo hospital, Surabaya, Indonesia. Int J Mycobacteriology. 2017 Jan;6(1):9-13. <https://www.ijmyco.org/text.asp?2017/6/1/9/201894>
4. Sousa S, Borges V, Joao I, Gomes JP, Jordao L. Nontuberculous mycobacteria persistence in a cell model mimicking alveolar macrophages. Microorganisms. 2019 May;7(113). <https://doi.org/10.3390/microorganisms7050113>
5. Restiawati NM, Burhan E. Diagnosis dan penatalaksanaan mycobacterium othe than tuberculosis (MOTT). J Respir Indo. 2011;31(3):156-64.
6. Juita LR, Fauzar F. Diagnosis dan tatalaksana penyakit paru nontuberculous mycobacteria. J Kesehat Andalas. 2018;7(Supplement 3):141. <https://doi.org/10.25077/jka.v7i0.858>
7. Best CA, Best TJ. *Mycobacterium smegmatis* infection of the hand. Hand. 2009;4(2):165-6. <https://doi.org/10.1007%2Fs11552-008-9147-6>

8. Kumala S, Bekti NDP. Aktivitas antibakteri dan antioksidan daun pecut kuda (*Stachytarpheta jamaicensis* L.) secara *in vitro*. J Farm Indones. 2016;8(2):137-43.
9. Novitasari D, Mahtuti E, Ibaadillah A. Uji daya hambat perasan daun pecut kuda (*Stachytarpheta jamaicensis* L. Vahl) terhadap pertumbuhan bakteri *Streptococcus pyogenes* secara *in vitro*. Garuda. 2017;8(2).
10. Thangiah AS. Phytochemical screening and antimicrobial evaluation of ethanolic-aqua extract of *Stachytarpheta jamaicensis* (L.) vahl leaves against some selected human pathogenic bacteria. Rasayan J Chem. 2019;12(1):300-7. <http://dx.doi.org/10.31788/RJC.2019.1215042>
11. Indarto I. Uji kualitatif Dan kuantitatif golongan senyawa organik dari kulit dan kayu batang tumbuhan *Artocarpus dadah* Miq. J Ilm Pendidik Fis Al-Biruni. 2015;4(1):75-84. <https://doi.org/10.24042/jipfalbiruni.v4i1.82>
12. Rijayanti RP, Luliana S, Trianto HF. Uji aktivitas antibakteri ekstrak etanol daun mangga bacang. Univ Tanjungpura. 2014;1(1):13-4.
13. Rakhmawatie MD, Wibawa T, Lisdiyanti P, Pratiwi WR, Mustofa. Evaluation of crystal violet decolorization assay and resazurin microplate assay for antimycobacterial screening. Heliyon. 2019 Aug;5(8):1-7. <https://doi.org/10.1016/j.heliyon.2019.e02263>
14. CLSI. Appendix J. Agar Disk Elution Method for Mycobacterium haemophilum. Susceptibility Testing of Mycobacteria, Nocardia spp., and Other Aerobic Actinomycetes. 2011. 57 p.
15. Wahyudi VA, Seqip P, Sahirah N. Formulasi permen pereda radang tenggorokan dari daun pecut kuda (*Stachytarpheta jamaicensis*) sebagai pangan fungsional. J Pangan dan Agroindustri. 2020;7(4):31-41. <https://doi.org/10.21776/ub.jpa.2019.007.04.4>
16. Sufitri RA, Nurdiana N, Krismayanti L. Uji ekstrak daun pecut kuda (*Stachytarpheta jamaicensis* L) sebagai penghambat bakteri *Staphylococcus aureus*. Biota. 2015;8(2):199-210. <https://doi.org/10.20414/jb.v8i2.69>
17. Ololade ZS, Ogunmola OO, Kuyooro SE, Abiona OO. *Stachytarpheta jamaicensis* leaf extract: Chemical composition, antioxidant, anti-arthritis, anti-inflammatory, and bactericidal potentials. J Sci Innov Res. 2017;6(4):119-25. <https://doi.org/10.31254/jsir.2017.6401>
18. Ramos DF, Leitão GG, Costa FDN, Abreu L, Villarreal JV, Leitão SG, et al. Investigation of the antimycobacterial activity of 36 plant extracts from the Brazilian Atlantic Forest. Rev Bras Ciencias Farm J Pharm Sci. 2008;44(4):669-74. <https://doi.org/10.1590/S1516-93322008000400013>
19. Lelovic N, Mitachi K, Yang J, Lemieux M, Ji Y, Kurosu M. Application of *Mycobacterium smegmatis* as a surrogate to evaluate drug leads against *Mycobacterium tuberculosis*. Physiol Behav. 2020;11(73):780-9. <https://doi.org/10.1038/s41429-020-0320-7>
20. Chatterji D, Dey A. Tracing the variation in physiological response to rifampicin across the microbial spectrum. J Bacteriol Virol. 2012;42(2):87-100. <http://dx.doi.org/10.4167/jbv.2012.42.2.87>
21. Sintha Suhirman. Kandungan Kimia Pecut Kuda. Pros Semin Nas. 2015;(April):93-7. <https://doi.org/10.25181/prosemnas.v0i0.516>
22. Rabaan AA, Alhumaid S, Albayat H, Alsaed M, Alofi FS, Al-Howaidi MH, et al. Promising antimycobacterial activities of flavonoids against *Mycobacterium sp.* drug targets: A comprehensive review. Molecules. 2022;27(5335):1-15. <https://doi.org/10.3390/molecules27165335>
23. Pawar A, Jha P, Chopra M, Chaudhry U, Saluja D. Screening of natural compounds that targets glutamate racemase of *Mycobacterium tuberculosis* reveals the anti-tubercular potential of flavonoids. Sci Rep. 2020;10(1):1-12. <https://doi.org/10.1038/s41598-020-57658-8>
24. Arabski M, Węgierek-Ciuk A, Czerwonka G, Lankoff A, Kaca W. Effects of saponins against clinical *E. coli* strains and eukaryotic cell line. J Biomed Biotechnol. 2012;2012. <https://doi.org/10.1155/2012/286216>
25. Saifudin A. Senyawa alam metabolit sekunder. Edisi 1. Deepublish.
26. Nyambuya T, Mautsa R, Mukanganyama S. Alkaloid extracts from *Combretum zeyheri* inhibit the growth of *Mycobacterium smegmatis*. BMC Complement Altern Med. 2017;17(1):1-11. <https://doi.org/10.1186/s12906-017-1636-0>
27. Wijaya V, Jand'ourek O, Křoustková J, Hradiská-Breiterová K, Korábečný J, Sobolová K, et al. Alkaloids of *Dicranostigma franchetianum* (Papaveraceae) and berberine derivatives as a new class of antimycobacterial agents. Biomolecules. 2022;12(844):1-13. <https://doi.org/10.3390/biom12060844>
28. Cowan MM. Plant products as antimicrobial agents. Clin Microbiol Rev. 1999;12(4):564-82. <https://doi.org/10.1128/CMR.12.4.564>
29. T JAS, J R, Rajan A, Shankar V. Features of the biochemistry of *Mycobacterium smegmatis*, as a possible model for *Mycobacterium tuberculosis*. J Infect Public Health. 2020;13(9):1255-64. <https://doi.org/10.1016/j.jiph.2020.06.023>

30. Handayani FW, Muhtadi A, Farmasi F, Padjadjaran U, Dara T, Manis K, et al. Review artikel: Aktivitas lipofilik protein LysB dari beberapa mikobakteriofaga. *Farmaka Suplemen*. 2013;14(1):1-15. <https://doi.org/10.24198/jf.v16i2.17627.g8706>

Turnitin Artikel Mutiara Medika dr. Ika

ORIGINALITY REPORT

21 %
SIMILARITY INDEX

18 %
INTERNET SOURCES

14 %
PUBLICATIONS

1 %
STUDENT PAPERS

PRIMARY SOURCES

1 repository.lppm.unila.ac.id
Internet Source 4%

2 ijpbs.com
Internet Source 2%

3 worldwidescience.org
Internet Source 1%

4 Maya Dian Rakhmawatie, Tri Wibawa, Puspita Lisdiyanti, Woro Rukmi Pratiwi, Mustofa.
"Evaluation of crystal violet decolorization assay and resazurin microplate assay for antimycobacterial screening", Heliyon, 2019
Publication 1%

5 www.researchgate.net
Internet Source 1%

6 Anthony Swamy Thangiah.
"PHYTOCHEMICAL SCREENING AND ANTIMICROBIAL EVALUATION OF ETHANOLIC-AQUA EXTRACT OF Stachytarpheta jamaicensis (L.) VAHL LEAVES AGAINST SOME

SELECTED HUMAN PATHOGENIC BACTERIA",
Rasayan Journal of Chemistry, 2019

Publication

7	bura.brunel.ac.uk Internet Source	1 %
8	Pia Fyhrquist, Enass Y. A. Salih, Satu Helenius, Into Laakso, Riitta Julkunen-Tiitto. "HPLC-DAD and UHPLC/QTOF-MS Analysis of Polyphenols in Extracts of the African Species Combretum padoides, C. zeyheri and C. psidioides Related to Their Antimycobacterial Activity", Antibiotics, 2020 Publication	1 %
9	journal.umy.ac.id Internet Source	1 %
10	www.mdpi.com Internet Source	1 %
11	Peter Masoko, Maano V. Masiphephethu. "Phytochemical Investigation, Antioxidant and Antimycobacterial Activities of (Lam) Thell Extracts Against ", Journal of Evidence-Based Integrative Medicine, 2019 Publication	1 %
12	link.springer.com Internet Source	1 %
13	www.nature.com Internet Source	1 %

14

Afzal Hussain, Sandeep Kumar Singh, Priya Ranjan Prasad Verma, Neeru Singh, Farhan Jalees Ahmad. " Experimental design-based optimization of lipid nanocarrier as delivery system against species: and evaluation ", Pharmaceutical Development and Technology, 2016

Publication

<1 %

15

Akane Kitayama, Calvin Kurnia, Vinna Kurniawati Sugiaman. "PERANAN AIR PERASAN JERUK NIPIS (Citrus aurantifolia) SEBAGAI ANTIMIKROBA TERHADAP *Prevotella intermedia* PADA PENCEGAHAN PERIIMPLANTITIS (SECARA IN-VITRO)", B-Dent: Jurnal Kedokteran Gigi Universitas Baiturrahmah, 2022

Publication

<1 %

16

Risyandi Anwar, Puspita Hajardhini. "Antibacterial Activity of Gallic Acid from the Leaves of *Altingia excelsa* Noronha to *Enterococcus faecalis*", Open Access Macedonian Journal of Medical Sciences, 2022

Publication

<1 %

17

Enass Y.A. Salih, Riitta Julkunen-Tiitto, Olavi Luukkanen, Mustafa K.M. Fahmi, Pia Fyhrquist. "Hydrolyzable tannins (ellagitannins), flavonoids, pentacyclic triterpenes and their glycosides in

<1 %

antimycobacterial extracts of the ethnopharmacologically selected Sudanese medicinal plant *Combretum hartmannianum* Schweinf", *Biomedicine & Pharmacotherapy*, 2021

Publication

18

academic.oup.com

Internet Source

<1 %

19

e-journals.unmul.ac.id

Internet Source

<1 %

20

ojs.iik.ac.id

Internet Source

<1 %

21

www.esrnexus.com

Internet Source

<1 %

22

Arga Setyo Adji, Nabila Atika, Yemima Billyana Kusbijantoro, Atiyatum Billah, Astrid Putri, Fitri Handajani. "A review of Leaves and Seeds *Moringa oleifera* Extract: The potential *Moringa oleifera* as Antibacterial, Anti-Inflammatory, Antidiarrhoeal, And Antiulcer Approaches To Bacterial Gastroenteritis", *Open Access Macedonian Journal of Medical Sciences*, 2022

Publication

<1 %

23

I Nyoman Jirna, I Gede Sudarmanto, Surya Bayu Kurniawan, Gusti Ayu Made Ratih, Burhannuddin Rasyid. "The potential of

<1 %

traditional balinese spices against the growth of Salmonella sp in vitro", Jurnal Teknologi Laboratorium, 2020

Publication

24

digilibadmin.unismuh.ac.id

Internet Source

<1 %

25

ijisrt.com

Internet Source

<1 %

26

repository2.unw.ac.id

Internet Source

<1 %

27

André C. Pereira, Beatriz Ramos, Ana C. Reis, Mónica V. Cunha. "Non-Tuberculous Mycobacteria: Molecular and Physiological Bases of Virulence and Adaptation to Ecological Niches", Microorganisms, 2020

Publication

<1 %

28

Armando Castañeda, Sergio Rajsbaum, Michel Raynal. "A Linearizability-based Hierarchy for Concurrent Specifications", Communications of the ACM, 2022

Publication

<1 %

29

C.J Rodgers. "Resistance of Yersinia ruckeri to antimicrobial agents in vitro", Aquaculture, 2001

Publication

<1 %

30

Tafadzwa Nyambuya, Ruvimbo Mautsa, Stanley Mukanganyama. "Alkaloid extracts

<1 %

from Combretum zeyheri inhibit the growth of Mycobacterium smegmatis", BMC Complementary and Alternative Medicine, 2017

Publication

31

Theo Audi Yanto, Mochammad Hatta, Agussalim Bukhari, Rosdiana Natzir.

"Molecular and Immunological Mechanisms of Miana Leaf (Coleus Scutellarioides [L] Benth) in Infectious Diseases", Biomedical and Pharmacology Journal, 2020

Publication

<1 %

Exclude quotes Off

Exclude matches Off

Exclude bibliography On