

Kalanchoe 2

by Risyandi 2 Anwar

Submission date: 30-Mar-2023 11:06PM (UTC+0800)

Submission ID: 2051071389

File name: Last_revision_KT.docx (192.38K)

Word count: 2822

Character count: 15814

•

Antimicrobial activity of β -sitosterol isolated from *Kalanchoe tomentosa* leaves against *Staphylococcus aureus* and *Klebsiella pneumonia*

Risyandi Anwar¹, Susi Sukmasari², Lilis Siti Aisyah³, Faridia Puspita Lestari⁴, Delia Ilfani⁵, Yenny Febriani Yun⁶, Pebrian Diki Prestya⁷

¹Paediatric Dentistry Department, Universitas Muhamadiyah Semarang, Indonesia.

² Paediatric Dentistry Department, Kuliyah of Dentistry, International Islamic University Malaysia.

^{3,4,5,6} Chemistry Study, Faculty of Science and Informatics, Universitas Jendral Ahmadyani, Bandung Indonesia.

⁷Oral Surgery Department, Universitas Muhamadiyah Semarang, Indonesia.

Corresponding author: Susi Sukmasari, email: sukmasari@iium.edu.my

<https://orcid.org/0000-0002-1955-0681>

LiveDNA: <https://livedna.net/?dna=62.37515>

Running title: Antimicrobial activity of β -sitosterol isolated from *Kalanchoe tomentosa*

Conflict of interest: The authors declare no conflict of interest

Author's Contribution: RA and PDP are conceptualizer, LSA, FPL, DI and YF as conducted experiments and data analysis. SS led the writing manuscript. All authors contributed to manuscript revision.

ABSTRACT

Background and Objective: Fresh *K. tomentosa* leaves obtained from Bandung, Indonesia was extracted using n-hexane followed by serial dichloromethane maceration. **Material and Method:** N-hexane and ethyl acetate were used to separate the dichloromethane extract using Vacuum Liquid Chromatography and the isolated compounds were recrystallized with n-hexane. **Result:** 37 mg of dichloromethane extract was obtained from the extraction process. Recrystallized compound isolates were identified as stigmast-5-en-3 β -ol or β -sitosterol. Both dichloromethane extract and β -sitosterol isolated compounds showed strong bacteriostatic activity against *S. aureus* with MIC=15.63 μ g/mL and 7.81 μ g/mL, and *K. pneumonia* with MIC=7.81 μ g/mL and 31.25 μ g/mL, respectively. However,

Commented [User1]: Write corresponding author name and registered LiveDNA number of the author.

If author is not registered then register him by following the link below:

Link for Science Alert registration:

<https://www.scialert.com/register.php>

Link for LiveDNA:

<https://livedna.net/>

Commented [User2]: •The running title of the study is missing.

•The author has to provide a running title of 7-9 words besides the main title in the manuscript and it must be the punch line of the main title.

Commented [User3]: •In the article, the authors must include a draft statement that discloses all relevant conflicts of interest and positions.

•Conflict of interest is a situation in which authors may be involved in multiple interests, financial or otherwise, and serving one interest could involve working against another.

•Clarify properly if there any conflict of interest?

Commented [User4]: The author's contribution is missing. Briefly describe each author contribution in this article.

only dichloromethane extract exhibited a bactericidal effect (7.81 µg/mL). **Conclusion:** The pure β-sitosterol compound was isolated from *K. tomentosa* dichloromethane extract. Both the dichloromethane extract and the isolated β-sitosterol compound had antibacterial effects against *S. aureus* and *K. pneumoniae*.

Keywords: *Kalanchoe tomentosa*, β-sitosterol, Oral infection, Minimum Inhibition Concentration, Minimum Bactericidal Concentration

Introduction

Oral cavity infections are often caused by both aerobic and anaerobic bacteria. Odontogenic infections such as osteomyelitis can cause symptoms such as oedema, pain, lymphadenitis, fever, cellulitis, and trismus^{1,2}.

Staphylococcus aureus is a bacterial pathogen responsible for a wide range of infections³. It can survive prolonged extreme conditions to cause infection, it causes abscesses when neutrophils enter the infection site in the invasion stage and can directly invade the lymph vessels and blood to produce bacteriotoxins that cause severe infection^{4,5}. Invasion of *S. aureus* into the heart tissue can cause acute endocarditis. *S. aureus* strains are resistant to several antibiotics including methicillin, nafcillin, and cephalosporins. These strains are known as methicillin-resistant *S. aureus* (MRSA)⁶. The presence of *S. aureus* is confirmed via blood culture where whitish-gold colonies are indicative of the pathogen. A clear halo around the bacterial colony is characteristic of this bacterium's release of haemolysin A toxin which ruptures red blood cells⁷.

Klebsiella pneumoniae is a gram-negative bacterium. Barlean et al. reported *K. pneumoniae* and *S. aureus* as the most commonly identified pathogens in surgical site infections in oral and maxillofacial surgery patients⁸. As with *S. aureus*, *K. pneumoniae* also exhibits antibiotic resistance^{9,10}. Both bacteria are present in the oral cavity and have been reported in postoperative head and neck wound infections. Adequate antibacterial administration is needed to treat bacterial infections. However, increased resistance with widespread antibacterial use has become a major public health issue worldwide. Therefore, it is necessary to search for novel antibacterial agents to treat infections without causing resistance. Natural ingredients have been a promising target for new antibacterial agents with fewer side effects^{4,8}.

Kalanchoe is a large genus of colourful succulent plants. It grows widely in Africa, Saudi Arabia, Asia, the Americas and Australia^{11,12}. Some species of *Kalanchoe* are used as traditional medicinal plants. *Kalanchoe tomentosa* is part of the family Crassulaceae; it is a succulent plant with dense, white, hair-like covering. It is also commonly known as the Panda plant. *K. tomentosa* is rich in alkaloids,

Commented [User5]: In the complete article, the author is advised to use the superscript form while citing references.
•References must be cited in the text in **superscript digits** at the end of sentence or paragraph before punctuation or full stop¹.
•In the case of two or more references, separate the superscript digits by **comma**^{2,4}.
•If there are more references but in continuous sequence then use (-) **dash** between superscript digits^{4,8}.

Commented [SS6R5]: Done change citation

triterpenes, glycosides, flavonoids, steroids and lipids.¹²⁻¹⁴. Ethanolic extracts of *K. tomentosa* (Crassulaceae) has been reported to contain 14 compounds including α -amyrin acetate, friedelin, glutinol, 1-dotriacontanol, phytol, stigmasta-7,25-dien-3 β -ol, β -sitosterol, isorhamnetin, 2,3-dihydroxypropyl tetradecanoate, eriodictyol, gallic acid, quercetin, kampferol-3-O-Rutinoside and isovitexin¹². The flavonoid profile of *K. tomentosa* increases cytotoxic activity against P-388 murine leukaemia cells¹³. However, literature reporting the antibacterial activity of *K. tomentosa* remains limited.

β -sitosterol is a bioactive phytosterol that is naturally derived from plant cell membranes¹⁵. It has been reported to exhibit antibacterial activity against *S. aureus* and *Escherichia coli*. Furthermore, it has been reported to show interesting anti-inflammatory and wound healing effects¹⁵. Pneumolysin is a toxin released by *S. pneumolysin* that is not targeted by currently available antibiotics, making it an interesting target for the development of therapeutics against this pathogen¹⁶. The phytosterol β -sitosterol has been shown to effectively protect against pneumolysin-induced cell lysis. β -sitosterol interacts with the toxin at Thr459 and Leu460.

This *in vitro* study aims to determine the potential of *K. tomentosa* extracts against *S. aureus* and *K. pneumoniae* bacteria, compared against β -sitosterol standards.

Materials and Methods

Study area:

The study conducted at Faculty of Sains and Informatika, Chemistry Department, Chemistry Lab., Jenderal Ahmad Yani University, Cimahi, Indonesia from 2019-2020.

Chemical and equipment:

A set of maceration and KCV. Aqueous sterile, acetone (CH₃COCH₃), ethyl acetate (CH₃COOC₂H₅), dichloromethane (CH₂Cl₂), methanol (CH₃OH), n-hexane (C₆H₁₄), chloroform (CHCl₃) p.a, Thin Layer Chromatography (TLC) plate F254 Merck Germany, Silica gel Merck 60 (0.2- 0.5 mm) Germany, Silica gel Merck 60 G Ultraviolet lamps; brand Vilber Lourmat VL-8. LC, rotary evaporator brand Heidolph Laborota 4000, a set of vacuum liquid chromatography (VLC), Ultraviolet spectrophotometer; Hewlett Packard 8453, Infrared spectrophotometer; Shimadzu Type Prestige-21, spectrometer; NMR ¹H brand JEOL Type JNM-ECA 500 MHz methods were used to elucidate the structures of β -sitosterol.

Plant Collection, Extraction and Isolation of compounds

K. tomentosa leaves were obtained from a nursery at Lembang, Bandung, West Java, Indonesia (6.8145°S latitude, 107.6230°E longitude, 2.254ft elevation). The plant was transported to our

Commented [User7]: •The author is advised to describe when and where the study was carried out? (year, location of study)
•(The study was carried out at Microbiology Department, Quality Control Lab, Egypt from January 2018 to March 2019).

Commented [User8]: Provide manufacturer's name and location (including city, state, and country).

laboratory in Bandung city and was identified by an expert before being cleaned and cut into smaller pieces.

20 kg of fresh *K. tomentosa* leaves were ground and extracted via maceration using n-hexane for 24 hours. Maceration was repeated until the extract was colourless (indicated using thin-layer chromatography (TLC)). The extract was filtered and re-extracted using methylene chloride for 24 hours. The final extract was filtered and concentrated using a rotary evaporator. A solid dark green methylene chloride extract (MCE) was separated using Vacuum Liquid Chromatography (VLC) (silica gel G60) with n-hexane-EtOAc solvent. TLC using n-hexane-EtOAc was performed on the crystals chosen to determine their purity and compare against pure β -sitosterol compounds.

Antibacterial Activity Test

100 μ L nutrient broth (NB) was added into the first column of a 96- well plate as the negative control. 5 μ L of *S. aureus* and *K. pneumonia* bacterial suspensions was added into 10 mL NB and vortexed to mix. 100 μ L of the bacterial suspension was added to columns 2-12. 100 μ L of MCE5 was added to each well and pipetted to mix. Next, 100 μ L was taken from the second column and serially diluted across the 3rd column until the 12th column. The plate was incubated at 37°C for 24 hours, and then the clear wells were observed. The lowest concentration where no microbial growth was detected was defined as the Minimum Bactericidal Concentration (MBC). 5 μ L of solution from the clear wells were transferred into nutrient agar (NA) and incubated at 37°C for 24 hours. The lowest concentration where no microbial growth was observed was defined as Minimum Inhibitory Concentration (MIC).

Result

Extraction and Isolation of compounds:

The final extract was filtered and concentrated using a rotary evaporator to yield 35mg of a solid dark green methylene chloride extract (MCE), then separation using VLC with n-hexane-EtOAc solvent obtained 10 fractions. The fifth fraction (MCE5) produced 8mg of white needle crystals when re-crystallized with n-hexane. The MCE5 was determined the purity and compared against pure β -sitosterol compounds.

UV and TLC analysis:

UV spectroscopy analysis of MCE5 was detected at 264 nm and 364 nm. TLC analysis of the MCE5 against the β -sitosterol standard is shown in **Figure 1**.



Figure 1. TLC analysis of MCE5 (left) against β -sitosterol standard (right). Both showed similar Rf values.

¹HNMR and IR analysis

Analysis of the first isolate using ¹HNMR (500 MHz, CDCl₃) is shown in Table 1. Only one signal was visible above δ H 5ppm, and visible signals accumulated in areas below δ H 2ppm.

Table 1. ¹HNMR data comparison between *K. tomentosa* MCE5 and β -sitosterol

Position of atom C	<i>K. tomentosa</i> MCE5 ¹ HNMR δ H (ppm) (Σ H:mult : J=Hz)	β -sitosterol*
1	0.82 : 0.86 (2H, dd, 10.5 : 5.5)	1.07 : 1.02 (2H, dd, 10.5 : 5.5)
2	1.44 : 1.47 (2H, td, 9.5 : 6.0)	1.44 : 1.48 (2H, td, 9.5 : 6.0)
3	3.49 : 3.55 (1H, m)	3.51 (1H, m)
4	2.23 : 2.31 (2H, m)	2.22 : 2.29 (2H, m)
5	-	-
6	5.35 (1H, t, 2.5)	5.35 (1H, br)
7	2.01 (2H, dt, 5.6 : 8.5)	1.85 : 2.01 (2H, dt, 5.6 : 8.5)
8	1.60 (1H, m)	1.57 (1H, m)
9	0.93 (1H, m)	0.93 (1H, m)
10	-	-
11	1.42 : 1.47 (2H, m)	1.42 : 1.49 (2H, m)
12	1.95 (2H, d, 5.6)	1.15 : 1.98 (2H, d, 5.6)
13	-	-
14	1.01 (1H, m)	1.00 (1H, m)

Commented [User9]: Author is advised to recheck the sign use in the table as we think decimal point must use here rather than inverted commas. Kindly use the correct sign in all values of the tables.

15	1.94 : 2.03 (2H, m) ¹³	1.57 (2H, m)
16	1.95 (2H, m)	1.84 (2H, m)
17	1.09 (1H, dt, 5.2 : 8.5) ⁹	1.09 (1H, dt, 5.2 : 8.5)
18	0.68 (3H, s)	0.68 (3H, s)
19	1.01 (3H, s)	1.01 (3H, s)
20	1.31 (1H, m)	1.36 (1H, m) ⁵
21	0.69 (3H, d : 6.1)	0.92 (3H, d, 6.1)
22	1.80 (2H, m)	1.38 (2H, m)
23	1.50 (2H, m) ⁵	1.54 (2H, m)
24	0.94 : 0.98 (1H, m)	0.93 (1H, m)
25	1.66 (1H, m) ⁷	1.66 (1H, m)
26	0.84 (3H, d, 6.2)	0.84 (3H, d, 6.2)
27	0.92 (3H, d, 6.7)	0.92 (3H, d, 6.7)
28	1.25 (2H, m)	1.26 (2H, m)
29	0.82 (3H, s)	0.83 (3H, s)

The various functional groups in MCE5 were elucidated using infrared (IR) spectrophotometry. The results of IR spectral data of MCE5 show absorption in the following areas: 3415 cm^{-1} area with widening intensity. Absorption at 1454 cm^{-1} and 1371 cm^{-1} with a sharp intensity indicated a C-H function group in CH₂ and CH₃ as shown in Table 2.

Table 2. IR data comparison between *K. tomentosa* MCE5 and β -sitosterol

Isolate 1 (cm^{-1})	β -sitosterol* (cm^{-1})	Creswell** (cm^{-1})	Functional Group
3415.93	3440.62	3450-3200	O-H
2945.30	2936.69	2800-3000	C-H aliphatic
1649.14	1646.55	1680-1620	C=C
1454.33	1463.42	1475-1300	C-H (in CH ₂)
1371.39	1381.65	1475-1300	C-H (in CH ₃)
1047.35	1053.89	1050-1260	C-O alcohol
962.48	970.32	995-710	=CH alkene

**Creswell dkk., 1982.

Commented [User10]: Author is advised to recheck the sign use in the table as we think decimal point must use here rather than inverted commas. Kindly use the correct sign in all values of the tables.

* Fitriani, 2013.

Antimicrobial Activity

The MIC values of MCE5 against *S. aureus* and *K. pneumonia* were 15.63 $\mu\text{g/mL}$ and 7.81 $\mu\text{g/mL}$, respectively. The MBC values are 7.81 $\mu\text{g/mL}$ as shown in Table 3.

Table 3. Antimicrobial activity of MCE5 and β -sitosterol against *S. aureus* and *K. tomentosa*

	MCE5		β -sitosterol	
	MIC ($\mu\text{g/mL}$)	MBC ($\mu\text{g/mL}$)	MIC ($\mu\text{g/mL}$)	MBC ($\mu\text{g/mL}$)
<i>S. aureus</i>	15.63	-	7.81	-
<i>K. pneumonia</i>	7.81	7.81	31.25	-

Commented [User11]: Author is advised to recheck the sign use in the table as we think decimal point must use here rather than inverted commas. Kindly use the correct sign in all values of the tables.

Discussion

UV spectroscopy analysis of MCE5 indicated strong absorption at 264nm and weak absorption at 364nm. The values indicate the presence of an unconjugated alkene system and the absence of aromatic systems. TLC analysis of the MCE5 against the β -sitosterol standard showed similar Rf values (Figure 1). Based on the values, it was predicted that MCE5 to be β -sitosterol. The above result is reinforced by data from ^1H NMR analysis. Analysis of the first isolate using ^1H NMR (500 MHz, CDCl_3) proved that the isolate was a steroid compound, as shown in Table 1. There was a distinctive oleophilic proton signal at δH 5ppm, and an oxygenated proton signal at δH 3ppm, which is commonly found in the steroid groups¹⁷.

In addition, a chemical shift in δH 1.42ppm was characteristic of a cyclohexane group in A, B and C in steroid compounds¹⁸. Proton signals found at δH 0.86 (2H, dd); 1.44 (2H, dt); 2.23 (2H, m); 2.01 (2H, dt); 1.42 (2H, m); 1.99 (2H, d); 1.94 (2H, m); 1.95 (2H, m) are signals for methylene protons

(CH₂). Proton signals at δ H 3.49(1H, m); 5.35(1H, t); 1.60(1H, m); 0.93(1H, m); 1.01(1H, m); 1.09 ppm (1H, dt) indicate the methine proton, δ H 3.49(1H, m) indicates the presence of hydrogen adjacent to the hydroxyl group suspected in C-3 in ring A present next to the molecular plane, on the same side as the methyl groups in C-10 and C-13. This configuration is recognized as β -configuration. A proton signal at δ H 5.35 (1H, t) shows oleophilic methane, indicating a double bond in C-5¹⁷. Each proton signal at δ H 0.68(3H, s) and 1.01 (3H, s) correspond to methyl groups as steroid substituent in the main framework at C-10 and C-13, respectively. These three signals (methylene, methine and methyl) indicate the presence of a steroid framework¹⁹ substituted by two methyl and one hydroxyl group.

Signals were observed in the aliphatic region (substituent at C-17) that indicate an alkane unit: three signals for the methylene group at δ H 1.80(2H, m); 1.50(2H, m) and 1.25 (2H, m), three signals for the methine group at δ H 1.31(1H, m); 0.94(1H, m); 1.66(1H, m), and four methyl group signals at δ H 0.69 (3H, d); 0.84(3H, d); 0.92(3H, d) and 0.82(3H, d). These ten proton signals correspond to alkyl skeletons²⁰ All the shifts were compared against β -sitosterol 10 compounds and many similarities were found. It was therefore concluded that MCE5 was a β -sitosterol compound.

The various functional groups in MCE5 were elucidated using infrared (IR) spectrophotometry. The results of IR spectral data of MCE5 show absorption in the following areas: 3415cm⁻¹ area with widening intensity, indicating alcohol group (-OH). This is reinforced by absorption at 1047cm⁻¹ with sharp intensity indicating alcohol(C-O), absorption at 2945 cm⁻¹ with sharp intensity indicating aliphatic functional group (CH), absorption at 1649 cm⁻¹ with widening intensity indicative of carbon group (C=C), absorption at 835 cm⁻¹ with moderate intensity indicating alkenes (=CH)²¹. According to Shaleh (2007), absorption at 1649 cm⁻¹ indicates the existence of a range of C=C functional groups with no conjugation that has a long-range wave of between 1620-1680 cm⁻¹¹².

The IR spectral data results reinforce that the isolates obtained are a group of steroid compounds that do not have conjugated double bonds. Absorption at 1454 cm⁻¹ and 1371 cm⁻¹ with a sharp intensity indicated a C-H function group in CH₂ and CH₃. The MCE5 IR spectral data results were compared against pure β -sitosterol compounds. Similar absorption patterns were identified, as shown in **Table 2**.

Results of TLC, ¹HNMR and IR analysis of MCE5 indicate that MCE5 from *K. tomentosa* may be a β -sitosterol compound, specifically, compound IUPAC stigmast-5-en-3 β -ol that belongs to the group stigmata in originating hydrocarbons steroid with the molecular formula C₂₉H₅₀O. ~~The molecular structure of this compound is shown in Figure 2.~~

MCE5 had bacteriostatic effects against *S. aureus* and *K. pneumonia* with MIC values of 15.63 µg/mL and 7.81 µg/mL, respectively. MCE did not have a bactericidal effect against *S. aureus*, but did against *K. pneumonia* with an MBC of 7.81 µg/mL. Therefore, MCE5 had strong activity against *S. aureus* and *K. pneumonia*, as shown in **Table 3**.

The MIC data show *K. pneumonia* required lower doses of MCE5 and β-sitosterol compared to *S. aureus* for effective antibacterial activity. Gram-positive and gram-negative bacteria have differences in their cell wall structures, which affect their susceptibility to antibacterial agents. Gram-positive bacterial cell walls have a single-layered peptidoglycan structure that is polar and has low lipid content (Pelczar). They also contain polysaccharides, which serve as positive ion transfers in and out of the cell. Hence, gram-positive bacterial cell walls are more polar and more susceptible to antibacterial agents.

MCE5 had one hydroxyl group (-OH) in its structure. The polar -OH group can penetrate the polar peptide and damage the bacterial cell wall by severing the peptidoglycan bonds to compromise the cell layer. This leaves the cytoplasmic membrane vulnerable to damage, causing the leak of important metabolites and activation of the bacterial enzyme system. Antibacterials target the peptidoglycan layer of cell walls in the bacteria. This layer is essential in preserving the bacteria from hypotonic environments, hence damage or loss of this layer will lead to loss of cell wall strength, resulting in death²³. Further research is needed to identify the other secondary metabolite compounds present in *K. tomentosa*, as well as elucidation of the structure of the compound via analysis using MS, CNMR, and two- and three-dimensional NMR. Isolation and identification of pure β-sitosterol compounds in the fifth fraction of the methyl chloride extract from the leaves of *K. tomentosa* that conferred bacteriostatic effects against *S. aureus* and *K. pneumonia*.

Conclusion

The pure β-sitosterol compound was isolated from *K. tomentosa* dichloromethane extract. Both the dichloromethane extract and the isolated β-sitosterol compound had antibacterial effects against *S. aureus* and *K. pneumonia*.


Significance Statement

This study discovers the β-sitosterol compound was isolated from *K. tomentosa* dichloromethane extract as one option of an antimicrobial drug discovery. This study will help the researcher to hone the Kalanchoe genus elucidation using different method of extraction that there was still very few papers on it. Thus, possibly other advantages combinations of the Kalanchoe, may be arrived at.

Commented [User12]: •The author is advised to state the significance of his research work after the conclusion section. The significance statement should provide the novelty aspect and significance of his research work for the existing literature and more generally to the society.

- It should be a summary that describes what this paper adds to and what was already known.
- Start this statement with the following words: This study discovers the ----- that can be beneficial for -----
- And the last sentence of this statement could be such as This study will help the researcher to uncover the critical areas of ----- that many researchers were not able to explore. Thus a new theory on ----- may be arrived at.
- A Model Significance Statement: This study discovers the possible synergistic effect of vitamin E, calcium, and vitamin D combination that can be beneficial for osteoporosis-induced ovariectomized rats. This study will help the researcher to uncover the critical area of postmenopausal bone loss that many researchers were not able to explore. Thus, a new theory on these micronutrients combination, and possibly other combinations, may be arrived at.

Commented [SS13R12]: Has been clarified



Kalanchoe 2

ORIGINALITY REPORT

10 %	%	%	10 %
SIMILARITY INDEX	INTERNET SOURCES	PUBLICATIONS	STUDENT PAPERS

PRIMARY SOURCES

1	Submitted to UC, Irvine Student Paper	3 %
2	Submitted to Universitas Negeri Semarang Student Paper	1 %
3	Submitted to Campbellville University Student Paper	1 %
4	Submitted to University of KwaZulu-Natal Student Paper	1 %
5	Submitted to Mae Fah Luang University Student Paper	1 %
6	Submitted to University of Newcastle upon Tyne Student Paper	<1 %
7	Submitted to Mahidol University Student Paper	<1 %
8	Submitted to Australian Catholic University Student Paper	<1 %
9	Submitted to Hanoi National University Student Paper	<1 %

10	Submitted to Mugla University Student Paper	<1 %
11	Submitted to Queen's University of Belfast Student Paper	<1 %
12	Submitted to Universitas Mulawarman Student Paper	<1 %
13	Submitted to Higher Education Commission Pakistan Student Paper	<1 %
14	Submitted to Universiti Sains Malaysia Student Paper	<1 %
15	Submitted to University of Queensland Student Paper	<1 %
16	Submitted to Nanyang Technological University, Singapore Student Paper	<1 %

Exclude quotes Off

Exclude matches Off

Exclude bibliography Off

Kalanchoe 2

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8

PAGE 9

PAGE 10
