

**JUDUL ARTIKEL:** Case Report Preeclamsia In Pregnant Women In Work Area  
Kismantoro Health Center Wonogiri With Dash Method

**E.ISBN:** 978-623-6974-71-1

Publish: The 2nd International Conference on Practice and Innovation Midwifery (The 2nd ICPIM)  
August 18th, 2022 – P.ISBN : 978-623-6974-70-4 / E.ISBN : 978-623-6974-71-1

Publisher: <https://icpim.unimus.ac.id/>

URL: <https://prosiding.unimus.ac.id/index.php/icpim/article/viewFile/1350/1355>

Tanggal Publisher: 18 Agustus 2022

**Urutan file ini sebagai berikut:**

1. Riwayat Submit
2. Manuskrip yang disubmit
3. Riwayat Review/review substansi
4. Manuskrip setelah revisi
5. In Press
6. Artikel sudah publish

## 1. RIWAYAT SUBMIT



## 2. MANUSKRIP YANG DISUBMIT

### LAPORAN KASUS PREEKLAMPSIA RINGAN PADA IBU HAMIL DI WILAYAH KERJA PUSKESMAS KISMANTORO WONOGIRI DENGAN METODE DASH

Izzah Al Fawaidah<sup>1</sup>, Indri Astuti Purwanti<sup>2</sup>, Novita Nining Angraeni<sup>3</sup>, Fitriani Nur Damayanti<sup>4</sup>  
<sup>1,2,3</sup> Program Studi Pendidikan Profesi Bidan, Universitas Muhammadiyah Semarang  
<sup>4</sup>Corresponding author: izzahalfawaidah@gmail.com

#### ABSTRAK

**Pendahuluan:** Preeklamsia merupakan salah satu penyebab tingginya angka kematian ibu dilihat dalam skala nasional maupun daerah dan biasanya ditandai dengan adanya gejala seperti hipertensi, edema, proteinuria. Pada tahun 2021 terdapat 23 kasus preeklamsia dan 11 kasus dengan preeklamsia yang dirujuk. Hasil wawancara pada 2 ibu yang mengalami preeklamsia penyebabnya adalah riwayat hipertensi sebelum hamil dan umur ibu > 35 tahun dengan multipara. DASH (Dietary Approaches to Stop Hypertension) adalah metode diet yang efektif untuk penderita hipertensi. Pendekatan diet DASH dapat dijadikan sebagai terapi non-farmakologis dan sebagai pengganti obat. **Presentasi Kasus:** hasil pengkajian menunjukkan bahwa partisipan memiliki keluhan pusing dan nyeri ulu hati sudah 3 hari, ibu hamil pada usia > 35 tahun dan mempunyai Riwayat darah tinggi sebelum hamil. Data obyektif menunjukkan hasil tekanan darah yang tinggi 150/100, oedema pada kaki kanan dan kiri, protein urine positif. Diagnosa pada kasus ini adalah preeklamsia dengan penanganan awal adalah kolaborasi dengan dokter untuk pemeriksaan dan penanganan lebih lanjut serta menganjurkan partisipan menerapkan pola makan dengan diet DASH setiap hari di rumah. Catatan perkembangan pada minggu pertama, partisipan sudah tidak pusing dan nyeri ulu hati serta oedema berkurang. Hasil minggu kedua, partisipan tidak mengalami oedema dan tekanan darah menurun. **Kesimpulan:** Penanganan preeklamsia dengan metode diet DASH dapat menurunkan tekanan darah bila diterapkan setiap hari dengan benar.

**Kata Kunci:** Preeklamsia, DASH, Kehamilan

### *PREECLAMPSIA CASE REPORT IN PREGNANT WOMEN IN WORK AREA KISMANTORO HEALTH CENTER WONOGIRI WITH DASH METHOD*

Izzah Al Fawaidah<sup>1</sup>, Indri Astuti Purwanti<sup>2</sup>, Novita Nining Angraeni<sup>3</sup>, Fitriani Nur Damayanti<sup>4</sup>  
<sup>1,2,3,4</sup> Midwifery Professional Education Study Program, University of Muhammadiyah Semarang

\*Corresponding author: [izzahalfawaidah@gmail.com](mailto:izzahalfawaidah@gmail.com)

### ABSTRACT

**Introduction:** Preeclampsia is one of the causes of the high maternal mortality rate seen on a national and regional scale and is usually characterized by symptoms such as hypertension, edema, proteinuria. In 2021 there were 23 cases of preeclampsia and 11 cases with preeclampsia referred. The results of interviews with 2 mothers who experienced preeclampsia were a history of hypertension before pregnancy and the mother's age was > 35 years with multipara. DASH (Dietary Approaches to Stop Hypertension) is an effective diet method for people with hypertension. The DASH diet approach can be used as a non-pharmacological therapy and as a substitute for drugs.

**Case Presentation:** the results of the study showed that participants had complaints of dizziness and heartburn for 3 days, pregnant women at the age of > 35 years and had a history of high blood pressure before pregnancy. Objective data showed high blood pressure 150/100, edema in the right and left legs, positive urine protein. The diagnosis in this case was preeclampsia with initial treatment in collaboration with doctors for further examination and treatment and recommending participants to adopt a DASH diet every day at home. Notes of development in the first week, participants were no longer dizzy and heartburn and edema were reduced. The results of the second week, participants did not experience edema and decreased blood pressure. **Conclusion:** Treatment of preeclampsia with the DASH diet method can lower blood pressure when properly applied every day.

**Keywords:** Preeclampsia, DASH, Pregnancy

## 1. Introduction

Preeclampsia is a contributor to the high maternal mortality rate is preeclampsia. Preeclampsia or eclampsia is a complication that is often experienced by pregnant women. Viewed on a national and regional scale, preeclampsia or eclampsia is a major cause of maternal death, namely bleeding, preeclampsia, and infection [Elsanti, 2020].

Preeclampsia is a collection of symptoms that occur in pregnant women, childbirth, and the puerperium consisting of hypertension, edema, and proteinuria that appear at 20 weeks gestation until the end of the first week after delivery. Preeclampsia is a serious problem and has a fairly high level of complexity. The magnitude of this problem is not only because preeclampsia affects the mother during pregnancy and childbirth, but also causes postpartum problems, such as the risk of cardiometabolic disease and other complications [Muzalfah et al., 2018].

According to the World Health Organization (2019), the maternal mortality rate in the world is 303,000 people. The Maternal Mortality Rate in Association of Southeast Asian Nations is 235 per 100,000 live births (ASEAN Secretariat, 2020). According to the Indonesian Demographic and Health Survey Data, the Maternal Mortality Rate in Indonesia increased from 228 per 100,000 live births in 2002-2007 to 359 per 100,000 live births in 2007-2012. MMR decreased in 2012-2015 to 305 per 100,000 live births and the number of

maternal deaths in Indonesia in 2019 was 4,221 cases [Kemenkes RI, 2021].

According to the Ministry of Health in Indonesia, the maternal mortality rate is 1712 per 100,000 live births. Meanwhile, the IMR in 2017 was 10294 per 1,000 live births [Kemenkes RI, 2021].

MMR achievement in 2020 is 98.6/100,000 life birth; IMR: 7.79/1000 life birth and infant mortality rate 8.99/1000 life birth) although this figure is much better than the national target (MMR: 226/100,000 life birth; IMR: 24/1,000 life birth) however, the MMR achievement is lower than the MMR achievement in 2019 (AKI 76.93/100,000 KH; IMR: 8.24/1000 KH and AKABA 9.65/1000 KH [Dinas Kesehatan Provinsi Jawa Tengah, 2021].

The MMR in 2020 is 111.45 per 100,000 live births, this has increased again compared to 2019 which was 73.87 per 100,000 births. The most common causes of maternal death in Wonogiri Regency in 2020 were bleeding 5 people, severe preeclampsia (PEB) 3, broncho pneumonia 1 person, hyperthyroidism 1 person, heart disease 1 person, amniotic fluid embolism 1 person [Dinas Kesehatan Wonogiri, 2021].

The results of the study [Wijaya, 2021] with the title Management of Preeclampsia in Pregnant Women at PMB Siti Azizah Wijaya, S.ST., Bd showed that participants 1 and 2 had complaints of swollen feet accompanied by headaches. On examination of vital signs, participants 1 and 2 showed high blood pressure results. After being

given implementation in the form of observing blood pressure, recommending bed rest, implementing a diet with the DASH diet and soaking the feet using warm water mixed with salt, it was found that participant 1 was resolved at week 3 and participant 2 was resolved at week 4. This occurs because participant 2 is more cooperative than participant 1.

Based on the results of a preliminary study at the Kismantoro Wonogiri Health Center, the number of cases of preeclampsia has increased from 2020 to 2021. In 2020 the number of pregnant women with preeclampsia was 11 cases and preeclampsia were referred as many as 7 cases. Meanwhile, in 2021 there were 23 cases of preeclampsia and 11 cases with preeclampsia who were referred. The results of interviews with 2 mothers who experienced preeclampsia were a history of hypertension before pregnancy and the mother's age was over thirty five years with multipara.

Based on (Hapsari, 2016) One way to control hypertension, especially in pregnancy, is to adjust the diet using the DASH method. Since the publication of the results of the DASH diet research in 1997, the DASH diet has been widely recommended to prevent and treat hypertension. The Dietary Approaches to Stop Hypertension (DASH) diet study states that a diet high in vegetables, fruit, and low-fat dairy products that are low in saturated fat and total fat and high in potassium, calcium, and magnesium can lower systolic blood pressure by 6-11 mmHg, and diastolic blood pressure of 3-6 mmHg [Rahadiyanti et al., 2015] .

The DASH diet approach is not only used as a non-pharmacological therapy but can be used as a substitute for drugs [Fitriyana &

Wirawati, 2022]. To anticipate the occurrence of unwanted things in pregnant women with symptoms that lead to preeclampsia, it is recommended to carry out routine and regular antenatal checks. This is useful for early detection if the mother's body finds signs and symptoms of preeclampsia, so that the mother can immediately get treatment. In addition, patients can also do bed rest early and apply a good diet so that their condition does not worsen and is expected to be more stable than the previous condition.

Therefore, in helping efforts to accelerate the decline in MMR and IMR, one of the authors is implementing continuous care or Continuity of Care. Continuity of Care in midwifery is a series of continuous and comprehensive service activities starting from pregnancy, childbirth, postpartum, newborn care and family planning services that link women's particular health needs and individual circumstances (Homer et al., 2014).

## 2. PATIENT IDENTITY

The patient named Mrs. S, 36 years old. His last education was junior high school, he worked to take care of the household, address Gesing Kismantoro.

The main complaints are dizziness and heartburn for 3 days. Pregnant with 2nd child and never miscarried. History of disease, namely having had hypertension in the first pregnancy. HPHT: 09-11-2021

## 3. CLINICAL FINDINGS

Good general condition, Compos mentis conscious, Blood pressure: 150/100mmHg, Respiration: 20 x/min, Pulse: 80 x/min, Temperature: 36.5oC, TFU: 28 cm, head presentation, convergent, contractions: none, FHR : 142 x/minute, right and left lower extremity edema.

## 4. TIMELINE

Tanggal dan jam	Subjektif	Objektif	Assessment	Planning
07.07.22 10.00	Dizziness and heartburn for 3 days	Blood pressure: 150/100 mmHg, Respiration: 20 x/minute, Pulse: 80 x/minute, Temperature: 36.5oC, TFU: 28 cm, head presentation, convergent,	Mrs. S. G2P1A0 36 years old 34 weeks pregnant single fetus live intra uteri, longitudinal position,	1. Explain to the mother about the current status of pregnancy 2. Explain to the mother about preeclampsia 3. Observe and ask the mother to sit on the left while sleeping 4. Advise the mother to soak the feet in warm water 5. Give moral support to the mother
				contraction: none, FHR: 142 x/minute, lower extremity edema. cephalic presentation, pucca, convergent with preeclampsia
				6. Give KIE to the mother regarding the management of preeclampsia with the DASH method 7. Collaborate with doctors for further examination 8. Advise mother to visit again in 1 week
				1. Explaining the results of the examination to the mother that the mother's condition has improved such as the mother is no longer dizzy and has heartburn, reduced blood pressure, reduced edema 2. Encourage mothers to continue implementing the DASH diet at home, namely a diet rich in fruits, vegetables, whole grains, nuts, fish, and low-fat milk. These foods are high in important

		convergent, contraction: none, FHR: 142 x/minute, lower extremities slightly edematous	position, cephalic presentation, puca, convergent with preeclampsia	nutrients, such as potassium, magnesium, calcium, fiber, and protein. 3. Advise the mother to rest enough during the day 1-2 hours and 7-8 hours at night 4. Advise the mother to continue to take vitamins from the doctor 5. Advise control mother for 1 more week to monitor general condition and vital sign
<b>21.07.22</b> <b>08.00</b>	Mom has no complaints, just wants to check her pregnancy	Blood pressure: 146/90 mmHg, Respiration: 20 x/minute, Pulse: 80 x/minute, Temperature: 36.5oC, TFU: 29 cm, head presentation, convergent, contraction: none, FHR: 150x/minute, extremities no edema	Mrs. S G2P1A0 36 years of age 36 weeks pregnant, single live intrauterine fetus, longitudinal position, cephalic presentation, puca, convergent with preeclampsia	1. Explain the results of the examination to the mother that her condition has improved, blood pressure has decreased, there is no edema, positive urine protein 2. Encourage mothers to continue implementing the DASH diet at home, namely a diet rich in fruits, vegetables, whole grains, nuts, fish, and low-fat milk. 3. Advise the mother to rest enough during the day 1-2 hours and 7-8 hours at night 4. Giving counseling to mothers with danger signs in pregnancy such as severe dizziness, blurred vision, heartburn, bleeding, seizures 5. Giving counseling to the mother for signs of labor such as urinating in the lower abdomen, removing blood mucus, removing fluid/amniotic fluid from the birth canal 6. Provide counseling to mothers regarding preparation for childbirth 7. Advise the mother to continue to take vitamins from the doctor 8. Instruct the mother for control in 1 week to monitor general condition, vital sign and signs of labor.

## 5. DIAGNOSTIC CHECK

Investigations carried out were urine protein examination with positive results. The diagnosis of this case was preeclampsia. This diagnosis is supported by research results which state that (Peres et al., 2018) Preeclampsia is a hypertensive condition found at gestational age over twenty weeks with or without urine protein and accompanied by other organ disorders.

The prognosis in this case is preterm delivery. This is in line with the theory Norwitz ER, Repke JT in Hidayati et al., 2018 One of the prognosis for preeclampsia in pregnancy is the risk of complications to the mother and fetus, such as impaired fetal growth, preterm birth, placental abruption, and Intra Uterin Fetal Death.

## 6. INTERVENTION THERAPY

The intervention carried out during the visit was to encourage mothers to apply the DASH method, namely a diet high in important nutrients, such as potassium, magnesium, calcium, fiber, and protein. (Nurhayati, 2021). Based on research (Porouw & Yulianingsih, 2019) The content of sodium and potassium in Ambon bananas is proven to be able to reduce high blood pressure in pregnant women when consumed regularly and with the right dose.

The next implementation is to encourage mothers to do bed rest (Novitasari et al., 2018), In addition, it is recommended that mothers soak their feet in warm water (Sabattani et al., 2016).

Management in cases of preeclampsia also provides referrals or recommendations to the hospital for further examination and treatment with the patient's consent.

## 7. FOLLOW UP AND RESULT

Monitoring is carried out every 1 week, by conducting home visits, at the first visit there is a development of the mother's condition, namely complaints of dizziness and heartburn have disappeared, reduced leg edema, there is a decrease in diastolic blood pressure of 5 mm Hg. The second visit showed progress with a decrease in systolic blood pressure of 4 mm Hg and diastolic 5 mm Hg, no edema, urine protein was still positive.

## 8. KESIMPULAN

Treatment of preeclampsia with the DASH method can lower blood pressure if applied every day properly. Collaboration with doctors is also very necessary for further examination and therapy.

## 9. INFORMED CONSENT

Informed consent has been done on July 7, 2022 and the patient is willing to be a respondent

## 10. REFERENCE


- Astuti. (2012). *Buku Ajaran Asuhan Kebidanan Ibu 1 (Kehamilan)*. Yogyakarta: Rahima Press
- Dea Prastika Hapsari, A. R. (2016). *Hubungan Pengetahuan dengan Perilaku Manajemen Hipertensi: Aktivitas Fisik dan Diet DASH Penderita Hipertensi di Desa Salamrejo*. III(2), 2016.
- Elsanti, B. D. (2020). *By Devita Elsanti*.
- Fitriyana, M., & Wirawati, M. K. (2022). Penerapan Pola Diet Dash Terhadap Tekanan Darah Pada Penderita Hipertensi Di Desa Kalikangkung Semarang. *Jurnal Manajemen Asuhan Keperawatan*, 6(1), 17–24.  
<https://doi.org/10.33655/mak.v6i1.126>
- Homer, C. S. E., Friberg, I. K., Dias, M. A. B., ten Hoope-Bender, P., Sandall, J., Speciale, A. M., & Bartlett, L. A. (2014). The projected effect of scaling up midwifery. *Lancet (London, England)*, 384(9948), 1146–1157.  
[https://doi.org/10.1016/S0140-6736\(14\)60790-X](https://doi.org/10.1016/S0140-6736(14)60790-X)
- Muzalfah, R., Santik, Y. D. P., & Wahyuningsih, A. S. (2018). Kejadian Preeklampsia pada Ibu Bersalin. *Higeia Journal Of Public Health Research Development*, 2(3), 1–12.  
<https://journal.unnes.ac.id/sju/index.php/higeia/article/view/21390/11738>
- Novitasari, P., Prasetyorini, H., & Prihati, D. R. (2018). Upaya Pemenuhan Kebutuhan Istirahat Tidur Pada Ibu Postpartum Dengan Preeklampsia. *Jurnal Manajemen Asuhan Keperawatan*, 2(1), 22–33.  
<https://doi.org/10.33655/mak.v2i1.31>
- Nurhayati, S. (2021). Penatalaksanaan Preeklampsia Pada Ibu Hamil. *STIKes Ngudia Husada Madura Oleh*.
- Nurianti, I., Saputri, I. N., & Crisdayanti Sitorus, B. (2021). Hubungan Dukungan Suami Dengan Kecemasan Ibu Hamil Dalam Menghadapi Proses Persalinan. *Jurnal Kebidanan Kestra (Jkk)*, 3(2),

163–169.

<https://doi.org/10.35451/jkk.v3i2.493>

- Peres, G. M., Mariana, M., & Cairrão, E. (2018). Pre-eclampsia and eclampsia: An update on the pharmacological treatment applied in Portugal. *Journal of Cardiovascular Development and Disease*, 5(1).  
<https://doi.org/10.3390/jcdd5010003>
- Porouw, H. S., & Yulianingsih, E. (2019). Pisang Ambon Dan Hipertensi Ibu Hamil. *Jambura Health and Sport Journal*, 1(2), 61–70.  
<https://doi.org/10.37311/jhsj.v1i2.2597>
- Putri, H. (2018). Determinan Kejadian Pre Eklamsi di RSD Kalisat Jember. *Jurnal Kesehatan Dr Soebandi*, 6(2), 21–29.
- Rahadiyanti, A., Setianto, B. Y., & Purba, M. B. (2015). Asupan makan DASH-like diet untuk mencegah risiko hipertensi pada wanita prediabetes. *Jurnal Gizi Klinik Indonesia*, 11(3), 115.  
<https://doi.org/10.22146/ijcn.19290>
- Sabattani et. al. (2016). Efektivitas rendam kaki dengan air hangat terhadap penurunan tekanan darah pada ibu hamil penderita preeklamsi di Puskesmas Ngaliyan Semarang. *Ilmu Keperawatan Dan Keidanan*, 5(1), 1–10.
- Yeyeh, A. R., Yolanda Sari, D., & Humaeroh, D. (2021). Hubungan Karakteristik Ibu Bersalin dengan Preeklampsia Berat di RSUD A Purwakarta Tahun 2020. *Jik*, 16–26.

### 3. RIWAYAT REVIEW/REVIEW SUBSTATANSI



## LAPORAN KASUS PREEKLAMPSIA RINGAN PADA IBU HAMIL DI WILAYAH KERJA PUSKESMAS KISMANTORO WONOGIRI DENGAN METODE DASH

Izzah Al Fawaidah<sup>1</sup>, Indri Astuti Purwanti<sup>2</sup>, Novita Nining Anggraini<sup>3</sup>, Fitriani Nur Damayanti<sup>4</sup>  
<sup>1,2,3,4</sup> Program Studi Pendidikan Profesi Bidan, Universitas Muhammadiyah Semarang  
\*Corresponding author: izzahalfawaidah@gmail.com

### ABSTRAK

**Pendahuluan:** Preeklamsia merupakan salah satu penyebab tingginya angka kematian ibu dilihat dalam skala nasional maupun daerah dan biasanya ditandai dengan adanya gejala seperti hipertensi, edema, proteinuria. Pada tahun 2021 terdapat 23 kasus preeklamsia dan 11 kasus dengan preeklamsia yang dirujuk. Hasil wawancara pada 2 ibu yang mengalami preeklamsia penyebabnya adalah riwayat hipertensi sebelum hamil dan umur ibu > 35 tahun dengan multipara. DASH (Dietary Approaches to Stop Hypertension) adalah metode diet yang efektif untuk penderita hipertensi. Pendekatan diet DASH dapat dijadikan sebagai terapi non-farmakologis dan sebagai pengganti obat. **Presentasi Kasus:** hasil pengkajian menunjukkan bahwa partisipan memiliki keluhan pusing dan nyeri ulu hati sudah 3 hari, ibu hamil pada usia > 35 tahun dan mempunyai Riwayat darah tinggi sebelum hamil. Data obyektif menunjukkan hasil tekanan darah yang tinggi 150/100, oedema pada kaki kanan dan kiri, protein urine positif. Diagnosa pada kasus ini adalah preeklamsia dengan penanganan awal adalah kolaborasi dengan dokter untuk pemeriksaan dan penanganan lebih lanjut serta menganjurkan partisipan menerapkan pola makan dengan diet DASH setiap hari di rumah. Catatan perkembangan pada minggu pertama, partisipan sudah tidak pusing dan nyeri ulu hati serta oedema berkurang. Hasil minggu kedua, partisipan tidak mengalami oedema dan tekanan darah menurun. **Kesimpulan:** Penanganan preeklamsia dengan metode diet DASH dapat menurunkan tekanan darah bila diterapkan setiap hari dengan benar.

**Kata Kunci:** Preeklamsia, DASH, Kehamilan

Commented [R1]: Delete text

## *PREECLAMPSIA CASE REPORT IN PREGNANT WOMEN IN WORK AREA KISMANTORO HEALTH CENTER WONOGIRI WITH DASH METHOD*

Izzah Al Fawaidah<sup>1</sup>, Indri Astuti Purwanti<sup>2</sup>, Novita Nining Anggraini<sup>3</sup>, Fitriani Nur Damayanti<sup>4</sup>  
<sup>1,2,3,4</sup> Midwifery Professional Education Study Program, University of Muhammadiyah Semarang



\*Corresponding author: [izzahalfawaidah@gmail.com](mailto:izzahalfawaidah@gmail.com)

### ABSTRACT

**Introduction:** Preeclampsia is one of the causes of the high maternal mortality rate seen on a national and regional scale and is usually characterized by symptoms such as hypertension, edema, proteinuria. In 2021 there were 23 cases of preeclampsia and 11 cases with preeclampsia referred. The results of interviews with 2 mothers who experienced preeclampsia were a history of hypertension before pregnancy and the mother's age was > 35 years with multipara. DASH (Dietary Approaches to Stop Hypertension) is an effective diet method for people with hypertension. The DASH diet approach can be used as a non-pharmacological therapy and as a substitute for drugs.

**Case Presentation:** the results of the study showed that participants had complaints of dizziness and heartburn for 3 days, pregnant women at the age of > 35 years and had a history of high blood pressure before pregnancy. Objective data showed high blood pressure 150/100, edema in the right and left legs, positive urine protein. The diagnosis in this case was preeclampsia with initial treatment in collaboration with doctors for further examination and treatment and recommending participants to adopt a DASH diet every day at home. Notes of development in the first week, participants were no longer dizzy and heartburn and edema were reduced. The results of the second week, participants did not experience edema and decreased blood pressure. **Conclusion:** Treatment of preeclampsia with the DASH diet method can lower blood pressure when properly applied every day.

**Keywords:** Preeclampsia, DASH, Pregnancy

## 1. Introduction

Preeclampsia is a contributor to the high maternal mortality rate is preeclampsia. Preeclampsia or eclampsia is a complication that is often experienced by pregnant women. Viewed on a national and regional scale, preeclampsia or eclampsia is a major cause of maternal death, namely bleeding, preeclampsia, and infection [Elsanti, 2020].

Preeclampsia is a collection of symptoms that occur in pregnant women, childbirth, and the puerperium consisting of hypertension, edema, and proteinuria that appear at 20 weeks gestation until the end of the first week after delivery. Preeclampsia is a serious problem and has a fairly high level of complexity. The magnitude of this problem is not only because preeclampsia affects the mother during pregnancy and childbirth, but also causes postpartum problems, such as the risk of cardiometabolic disease and other complications [Muzalfah et al., 2018].

According to the World Health Organization (2019), the maternal mortality rate in the world is 303,000 people. The Maternal Mortality Rate in Association of Southeast Asian Nations is 235 per 100,000 live births (ASEAN Secretariat, 2020). According to the Indonesian Demographic and Health Survey Data, the Maternal Mortality Rate in Indonesia increased from 228 per 100,000 live births in 2002-2007 to 359 per 100,000 live births in 2007-2012. MMR decreased in 2012-2015 to 305 per 100,000 live births and the number of

maternal deaths in Indonesia in 2019 was 4,221 cases [Kemenkes RI, 2021].

According to the Ministry of Health in Indonesia, the maternal mortality rate is 1712 per 100,000 live births. Meanwhile, the IMR in 2017 was 10294 per 1,000 live births [Kemenkes RI, 2021].

MMR achievement in 2020 is 98.6/100,000 life birth; IMR: 7.79/1000 life birth and infant mortality rate 8.99/1000 life birth) although this figure is much better than the national target (MMR: 226/100,000 life birth; IMR: 24/1,000 life birth) however, the MMR achievement is lower than the MMR achievement in 2019 (AKI 76.93/100,000 KH; IMR: 8.24/1000 KH and AKABA 9.65/1000 KH [Dinas Kesehatan Provinsi Jawa Tengah, 2021].

The MMR in 2020 is 111.45 per 100,000 live births, this has increased again compared to 2019 which was 73.87 per 100,000 births. The most common causes of maternal death in Wonogiri Regency in 2020 were bleeding 5 people, severe preeclampsia (PEB) 3, broncho pneumonia 1 person, hyperthyroidism 1 person, heart disease 1 person, amniotic fluid embolism 1 person [Dinas Kesehatan Wonogiri, 2021].

The results of the study [Wijaya, 2021] with the title Management of Preeclampsia in Pregnant Women at PMB Siti Azizah Wijaya, S.ST., Bd showed that participants 1 and 2 had complaints of swollen feet accompanied by headaches. On examination of vital signs, participants 1 and 2 showed high blood pressure results. After being

given implementation in the form of observing blood pressure, recommending bed rest, implementing a diet with the DASH diet and soaking the feet using warm water mixed with salt, it was found that participant 1 was resolved at week 3 and participant 2 was resolved at week 4. This occurs because participant 2 is more cooperative than participant 1.

Based on the results of a preliminary study at the Kismantoro Wonogiri Health Center, the number of cases of preeclampsia has increased from 2020 to 2021. In 2020 the number of pregnant women with preeclampsia was 11 cases and preeclampsia were referred as many as 7 cases. Meanwhile, in 2021 there were 23 cases of preeclampsia and 11 cases with preeclampsia who were referred. The results of interviews with 2 mothers who experienced preeclampsia were a history of hypertension before pregnancy and the mother's age was over thirty five years with multipara.

Based on (Hapsari, 2016) One way to control hypertension, especially in pregnancy, is to adjust the diet using the DASH method. Since the publication of the results of the DASH diet research in 1997, the DASH diet has been widely recommended to prevent and treat hypertension. The Dietary Approaches to Stop Hypertension (DASH) diet study states that a diet high in vegetables, fruit, and low-fat dairy products that are low in saturated fat and total fat and high in potassium, calcium, and magnesium can lower systolic blood pressure by 6-11 mmHg, and diastolic blood pressure of 3-6 mmHg [Rahadiyanti et al., 2015] .

The DASH diet approach is not only used as a non-pharmacological therapy but can be used as a substitute for drugs [Fitriyana &

Wirawati, 2022]. To anticipate the occurrence of unwanted things in pregnant women with symptoms that lead to preeclampsia, it is recommended to carry out routine and regular antenatal checks. This is useful for early detection if the mother's body finds signs and symptoms of preeclampsia, so that the mother can immediately get treatment. In addition, patients can also do bed rest early and apply a good diet so that their condition does not worsen and is expected to be more stable than the previous condition.

Therefore, in helping efforts to accelerate the decline in MMR and IMR, one of the authors is implementing continuous care or Continuity of Care. Continuity of Care in midwifery is a series of continuous and comprehensive service activities starting from pregnancy, childbirth, postpartum, newborn care and family planning services that link women's particular health needs and individual circumstances (Homer et al., 2014).

## 2. PATIENT IDENTITY

The patient named Mrs. S, 36 years old. His last education was junior high school, he worked to take care of the household, address Gesing Kismantoro.

The main complaints are dizziness and heartburn for 3 days. Pregnant with 2nd child and never miscarried. History of disease, namely having had hypertension in the first pregnancy. HPHT: 09-11-2021

## 3. CLINICAL FINDINGS

Good general condition, Compos mentis conscious, Blood pressure: 150/100mmHg, Respiration: 20 x/min, Pulse: 80 x/min, Temperature: 36.5oC, TFU: 28 cm, head presentation, convergent, contractions: none, FHR : 142 x/minute, right and left lower extremity edema.

## 4. TIMELINE

Tanggal dan jam	Subjektif	Objektif	Assessment	Planning
07.07.22 10.00	Dizziness and heartburn for 3 days	Blood pressure: 150/100 mmHg, Respiration: 20 x/minute, Pulse: 80 x/minute, Temperature: 36.5oC, TFU: 28 cm, head presentation, convergent,	Mrs. S. G2P1A0 36 years old 34 weeks pregnant single fetus live intra uteri, longitudinal position,	9. Explain to the mother about the current status of pregnancy 10. Explain to the mother about preeclampsia 11. Observe the mother's vital signs, especially blood pressure, and ask the mother to lie on her left side while sleeping 12. Advise the mother to soak the feet in warm water 13. Give moral support to the mother
				contraction: none, FHR: 142 x/minute, lower extremity edema. cephalic presentation, puca, convergent with preeclampsia Mrs. S G2P1A0 36 weeks of age pregnant, single live intrauterine fetus, longitudinal 14. Give KIE to the mother regarding the method 15. Collaborate with doctors for further examination 16. Advise mother to visit again in 1 week 6. Explaining the results of the examination to the mother that the mother's condition has improved such as the mother is no longer dizzy and has heartburn, reduced blood pressure, reduced edema 7. Encourage mothers to continue implementing the DASH diet at home, namely a diet rich in fruits, vegetables, whole grains, nuts, fish, and low-fat milk. These foods are high in important

Commented [R2]: Tidy up writing method

		convergent, contraction: none, FHR: 142 x/minute, lower extremities slightly edematous	position, cephalic presentation, puca, convergent with preeclampsia	nutrients, such as potassium, magnesium, calcium, fiber, and protein. 8. Advise the mother to rest enough during the day 1-2 hours and 7-8 hours at night 9. Advise the mother to continue to take vitamins from the doctor 10. Advise control mother for 1 more week to monitor general condition and vital sign
<b>21.07.22 08.00</b>	Mom has no complaints, just wants to check her pregnancy	Blood pressure: 146/90 mmHg, Respiration: 20 x/minute, Pulse: 80 x/minute, Temperature: 36.5oC, TFU: 29 cm, head presentation, convergent, contraction: none, FHR: 150x/minute, extremities no edema	Mrs. S G2P1A0 36 years of age 36 weeks pregnant, single live intrauterine fetus, longitudinal position, cephalic presentation, puca, convergent with preeclampsia	9. Explain the results of the examination to the mother that her condition has improved, blood pressure has decreased, there is no edema, positive urine protein 10. Encourage mothers to continue implementing the DASH diet at home, namely a diet rich in fruits, vegetables, whole grains, nuts, fish, and low-fat milk. 11. Advise the mother to rest enough during the day 1-2 hours and 7-8 hours at night 12. Giving counseling to mothers with danger signs in pregnancy such as severe dizziness, blurred vision, heartburn, bleeding, seizures 13. Giving counseling to the mother for signs of labor such as urinating in the lower abdomen, removing blood mucus, removing fluid/amniotic fluid from the birth canal 14. Provide counseling to mothers regarding preparation for childbirth 15. Advise the mother to continue to take vitamins from the doctor 16. Instruct the mother for control in 1 week to monitor general condition, vital sign and signs of labor.

## 5. DIAGNOSTIC CHECK

Investigations carried out were urine protein examination with positive results. The diagnosis of this case was preeclampsia. This diagnosis is supported by research results which state that (Peres et al., 2018) Preeclampsia is a hypertensive condition found at gestational age over twenty weeks with or without urine protein and accompanied by other organ disorders.

The prognosis in this case is preterm delivery. This is in line with the theory Norwitz ER, Repke JT in Hidayati et al., 2018 One of the prognosis for preeclampsia in pregnancy is the risk of complications to the mother and fetus, such as impaired fetal growth, preterm birth, placental abruption, and Intra Uterin Fetal Death.

## 6. INTERVENTION THERAPY

The intervention carried out during the visit was to encourage mothers to apply the DASH method, namely a diet high in important nutrients, such as potassium, magnesium, calcium, fiber, and protein. (Nurhayati, 2021). Based on research (Porouw & Yulianingsih, 2019) The content of sodium and potassium in Ambon bananas is proven to be able to reduce high blood pressure in pregnant women when consumed regularly and with the right dose.

The next implementation is to encourage mothers to do bed rest (Novitasari et al., 2018), In addition, it is recommended that mothers soak their feet in warm water (Sabattani et al., 2016).

Management in cases of preeclampsia also provides referrals or recommendations to the hospital for further examination and treatment with the patient's consent.

## 7. FOLLOW UP AND RESULT

Monitoring is carried out every 1 week, by conducting home visits, at the first visit there is a development of the mother's condition, namely complaints of dizziness and heartburn have disappeared, reduced leg edema, there is a decrease in diastolic blood pressure of 5 mm Hg. The second visit showed progress with a decrease in systolic blood pressure of 4 mm Hg and diastolic 5 mm Hg, no edema, urine protein was still positive.

## 8. KESIMPULAN

Treatment of preeclampsia with the DASH method can lower blood pressure if applied every day properly. Collaboration with doctors is also very necessary for further examination and therapy.

## 9. INFORMED CONSENT

Informed consent has been done on July 7, 2022 and the patient is willing to be a respondent

## 10. REFERENCE

- Astuti. (2012). *Buku Ajaran Asuhan Kebidanan Ibu 1 (Kehamilan)*. Yogyakarta: Rahima Press
- Dea Prastika Hapsari, A. R. (2016). *Hubungan Pengetahuan dengan Perilaku Manajemen Hipertensi: Aktivitas Fisik dan Diet DASH Penderita Hipertensi di Desa Salamrejo*. III(2), 2016.
- Elsanti, B. D. (2020). *By Devita Elsanti*.
- Fitriyana, M., & Wirawati, M. K. (2022). Penerapan Pola Diet Dash Terhadap Tekanan Darah Pada Penderita Hipertensi Di Desa Kalikangkung Semarang. *Jurnal Manajemen Asuhan Keperawatan*, 6(1), 17–24.  
<https://doi.org/10.33655/mak.v6i1.126>
- Homer, C. S. E., Friberg, I. K., Dias, M. A. B., ten Hoope-Bender, P., Sandall, J., Speciale, A. M., & Bartlett, L. A. (2014). The projected effect of scaling up midwifery. *Lancet (London, England)*, 384(9948), 1146–1157.  
[https://doi.org/10.1016/S0140-6736\(14\)60790-X](https://doi.org/10.1016/S0140-6736(14)60790-X)
- Muzalfah, R., Santik, Y. D. P., & Wahyuningsih, A. S. (2018). Kejadian Preeklampsia pada Ibu Bersalin. *Higeia Journal Of Public Health Research Development*, 2(3), 1–12.  
<https://journal.unnes.ac.id/sju/index.php/higeia/article/view/21390/11738>
- Novitasari, P., Prasetyorini, H., & Prihati, D. R. (2018). Upaya Pemenuhan Kebutuhan Istirahat Tidur Pada Ibu Postpartum Dengan Preeklampsia. *Jurnal Manajemen Asuhan Keperawatan*, 2(1), 22–33.  
<https://doi.org/10.33655/mak.v2i1.31>
- Nurhayati, S. (2021). Penatalaksanaan Preeklampsia Pada Ibu Hamil. *STIKes Ngudia Husada Madura Oleh*.
- Nurianti, I., Saputri, I. N., & Crisdayanti Sitorus, B. (2021). Hubungan Dukungan Suami Dengan Kecemasan Ibu Hamil Dalam Menghadapi Proses Persalinan. *Jurnal Kebidanan Kestra (Jkk)*, 3(2),

163–169.

<https://doi.org/10.35451/jkk.v3i2.493>

- Peres, G. M., Mariana, M., & Cairrão, E. (2018). Pre-eclampsia and eclampsia: An update on the pharmacological treatment applied in Portugal. *Journal of Cardiovascular Development and Disease*, 5(1).  
<https://doi.org/10.3390/jcdd5010003>
- Porouw, H. S., & Yulianingsih, E. (2019). Pisang Ambon Dan Hipertensi Ibu Hamil. *Jambura Health and Sport Journal*, 1(2), 61–70.  
<https://doi.org/10.37311/jhsj.v1i2.2597>
- Putri, H. (2018). Determinan Kejadian Pre Eklamsi di RSD Kalisat Jember. *Jurnal Kesehatan Dr Soebandi*, 6(2), 21–29.
- Rahadiyanti, A., Setianto, B. Y., & Purba, M. B. (2015). Asupan makan DASH-like diet untuk mencegah risiko hipertensi pada wanita prediabetes. *Jurnal Gizi Klinik Indonesia*, 11(3), 115.  
<https://doi.org/10.22146/ijcn.19290>
- Sabattani et. al. (2016). Efektivitas rendam kaki dengan air hangat terhadap penurunan tekanan darah pada ibu hamil penderita preeklamsi di Puskesmas Ngaliyan Semarang. *Ilmu Keperawatan Dan Keidanan*, 5(1), 1–10.
- Yeyeh, A. R., Yolanda Sari, D., & Humaeroh, D. (2021). Hubungan Karakteristik Ibu Bersalin dengan Preeklamsia Berat di RSUD A Purwakarta Tahun 2020. *Jik*, 16–26.

#### 4. MANUSKRIP SETELAH REVISI

## PREECLAMPSIA CASE REPORT IN PREGNANT WOMEN IN WORK AREA KISMANTORO HEALTH CENTER WONOGIRI WITH DASH METHOD

Izzah Al Fawaidah<sup>1</sup>, Indri Astuti Purwanti<sup>2</sup>, Novita Nining Anggraini<sup>3</sup>, Fitriani Nur Damayanti<sup>4</sup>  
<sup>1,2,3,4</sup> Midwifery Professional Education Study Program, University of Muhammadiyah Semarang

\*Corresponding author: [izzahalfawaidah@gmail.com](mailto:izzahalfawaidah@gmail.com)

### ABSTRACT

**Introduction:** Preeclampsia is one of the causes of the high maternal mortality rate seen on a national and regional scale and is usually characterized by symptoms such as hypertension, edema, proteinuria. In 2021 there were 23 cases of preeclampsia and 11 cases with preeclampsia referred. The results of interviews with 2 mothers who experienced preeclampsia were a history of hypertension before pregnancy and the mother's age was > 35 years with multipara. DASH (Dietary Approaches to Stop Hypertension) is an effective diet method for people with hypertension. The DASH diet approach can be used as a non-pharmacological therapy and as a substitute for drugs. **Case Presentation:** the results of the study showed that participants had complaints of dizziness and heartburn for 3 days, pregnant women at the age of > 35 years and had a history of high blood pressure before pregnancy. Objective data showed high blood pressure 150/100, edema in the right and left legs, positive urine protein. The diagnosis in this case was preeclampsia with initial treatment in collaboration with doctors for further examination and treatment and recommending participants to adopt a DASH diet every day at home. Notes of development in the first week, participants were no longer dizzy and heartburn and edema were reduced. The results of the second week, participants did not experience edema and decreased blood pressure. **Conclusion:** Treatment of preeclampsia with the DASH diet method can lower blood pressure when properly applied every day. **Keywords:** Preeclampsia, DASH, Pregnancy

### 1. Introduction

Preeclampsia is a contributor to the high maternal mortality rate is preeclampsia. Preeclampsia or eclampsia is a complication that is often experienced by pregnant women. Viewed on a national and regional scale, preeclampsia or eclampsia is a major cause of maternal death, namely bleeding, preeclampsia, and infection [Elsanti, 2020].

Preeclampsia is a collection of symptoms that occur in pregnant women, childbirth, and the puerperium consisting of hypertension, edema, and proteinuria that appear at 20 weeks gestation until the end of the first week after delivery. Preeclampsia is a serious problem and has a fairly high level of complexity. The magnitude of this problem is not only because preeclampsia affects the mother during pregnancy and childbirth, but also causes postpartum problems, such as the risk of

cardiometabolic disease and other complications [Muzalfah et al., 2018].

According to the World Health Organization (2019), the maternal mortality rate in the world is 303,000 people. The Maternal Mortality Rate in Association of Southeast Asian Nations is 235 per 100,000 live births (ASEAN Secretariat, 2020). According to the Indonesian Demographic and Health Survey Data, the Maternal Mortality Rate in Indonesia increased from 228 per 100,000 live births in 2002-2007 to 359 per 100,000 live births in 2007-2012. MMR decreased in 2012-2015 to 305 per 100,000 live births and the number of maternal deaths in Indonesia in 2019 was 4,221 cases [Kemenkes RI, 2021].

According to the Ministry of Health in Indonesia, the maternal mortality rate is 1712 per 100,000 live births. Meanwhile, the IMR in 2017 was 10294 per 1,000 live births [Kemenkes RI, 2021].

MMR achievement in 2020 is 98.6/100,000 life birth; IMR: 7.79/1000 life birth and infant mortality rate 8.99/1000 life birth) although this figure is much better than the national target (MMR: 226/100,000 life birth; IMR: 24/1,000 life birth) however, the MMR achievement is lower than the MMR achievement in 2019 (AKI 76.93/100,000 KH; IMR: 8.24/1000 KH and AKABA 9.65/1000 KH [Dinas Kesehatan Provinsi Jawa Tengah, 2021].

The MMR in 2020 is 111.45 per 100,000 live births, this has increased again compared to 2019 which was 73.87 per 100,000 births. The most common causes of maternal death in Wonogiri Regency in 2020 were bleeding 5 people, severe preeclampsia (PEB) 3, broncho pneumonia 1 person, hyperthyroidism 1 person, heart disease 1 person, amniotic fluid embolism 1 person [Dinas Kesehatan Wonogiri, 2021].

The results of the study [Wijaya, 2021] with the title Management of Preeclampsia in Pregnant Women at PMB Siti Azizah Wijaya, S.ST., Bd showed that participants 1 and 2 had complaints of swollen feet accompanied by headaches. On examination of vital signs, participants 1 and 2 showed high blood pressure results. After being given implementation in the form of observing blood pressure, recommending bed rest, implementing a diet with the DASH diet and soaking the feet using warm water mixed with salt, it was found that participant 1 was resolved at week 3 and participant 2 was resolved at week 4. This occurs because participant 2 is more cooperative than participant 1.

Based on the results of a preliminary study at the Kismantoro Wonogiri Health Center, the number of cases of preeclampsia has increased from 2020 to 2021. In 2020 the number of pregnant women with preeclampsia was 11 cases and preeclampsia were referred as many as 7 cases. Meanwhile, in 2021 there were 23 cases of preeclampsia and 11 cases with preeclampsia who were referred. The results of interviews with 2 mothers who experienced preeclampsia were a history of hypertension before pregnancy and the mother's age was over thirty five years with multipara.

Based on (Hapsari, 2016) One way to control hypertension, especially in pregnancy, is to adjust the diet using the DASH method. Since the publication of the

results of the DASH diet research in 1997, the DASH diet has been widely recommended to prevent and treat hypertension. The Dietary Approaches to Stop Hypertension (DASH) diet study states that a diet high in vegetables, fruit, and low-fat dairy products that are low in saturated fat and total fat and high in potassium, calcium, and magnesium can lower systolic blood pressure by 6-11 mmHg. and diastolic blood pressure of 3-6 mmHg [Rahadiyanti et al., 2015] .

The DASH diet approach is not only used as a non-pharmacological therapy but can be used as a substitute for drugs [Fitriyana & Wirawati, 2022]. To anticipate the occurrence of unwanted things in pregnant women with symptoms that lead to preeclampsia, it is recommended to carry out routine and regular antenatal checks. This is useful for early detection if the mother's body finds signs and symptoms of preeclampsia, so that the mother can immediately get treatment. In addition, patients can also do bed rest early and apply a good diet so that their condition does not worsen and is expected to be more stable than the previous condition.

Therefore, in helping efforts to accelerate the decline in MMR and IMR, one of the authors is implementing continuous care or Continuity of Care. Continuity of Care in midwifery is a series of continuous and comprehensive service activities starting from pregnancy, childbirth, postpartum, newborn care and family planning services that link women's particular health needs and individual circumstances (Homer et al., 2014).

## **2. PATIENT IDENTITY**

The patient named Mrs. S, 36 years old. His last education was junior high school, he worked to take care of the household, address Gesing Kismantoro.

The main complaints are dizziness and heartburn for 3 days. Pregnant with 2nd child and never miscarried. History of disease, namely having had hypertension in the first pregnancy. HPHT: 09-11-2021

## **3. CLINICAL FINDINGS**

Good general condition, Compos mentis consciousness, Blood pressure: 150/100mmHg, Respiration: 20 x/min, Pulse: 80 x/min, Temperature: 36.5oC, TFU: 28 cm, head presentation, convergent, contractions: none, FHR : 142 x/minute, right and left lower extremity edema.

#### 4. TIMELINE

Tanggal dan jam	Subjektif	Objektif	Assessment	Planning
07.07.22 10.00	Dizziness and heartburn for 3 days	Blood pressure: 150/100 mmHg, Respiration: 20 x/minute, Pulse: 80 x/minute, Temperature: 36.5oC, TFU: 28 cm, head presentation, convergent, contraction: none, FHR: 142 x/minute, lower extremity edema.	Mrs. S G2P1A0 36 years old 34 weeks pregnant single fetus live intra uteri, longitudinal position, cephalic presentation, puca, convergent with preeclampsia	17. Explain to the mother about the current state of her pregnancy 18. Explain to the mother about preeclampsia 19. Observation of KU, vital signs, especially blood pressure, and ask the mother to tilt to the left while sleeping. 20. Advise the mother to bed rest and soak the feet in warm water 21. Give moral support to mother 22. Give KIE to the mother regarding the management of preeclampsia with the DASH method 23. Collaborate with doctors for further examination 24. Advise mother to visit again in 1 week
14.07.22 09.00	Mom is no longer dizzy and no pain in the pit of the heart	Blood pressure: 150/95 mmHg, Respiration: 20 x/minute, Pulse: 80 x/minute, Temperature: 36.5oC, TFU: 28 cm, head presentation, convergent, contraction: none, FHR: 142 x/minute, lower extremities slightly edematous	Mrs. S G2P1A0 36 years of age 35 weeks pregnant, single live intrauterine fetus, longitudinal position, cephalic presentation, puca, convergent with preeclampsia	11. Explaining the results of the examination to the mother that the mother's condition has improved such as the mother is no longer dizzy and has heartburn, reduced blood pressure, reduced edema 12. Encourage mothers to continue implementing the DASH diet at home, namely a diet rich in fruits, vegetables, whole grains, nuts, fish, and low-fat milk. These foods are high in important nutrients, such as potassium, magnesium, calcium, fiber, and protein. 13. Advise the mother to rest enough during the day 1-2 hours and 7-8 hours at night 14. Advise the mother to continue to take vitamins from the doctor 15. Advise control mother for 1 more week to monitor general condition and vital sign
21.07.22 08.00	Mom has no complaints, just wants to check her pregnancy	Blood pressure: 146/90 mmHg, Respiration: 20 x/minute, Pulse: 80 x/minute, Temperature: 36.5oC, TFU: 29 cm, head presentation, convergent, contraction: none, FHR: 150x/minute, extremities no edema	Mrs. S G2P1A0 36 years of age 36 weeks pregnant, single live intrauterine fetus, longitudinal position, cephalic presentation, puca, convergent with preeclampsia	17. Explain the results of the examination to the mother that her condition has improved, blood pressure has decreased, there is no edema, positive urine protein 18. Encourage mothers to continue implementing the DASH diet at home, namely a diet rich in fruits, vegetables, whole grains, nuts, fish, and low-fat milk. 19. Advise the mother to rest enough during the day 1-2 hours and 7-8 hours at night 20. Giving counseling to mothers with danger signs in pregnancy such as severe dizziness, blurred vision, heartburn, bleeding, seizures 21. Giving counseling to the mother for signs of labor such as urinating in the lower abdomen, removing blood mucus, removing fluid/amniotic fluid from the birth canal 22. Provide counseling to mothers regarding preparation for childbirth 23. Advise the mother to continue to take vitamins from the doctor



---

24. Instruct the mother for control in 1 week to monitor general condition, vital sign and signs of labor.

---

## 5. DIAGNOSTIC CHECK

Investigations carried out were urine protein examination with positive results. The diagnosis of this case was preeclampsia. This diagnosis is supported by research results which state that (Peres et al., 2018) Preeclampsia is a hypertensive condition found at gestational age over twenty weeks with or without urine protein and accompanied by other organ disorders.

The prognosis in this case is preterm delivery. This is in line with the theory Norwitz ER, Repke JT in Hidayati et al., 2018 One of the prognosis for preeclampsia in pregnancy is the risk of complications to the mother and fetus, such as impaired fetal growth, preterm birth, placental abruption, and Intra Uterin Fetal Death.

## 6. INTERVENTION THERAPY

The intervention carried out during the visit was to encourage mothers to apply the DASH method, namely a diet high in important nutrients, such as potassium, magnesium, calcium, fiber, and protein. (Nurhayati, 2021). Based on research (Porouw & Yulianingsih, 2019) The content of sodium and potassium in Ambon bananas is proven to be able to reduce high blood pressure in pregnant women when consumed regularly and with the right dose.

The next implementation is to encourage mothers to do bed rest (Novitasari et al., 2018), In addition, it is recommended that mothers soak their feet in warm water (Sabattani et al., 2016).

Management in cases of preeclampsia also provides referrals or recommendations to the hospital for further examination and treatment with the patient's consent.

## FOLLOW UP AND RESULT

Monitoring is carried out every 1 week, by conducting home visits, at the first visit there is a development of the mother's condition, namely complaints of dizziness and heartburn have disappeared, reduced leg edema, there is a decrease in diastolic blood pressure of 5 mm Hg. The second visit showed progress with a decrease in systolic

blood pressure of 4 mm Hg and diastolic 5 mm Hg, no edema, urine protein was still positive.

## 8. KESIMPULAN

Treatment of preeclampsia with the DASH method can lower blood pressure if applied every day properly. Collaboration with doctors is also very necessary for further examination and therapy.

## 9. INFORMED CONSENT

Informed consent has been done on July 7, 2022 and the patient is willing to be a respondent

## 10. REFERENCE

- Astuti. (2012). *Buku Ajaran Asuhan Kebidanan Ibu 1 (Kehamilan)*. Yogyakarta: Rahima Press
- Dea Prastika Hapsari, A. R. (2016). *Hubungan Pengetahuan dengan Perilaku Manajemen Hipertensi: Aktivitas Fisik dan Diet DASH Penderita Hipertensi di Desa Salamrejo. III(2)*, 2016.
- Elsanti, B. D. (2020). *By Devita Elsanti*.
- Fitriyana, M., & Wirawati, M. K. (2022). Penerapan Pola Diet Dash Terhadap Tekanan Darah Pada Penderita Hipertensi Di Desa Kalikangkung Semarang. *Jurnal Manajemen Asuhan Keperawatan*, 6(1), 17–24.  
<https://doi.org/10.33655/mak.v6i1.126>
- Homer, C. S. E., Friberg, I. K., Dias, M. A. B., ten Hoope-Bender, P., Sandall, J., Speciale, A. M., & Bartlett, L. A. (2014). The projected effect of scaling up midwifery. *Lancet (London, England)*, 384(9948), 1146–1157.  
[https://doi.org/10.1016/S0140-6736\(14\)60790-X](https://doi.org/10.1016/S0140-6736(14)60790-X)
- Muzalfah, R., Santik, Y. D. P., & Wahyuningsih, A. S. (2018). Kejadian Preeklampsia pada Ibu Bersalin. *Higeia Journal Of Public Health Research Development*, 2(3), 1–12.  
<https://journal.unnes.ac.id/sju/index.php/higeia/article/view/21390/11738>
- Novitasari, P., Prasetyorini, H., & Prihati, D. R. (2018). Upaya Pemenuhan Kebutuhan Istirahat Tidur Pada Ibu Postpartum Dengan Preeklampsia. *Jurnal Manajemen Asuhan Keperawatan*, 2(1), 22–33.  
<https://doi.org/10.33655/mak.v2i1.31>

- Nurhayati, S. (2021). Penatalaksanaan Preeklamsia Pada Ibu Hamil. *STIKes Ngudia Husada Madura Oleh*.
- Nurianti, I., Saputri, I. N., & Crisdayanti Sitorus, B. (2021). Hubungan Dukungan Suami Dengan Kecemasan Ibu Hamil Dalam Menghadapi Proses Persalinan. *Jurnal Kebidanan Kestra (Jkk)*, 3(2), 163–169.  
<https://doi.org/10.35451/jkk.v3i2.493>
- Peres, G. M., Mariana, M., & Cairrão, E. (2018). Pre-eclampsia and eclampsia: An update on the pharmacological treatment applied in Portugal. *Journal of Cardiovascular Development and Disease*, 5(1).  
<https://doi.org/10.3390/jcdd5010003>
- Porouw, H. S., & Yulianingsih, E. (2019). Pisang Ambon Dan Hipertensi Ibu Hamil. *Jambura Health and Sport Journal*, 1(2), 61–70.
- <https://doi.org/10.37311/jhsj.v1i2.2597>
- Putri, H. (2018). Determinan Kejadian Pre Eklamsi di RSD Kalisat Jember. *Jurnal Kesehatan Dr Soebandi*, 6(2), 21–29.
- Rahadiyanti, A., Setianto, B. Y., & Purba, M. B. (2015). Asupan makan DASH-like diet untuk mencegah risiko hipertensi pada wanita prediabetes. *Jurnal Gizi Klinik Indonesia*, 11(3), 115.  
<https://doi.org/10.22146/ijcn.19290>
- Sabattani et. al. (2016). Efektivitas rendam kaki dengan air hangat terhadap penurunan tekanan darah pada ibu hamil penderita preeklamsi di Puskesmas Ngaliyan Semarang. *Ilmu Keperawatan Dan Keidanan*, 5(1), 1–10.
- Yeyeh, A. R., Yolanda Sari, D., & Humaeroh, D. (2021). Hubungan Karakteristik Ibu Bersalin dengan Preeklampsia Berat di RSU A Purwakarta Tahun 2020. *Jik*, 16–26.

## 5. IN PRESS



**International Conference on Practice and Innovation  
Midwifery (ICPIM)**

**Universtas Muhammadiyah Semarang**  
Jl. Kedungmundu Raya No. 18 Semarang – ID 50273  
Phone: +62 24-76740288 Fax: + 62 24-76740287 E-mail: icpim@unimus.ac.id



**ORAL PRESENTATION ACCEPTANCE**

August, 12<sup>th</sup> 2022

Dear **Izzah Al Fawaidah**,

Thank you for submitting a paper to present at “**THE 2nd INTERNATIONAL CONFERENCE ON PRACTICE AND INNOVATION IN MIDWIFERY**”. On behalf of the Organizing Committee, I am delighted to inform you that your abstract has been accepted for an oral presentation at the conference.

Details of your abstract are as follows:

ID : **PR144**  
Title : **Case Report Preeclamsia In Pregnant Women In Work Area Kismantoro Health Center Wonogiri With Dash Method**  
Authors : Izzah Al Fawaidah  
Presenting Author : Izzah Al Fawaidah

Please check the above details of your presentation carefully as all conference material will be printed this information. If there are any corrections please inform the committee as soon as possible by email. You will be informed in a separate email about the scheduling of your presentation and you will be given a final programme at the conference.

Furthermore, you are requested to immediately complete the administrative requirements related to payment as confirmation of your participation as a presenter and payment receipt can be sent via WhatsApp to **Mrs. Siti Istiana: +62 856-4027-0631**. Please send the presentation file by August 15<sup>th</sup>, 2022 in ppt/pptx format regarding the articles to be presented. PPT files please sent to email with the subject : **File PPT PR144**.

Please be advised that your registration fee does not include travel insurance, please ensure that you have your own insurance policy before travelling.

Kind regards,

Indri Astuti Purwanti

Conference Content Executive.

**6. ARTIKEL SUDAH PUBLISH**



Articles

1 result (0.00 sec)

My profile My library

Any time

Since 2023

Since 2022

Since 2019

Custom range ...

Sort by relevance

Sort by date

Any type

Review articles

include patents

include citations

Create alert

(pdf) CASE REPORT PREECLAMISIA IN PREGNANT WOMEN IN WORK AREA KISMANTORO HEALTH CENTER WONOGIRI WITH DASH METHOD

[PDF] unimus.ac.id

I A) Fawaidah, (S. Darmas) on Practice and ... 2022 - prosiding.unimus.ac.id

The DASH diet approach can be used as a non-pharmacological therapy and as a substitute for drugs. The results of the study showed that participants had complaints of dizziness and heartburn for 3 days, pregnant women at the age of > 35 years and had a history of high blood pressure before pregnancy. Objective data showed high blood pressure 150/100, edema in the right and left legs, positive urine protein. The diagnosis in this ... Based on the results of a preliminary study at the Kismantoro Wonogiri Health Center, the ...

☆ Save ⓘ Cite Related articles 00



## CASE REPORT PREECLAMPSIA IN PREGNANT WOMEN IN WORK AREA KISMANTORO HEALTH CENTER WONOGIRI WITH DASH METHOD

Izzah Al Fawaidah<sup>1</sup>, Indri Astuti Purwanti<sup>2</sup>, Novita Nining Anggraini<sup>3</sup>, Fitriani Nur Damayanti<sup>4</sup>

<sup>1</sup> Student of the Midwife Profession Education Department, Universitas Muhammadiyah Semarang

<sup>2,3,4</sup> Lecturer of Midwifery Department, Universitas Muhammadiyah Semarang

Corresponding author : [izzahalfawaidah@gmail.com](mailto:izzahalfawaidah@gmail.com)

### ABSTRACT

Preeclampsia is one of the causes of the high maternal mortality rate seen on a national and regional scale and is usually characterized by symptoms such as hypertension, edema, proteinuria. In 2021 there were 23 cases of preeclampsia and 11 cases with preeclampsia referred. The results of interviews with 2 mothers who experienced preeclampsia were a history of hypertension before pregnancy and the mother's age was > 35 years with multipara. DASH (Dietary Approaches to Stop Hypertension) is an effective diet method for people with hypertension. The DASH diet approach can be used as a non-pharmacological therapy and as a substitute for drugs. The results of the study showed that participants had complaints of dizziness and heartburn for 3 days, pregnant women at the age of > 35 years and had a history of high blood pressure before pregnancy. Objective data showed high blood pressure 150/100, edema in the right and left legs, positive urine protein. The diagnosis in this case was preeclampsia with initial treatment in collaboration with doctors for further examination and treatment and recommending participants to adopt a DASH diet every day at home. Notes of development in the first week, participants were no longer dizzy and heartburn and edema were reduced. The results of the second week, participants did not experience edema and decreased blood pressure. Treatment of preeclampsia with the DASH diet method can lower blood pressure when properly applied every day.

**Keywords:** Preeclampsia, DASH, Pregnancy

### 1. INTRODUCTION

Preeclampsia is a contributor to the high maternal mortality rate is preeclampsia. Preeclampsia or eclampsia is a complication that is often experienced by pregnant women. Viewed on a national and regional scale, preeclampsia or eclampsia is a major cause of maternal death, namely bleeding, preeclampsia, and infection [Elsanti, 2020].

Preeclampsia is a collection of symptoms that occur in pregnant women, childbirth, and the puerperium consisting of hypertension, edema, and proteinuria that appear at 20 weeks gestation until the end of the first week after delivery. Preeclampsia is a serious problem and has a fairly high level of complexity. The magnitude of this problem is not only because preeclampsia affects the mother during pregnancy and childbirth, but also causes postpartum problems, such as the risk of cardiometabolic disease and other complications [Muzalfah et al., 2018].

According to the World Health Organization (2019), the maternal mortality rate in the world is 303,000 people. The Maternal Mortality Rate in Association of Southeast Asian Nations is 235 per 100,000 live births (ASEAN Secretariat, 2020). According to the Indonesian Demographic and Health Survey Data, the Maternal Mortality Rate in Indonesia increased from 228 per 100,000 live births in 2002-2007 to 359 per 100,000 live births in 2007-2012. MMR decreased in 2012-2015 to 305 per 100,000 live births and the number of maternal deaths in Indonesia in 2019 was 4,221 cases [Kemenkes RI, 2021].

According to the Ministry of Health in Indonesia, the maternal mortality rate is 1712 per 100,000 live births. Meanwhile, the IMR in 2017 was 10294 per 1,000 live births [Kemenkes RI, 2021].

MMR achievement in 2020 is 98.6/100,000 life birth; IMR: 7.79/1000 life birth and infant mortality rate 8.99/1000 life birth) although this figure is much better than the national target



(MMR: 226/100,000 life birth; IMR: 24/1,000 life birth) however, the MMR achievement is lower than the MMR achievement in 2019 (AKI 76.93/100,000 KH; IMR: 8.24/1000 KH and AKABA 9.65/1000 KH [Dinas Kesehatan Provinsi Jawa Tengah, 2021].

The MMR in 2020 is 111.45 per 100,000 live births, this has increased again compared to 2019 which was 73.87 per 100,000 births. The most common causes of maternal death in Wonogiri Regency in 2020 were bleeding 5 people, severe pre-eclampsia (PEB) 3, broncho pneumonia 1 person, hyperthyroidism 1 person, heart disease 1 person, amniotic fluid embolism 1 person [Dinas Kesehatan Wonogiri, 2021].

The results of the study [Wijaya, 2021] with the title Management of Preeclampsia in Pregnant Women at PMB Siti Azizah Wijaya, S.ST., Bd showed that participants 1 and 2 had complaints of swollen feet accompanied by headaches. On examination of vital signs, participants 1 and 2 showed high blood pressure results. After being given implementation in the form of observing blood pressure, recommending bed rest, implementing a diet with the DASH diet and soaking the feet using warm water mixed with salt, it was found that participant 1 was resolved at week 3 and participant 2 was resolved at week 4. This occurs because participant 2 is more cooperative than participant 1.

Based on the results of a preliminary study at the Kismantoro Wonogiri Health Center, the number of cases of preeclampsia has increased from 2020 to 2021. In 2020 the number of pregnant women with preeclampsia was 11 cases and preeclampsia were referred as many as 7 cases. Meanwhile, in 2021 there were 23 cases of preeclampsia and 11 cases with preeclampsia who were referred. The results of interviews with 2 mothers who experienced preeclampsia were a history of hypertension before pregnancy and the mother's age was over thirty five years with multipara.

Based on (Hapsari, 2016) One way to control hypertension, especially in pregnancy, is to adjust the diet using the DASH method. Since the publication of the results of the DASH diet research in 1997, the DASH diet has been widely recommended to prevent and treat hypertension. The Dietary Approaches to Stop Hypertension (DASH) diet study states that a diet high in vegetables, fruit, and low-fat dairy products that

are low in saturated fat and total fat and high in potassium, calcium, and magnesium can lower systolic blood pressure by 6-11 mmHg. and diastolic blood pressure of 3-6 mmHg [Rahadiyanti et al., 2015].

The DASH diet approach is not only used as a non-pharmacological therapy but can be used as a substitute for drugs [Fitriyana & Wirawati, 2022]. To anticipate the occurrence of unwanted things in pregnant women with symptoms that lead to preeclampsia, it is recommended to carry out routine and regular antenatal checks. This is useful for early detection if the mother's body finds signs and symptoms of preeclampsia, so that the mother can immediately get treatment. In addition, patients can also do bed rest early and apply a good diet so that their condition does not worsen and is expected to be more stable than the previous condition.

Therefore, in helping efforts to accelerate the decline in MMR and IMR, one of the authors is implementing continuous care or Continuity of Care. Continuity of Care in midwifery is a series of continuous and comprehensive service activities starting from pregnancy, childbirth, postpartum, newborn care and family planning services that link women's particular health needs and individual circumstances (Homer et al., 2014).

## 2. PATIENT IDENTITY

The patient named Mrs. S, 36 years old. His last education was junior high school, he worked to take care of the household, address Gesing Kismantoro.

The main complaints are dizziness and heartburn for 3 days. Pregnant with 2nd child and never miscarried. History of disease, namely having had hypertension in the first pregnancy.

## 3. CLINICAL FINDINGS

Good general condition, Compos mentis consciousness, Blood pressure: 150/100mmHg, Respiration: 20 x/min, Pulse: 80 x/min, Temperature: 36.5oC, TFU: 28 cm, head presentation, convergent, contractions: none, FHR : 142 x/minute, right and left lower extremely edema.



#### 4. TIMELINE

Time	Subjective	Objective	Assessment	Planning
07.07.22 10.00	Dizziness and heartburn for 3 days	Blood pressure: 150/100 mmHg. Respiration: 20 x/minute, Pulse: 80 x/minute, Temperature: 36.5oC, TFU: 28 cm, head presentation, convergent, contraction: none, FHR: 142 x/minute, lower extremity edema.	Mrs. S G2P1A0 36 years old 34 weeks pregnant, single fetus live intra uteri, longitudinal position, cephalic presentation, puca, convergent with preeclampsia	1. Explain to the mother about the current state of her pregnancy 2. Explain to the mother about preeclampsia 3. Observation of KU, vital signs, especially blood pressure, and ask the mother to tilt to the left while sleeping. 4. Advise the mother to bed rest and soak the feet in warm water 5. Give moral support to mother 6. Give KIE to the mother regarding the management of preeclampsia with the DASH method 7. Collaborate with doctors for further examination 8. Advise mother to visit again in 1 week
14.07.22 09.00	Mom is no longer dizzy and no pain in the pit of the heart	Blood pressure: 150/95 mmHg. Respiration: 20 x/minute, Pulse: 80 x/minute, Temperature: 36.5oC, TFU: 28 cm, head presentation, convergent, contraction: none, FHR: 142 x/minute, lower extremities slightly edematous	Mrs. S G2P1A0 36 years of age 35 weeks pregnant, single live intrauterine fetus, longitudinal position, cephalic presentation, puca, convergent with preeclampsia	1. Explaining the results of the examination to the mother that the mother's condition has improved such as the mother is no longer dizzy and has heartburn, reduced blood pressure, reduced edema 2. Encourage mothers to continue implementing the DASH diet at home, namely a diet rich in fruits, vegetables, whole grains, nuts, fish, and low-fat milk. These foods are high in important nutrients, such as potassium, magnesium, calcium, fiber, and protein. 3. Advise the mother to rest enough during the day 1-2 hours and 7-8 hours at night 4. Advise the mother to continue to take vitamins from the doctor 5. Advise control mother for 1 more week to monitor general condition and vital sign
21.07.22 08.00	Mom has no complaints, just wants to check her pregnancy	Blood pressure: 146/90 mmHg. Respiration: 20 x/minute, Pulse: 80 x/minute, Temperature: 36.5oC, TFU: 29 cm, head presentation, convergent, contraction: none, FHR: 150x/minute, extremities no edema	Mrs. S G2P1A0 36 years of age 36 weeks pregnant, single live intrauterine fetus, longitudinal position, cephalic presentation, puca, convergent with preeclampsia	1. Explain the results of the examination to the mother that her condition has improved, blood pressure has decreased, there is no edema, positive urine protein 2. Encourage mothers to continue implementing the DASH diet at home, namely a diet rich in fruits, vegetables, whole grains, nuts, fish, and low-fat milk. 3. Advise the mother to rest enough during the day 1-2 hours and 7-8 hours at night 4. Giving counseling to mothers with danger signs in pregnancy such as severe dizziness, blurred vision, heartburn, bleeding, seizures 5. Giving counseling to the mother for signs of labor such as urinating in the lower abdomen, removing blood mucus, removing fluid/amniotic fluid from the birth canal 6. Provide counseling to mothers regarding preparation for childbirth 7. Advise the mother to continue to take vitamins from the doctor 8. Instruct the mother for control in 1 week to monitor general condition, vital sign and signs of labor.

#### 5. DIAGNOSTIC CHECK

Investigations carried out were urine protein examination with positive results. The diagnosis of this case was preeclampsia. This diagnosis is supported by research results which state that (Peres et al., 2018) Preeclampsia is a hypertensive condition found at gestational age over twenty

weeks with or without urine protein and accompanied by other organ disorders.

The prognosis in this case is preterm delivery. This is in line with the theory Norwitz ER, Repke JT in Hidayati et al., 2018 One of the prognosis for preeclampsia in pregnancy is the risk of complications to the mother and fetus, such as



impaired fetal growth, preterm birth, placental abruption, and Intra Uterin Fetal Death.

#### 6. INTERVENTION THERAPY

The intervention carried out during the visit was to encourage mothers to apply the DASH method, namely a diet high in important nutrients, such as potassium, magnesium, calcium, fiber, and protein. (Nurhayati, 2021). Based on research (Porouw & Yulianingsih, 2019) The content of sodium and potassium in Ambon bananas is proven to be able to reduce high blood pressure in pregnant women when consumed regularly and with the right dose.

The next implementation is to encourage mothers to do bed rest (Novitasari et al., 2018). In addition, it is recommended that mothers soak their feet in warm water (Sabattani et al., 2016).

Management in cases of preeclampsia also provides referrals or recommendations to the hospital for further examination and treatment with the patient's consent.

#### 7. FOLLOW UP AND RESULT

Monitoring is carried out every 1 week, by conducting home visits, at the first visit there is a development of the mother's condition, namely complaints of dizziness and heartburn have disappeared, reduced leg edema, there is a decrease in diastolic blood pressure of 5 mm Hg. The second visit showed progress with a decrease in systolic blood pressure of 4 mm Hg and diastolic 5 mm Hg, no edema, urine protein was still positive.

#### 8. CONCLUSION

Treatment of preeclampsia with the DASH method can lower blood pressure if applied every day properly. Collaboration with doctors is also very necessary for further examination and therapy.

#### 9. INFORMED CONSENT

Informed consent has been done on July 7, 2022 and the patient is willing to be a respondent.

#### REFERENCE

- [1] Astuti. (2012). Buku Ajaran Asuhan Kebidanan Ibu 1 (Kehamilan). Yogyakarta: Rahima Press
- [2] Dea Prastika Hapsari, A. R. (2016). *Hubungan Pengetahuan dengan Perilaku Manajemen Hipertensi: Aktivitas Fisik dan Diet DASH Penderita Hipertensi di Desa Salamrejo*. III(2), 2016.
- [3] Elsanti, B. D. (2020). *By Devita Elsanti*.
- [4] Fitriyana, M., & Wirawati, M. K. (2022). Penerapan Pola Diet Dash Terhadap Tekanan Darah Pada Penderita Hipertensi Di Desa Kalikangkung Semarang. *Jurnal Manajemen Asuhan Keperawatan*, 6(1), 17-24. <https://doi.org/10.33655/mak.v6i1.126>
- [5] Homer, C. S. E., Friberg, I. K., Dias, M. A. B., ten Hoop-Bender, P., Sandall, J., Speciale, A. M., & Bartlett, L. A. (2014). The projected effect of scaling up midwifery. *Lancet (London, England)*, 384(9948), 1146-1157. [https://doi.org/10.1016/S0140-6736\(14\)60790-X](https://doi.org/10.1016/S0140-6736(14)60790-X)
- [6] Muzalfah, R., Santik, Y. D. P., & Wahyuningsih, A. S. (2018). Kejadian Preklampsia pada Ibu Bersalin. *Higeia Journal Of Public Health Research Development*, 2(3), 1-12. <https://journal.unnes.ac.id/sju/index.php/higci/a/article/view/21390/11738>
- [7] Novitasari, P., Prasetyorini, H., & Prihati, D. R. (2018). Upaya Pemenuhan Kebutuhan Istirahat Tidur Pada Ibu Postpartum Dengan Preklampsia. *Jurnal Manajemen Asuhan Keperawatan*, 2(1), 22-33. <https://doi.org/10.33655/mak.v2i1.31>
- [8] Nurhayati, S. (2021). Penatalaksanaan Preklamsia Pada Ibu Hamil. *STIKes Ngudia Husada Madura Oleh*.
- [9] Nurianti, I., Saputri, I. N., & Crisdayanti Sitorus, B. (2021). Hubungan Dukungan Suami Dengan Kecemasan Ibu Hamil Dalam Menghadapi Proses Persalinan. *Jurnal Kebidanan Kestra (Jkk)*, 3(2), 163-169. <https://doi.org/10.35451/jkk.v3i2.493>
- [10] Peres, G. M., Mariana, M., & Cairão, E. (2018). Pre-eclampsia and eclampsia: An update on the pharmacological treatment applied in Portugal. *Journal of Cardiovascular Development and Disease*, 5(1). <https://doi.org/10.3390/jcdd5010003>
- [11] Porouw, H. S., & Yulianingsih, E. (2019). Pisang Ambon Dan Hipertensi Ibu Hamil. *Jambura Health and Sport Journal*, 1(2), 61-70. <https://doi.org/10.37311/jhsj.v1i2.2597>
- [12] Putri, H. (2018). Determinan Kejadian Pre Eklamsi di RSD Kalisat Jember. *Jurnal Kesehatan Dr Soebandi*, 6(2), 21-29.
- [13] Rahadiyanti, A., Setianto, B. Y., & Purba, M. B. (2015). Asupan makan DASH-like diet untuk mencegah risiko hipertensi pada wanita prediabetes. *Jurnal Gizi Klinik Indonesia*, 11(3), 115. <https://doi.org/10.22146/jcn.19290>
- [14] Sabattani et al. (2016). Efektivitas rendam kaki dengan air hangat terhadap penurunan tekanan darah pada ibu hamil penderita preklamsi di Puskesmas Ngaliyan Semarang. *Ilmu Keperawatan Dan Keidanan*, 5(1), 1-10.
- [15] Yeyeh, A. R., Yolanda Sari, D., & Humaeroh, D. (2021). Hubungan Karakteristik Ibu Bersalin dengan Preklampsia Berat di RSU A Purwakarta Tahun 2020. *Jik*, 16-26.