

Description of Hemoglobin Levels for Pregnant Women Workers of The Djarum Cigarette Factory at The Mejobo Kudus Health Center

by Budi Santosa

Submission date: 22-May-2023 09:38AM (UTC+0700)

Submission ID: 2098782866

File name: 11723-36470-1-PB.pdf (236.23K)

Word count: 2246

Character count: 11294



Description of Hemoglobin Levels for Pregnant Women Workers of The Djarum Cigarette Factory at The Mejobo Kudus Health Center

Budi Santosa^{1*}, Hasna Widad Salsabila², Andri Sukeksi³

^{1,2,3}Department of Medical/Clinical Laboratory Science, Universitas Muhammadiyah Semarang, Central Java, Indonesia

Abstract

During pregnancy a woman is very susceptible to disorders of low levels of hemoglobin in the blood. Low hemoglobin values are associated with clinical problems such as anemia. Anemia can be more severe due to several factors, such as trimester of pregnancy, type of work, education, infectious diseases and so on. Most of the pregnant women in the Kudus Regency are pregnant women who work in cigarette factories. Pregnant women who work in cigarette factories will have an impact on their pregnancy, based on the theory that tobacco as a raw material for cigarettes contains addictive substances that are harmful to health. The purpose of this study was to find out the description of hemoglobin in pregnant women working in the Djarum cigarette factory at the Mejobo Kudus Health Center. This type of research is descriptive. The research was carried out in May 2022 and the examination was carried out at the Mejobo Health Center. Sampling with purposive sampling technique as many as 25 respondents. Analysis and data processing using tables. The results obtained from 25 respondents mostly had an average hemoglobin level of 12.3 g/dl, where the majority of respondents had normal hemoglobin of 88%. The majority of respondents do not experience anemia, this can occur because when working pregnant women use complete APD, from head coverings, masks, and aprons to protect themselves.

Keywords: hemoglobin; pregnant mother; cigarette factory workers

*Corresponding Author: Budi Santosa (email: budisantosa@unimus.ac.id), Kedungmundu Raya, Semarang.

Introduction

During pregnancy a woman is very susceptible to experiencing low levels of hemoglobin in the blood. Low hemoglobin values are associated with clinical problems such as anemia. The Centers for Disease Control and Prevention (CDC) defines anemia as a hemoglobin level lower than 11 gr/dl in the first and third trimesters, less than 10.5 gr/dl in the second trimester. Anemia can be more severe due to several factors, such as the third trimester of pregnancy, type of work, education, infection and so forth. The majority of pregnant women in the Kudus Regency area are pregnant women who work in cigarette factories. Pregnant women who work in cigarette factories will have an impact on their pregnancy, based on the theory that tobacco as a raw material for cigarettes contains addictive substances that are harmful to health. This can be one of the factors causing the mother to have symptoms of anemia or anemia. Based on the background of the problems above, it prompted researchers to conduct research by taking the title "Description of Hb Levels of Pregnant Women Djarum Cigarette Factory Workers at the Mejoko Kudus Community Health Center".

Method

Research Tools and Materials

The tools used in this study were a hematology analyzer and a tourniquet.

Materials used in the venous blood study, 70% alcohol, 3 cc syringe, alcohol swab, plaster, EDTA tube, and hematology analyzer tool.

Ways of working

How to Take Venous Blood

The area to be taken for blood (median cubital vein) is cleaned with 70% alcohol cotton, let it dry and a tourniquet is placed on the upper arm, approximately above the elbow. The patient is asked to clench his fist so that the veins are clearly visible. The syringe needle is inserted into the vein with the needle eye facing up. After it appears that blood is flowing in the syringe, the syringe suction is withdrawn slowly until as much as 3 ml of blood can be obtained. Remove the tourniquet and take the cotton and place it over the puncture site. The needle and syringe are removed slowly, the puncture marks are pressed using alcohol cotton. The needle is removed from the syringe, and blood is drawn into the EDTA tube through the tube wall.

How to Check Hb Levels using a Hematology Analyzer

Prepare an EDTA tube containing the patient's blood sample. Press the [MENU] key and select "count" then press enter. Press the [MODE] button then the examination method ("Whole Blood-ALL", "WB-WBC/HGH), or "WB-RBC/PLT" will appear on the top screen with a blue display. Then press the [F1] button to write down patient data, the blood sample in the EDTA tube is homogenized and inserted into the sample probe until it touches the bottom of the tube. Press the probe button for the calculation process and the examination results will appear on the screen, then the results are read.

Results and Discussion

Description of Hemoglobin Levels for Pregnant Women Workers of The Djarum Cigarette Factory ...

This research was carried out at the Mejobo Health Center Laboratory in Kudus Regency. The sample for the examination is the blood of pregnant women who carry out examinations at the Mejobo Health Center with EDTA anticoagulant. The data used in this research is primary data from the results of examination of hemoglobin levels of pregnant women who work in cigarette factories who carry out examinations at the Mejobo Kudus Health Center.

Table 1. Hemoglobin levels based on age in pregnant women who work at the Djarum cigarette factory at the Mejobo Kudus Health Center.

Age	Hemoglobin			Total
	Normal	Anemia Light	Anemia Heavy	
	Amount (%)	Amount (%)	Amount (%)	Amount (%)
20-29	12 (48%)	1 (4%)	1 (4%)	14 (56%)
30-39	8 (32%)	1 (4%)	0 (0)	9 (36%)
>40	2 (8%)	0 (0)	0 (0)	2 (8%)
Amount	14 (88%)	2 (8%)	1 (4%)	25 (100%)

Based on table 6, it is known that there are hemoglobin levels with an average 12 pregnant women working in cigarette factories aged 20-29 who have normal hemoglobin level of 13.4 g/dl.

Table 2. Hemoglobi levels based on the trimester of pregnancy in pregnant women who work at the Djarum cigarette factory at the Mejobo Kudus Health Center.

Pregnancy Trimesters	Hemoglobin			Total
	Normal	Anemia Light	Anemia Heavy	
	Amount (%)	Amount (%)	Amount (%)	Amount (%)
Trimester I	15 (60%)	1 (4%)	1 (4%)	17 (68%)
Trimesters II	4 (16%)	1 (4%)	0 (0)	5 (20%)
Trimesters III	3 (12%)	0 (0)	0 (0)	3 (12%)
Amount	22 (88%)	2 (8%)	1 (4%)	25 (100%)

Based on table 7, it is known that there are 15 pregnant women working in a cigarette factory with a gestational age in the first trimester who have normal hemoglobin levels with an average hemoglobin level of 12.8 g/dl.

Table 3. Hemoglobin levels based on length of work in pregnant women who work at the Djarum cigarette factory at the Mejobo Kudus Health Center.

Length of work	Hemoglobin			Total Amount (%)
	Normal Amount (%)	Anemia Light Amount (%)	Anemia Heavy Amount (%)	
< 2 yrs	11 (44%)	1 (4%)	1 (4%)	13 (52%)
2-5 yrs	9 (36%)	1 (4%)	0 (0)	10 (40%)
>5 yrs	2 (8%)	0 (0)	0 (0)	2 (8%)
Amount	22 (88%)	2 (8%)	1 (4%)	25 (100%)

Based on table 8, it is known that there are 11 pregnant women who work in cigarette factories with a length of service of <2 years who have normal hemoglobin levels with an average hemoglobin level of 13.0 g/dl.

Discussion

The results of examination of hemoglobin levels in pregnant women working in the cigarette factory at the Mejobo Health Center showed that out of 25 respondents, most of the respondents had an average hemoglobin level of 12.3 g/dl, where the majority of respondents had a normal hemoglobin of 88%. This shows that the majority of respondents did not have anemia. The results of the hemoglobin levels of the respondents based on the age of the pregnant women showed that the majority of respondents aged 20-40 years were normal or had hemoglobin levels > 11.00 g/dl, or did not have anemia. The results of the respondents' hemoglobin

levels based on the trimester of pregnancy showed that the majority of respondents with the first to third trimesters of gestation had normal hemoglobin levels, or did not experience anemia. The results of the respondents' hemoglobin levels based on the length of time pregnant women worked in cigarette factories showed that the majority of respondents with a length of service of <2 years to >5 years had normal hemoglobin levels, or did not experience anemia. Pregnant women who do not experience anemia even though they have worked in a factory for more than 5 years can occur because when working pregnant women use complete PPE, from head coverings, masks, and aprons to protect themselves.

In the study of examining hemoglobin levels of pregnant women who worked at the Djarum cigarette factory at the Mejobo Health Center, the incidence of anemia in pregnant women was still low, the percentage was 12%.

There are many ways that can be done to increase hemoglobin levels in pregnant women so that anemia does not occur which has a negative impact on the health of the mother and fetus in the womb, one of which is by using complete PPE when working in factories. Personal protective equipment is a must that must be carried out by the parties. Djarum factory management in each region. The goal is to reduce the number of work accidents that might occur. Personal protective equipment provided by the Djarum factory SKT unit.

For pregnant women, it is better to carry out routine control of the midwife. Have regular check-ups to control fetal growth and the health of pregnant women, especially routinely checking hemoglobin levels in the blood so that they can be detected earlier if anemia occurs and can be treated immediately, so as not to have a negative impact on the health of the mother and fetus in the womb.

Conclusion

1. The average age based on 20-29 years has a hemoglobin level of 12.8 g/dl, while those aged 30-39 years have an average hemoglobin level of 11.9 g/dl, those aged > 40 years have an average hemoglobin level of 12.6 gr/dl .
2. The mean based on gestational age in the first trimester has a hemoglobin level of 12.4 g/dl, whereas in the second trimester the gestational age has an average hemoglobin level of 12.3 g/dl, in the third trimester the gestational age has an average hemoglobin level of 13.3 gr/dl.

3. The mean based on length of work <2 years has a hemoglobin level of 11.4 g/dl, while those with a length of work of 2-5 years have an average hemoglobin level of 12.5 g/dl, for a length of work > 5 years have an average hemoglobin level of 12.6 g /dl.
4. The majority of respondents did not experience anemia with a normal hemoglobin level of 88%.

Reference

- Ageng, S. Z. 2020. Perbedaan Status Anemia Berdasarkan Konsumsi Zat Besi (Fe) dan Lama Menstruasi Pada Siswi SMK Negeri 1 Tabanan. <http://repository.poltekkes-denpasar.ac.id/4099/>. Diakses tanggal 26 Februari 2022.
- Andita, F. 2018. Analisa Faktor-Faktor yang Mempengaruhi Anemia Kehamilan di Puskesmas Padang Bulan. Karya Tulis Ilmiah. Universitas Sumatera Utara.
- Aulia, V. 2019. Gambaran Hasil Kadar Hemoglobin (Hb) pada Ibu Hamil Trimester I dan Trimester II di Puskesmas Muara Labuh. Karya Tulis Ilmiah. Sekolah Tinggi Ilmu Kesehatan Perintis Padang.
- Minarsih, F. N. N. 2018. Faktor-Faktor yang Berhubungan dengan Kejadian Anemia pada Ibu Hamil di Puskesmas Moramo Tahun 2018. Karya Tulis Ilmiah. Politeknik Kesehatan Kendari.
- Ningrum, D. P. 2014. Gambaran Hemoglobin pada Pekerja yang Terpapar Asap Pembakaran Sate. Karya Tulis Ilmiah. Universitas

Muhammadiya

- h Semarang.
- Prayogi, K. R. 2012. Evaluasi Pelaksanaan Program Kesehatan dan Keselamatan Kerja (K3) (Studi pada PT.Djarum Unit SKT Megawon II). Skripsi. Universitas Pembangunan Nasional "Veteran" Yogyakarta.
- Rumarjianti, S. I. 2015. Hubungan Antara Jenis Pekerjaan dengan Anemia Ibu Hamil Pekerja Pabrik Rokok di Puskesmas Rejosari Kabupaten Kudus Jawa Tengah. Tesis. Universitas Airlangga.
- Syafi'l, M. I. 2015. Gambaran Kadar Hemoglobin (hb) pada Pekerja Pabrik Daur Ulang Accu di Kota Semarang Akibat Paparan Timbal (pb). Karya Tulis Ilmiah. Universitas Muhammadiyah Semarang.
- Sri, L. 2018. Gambaran Kadar Hemoglobin pada Pedagang Sate di Kecamatan Gianyar. <http://repository.poltekkes-denpasar.ac.id/1118/>. Diakses tanggal 18 Maret 2022.
- Supini, S. 2018. Gambaran Kadar Hemoglobin Pada Ibu Hamil Trimester III dengan Anemia di Puskesmas Gatak Sukoharjo. Karya Tulis Ilmiah. Universitas Muhammadiyah Semarang.
- Suriani, L. 2019. Analisa Kadar Haemoglobin pada Ibu Hamil Trimester III di Puskesmas Pargarutan Tapanuli Selatan. Karya tulis ilmiah. Politeknik Kesehatan Medan.
- Utami, L. P. 2022. Pemeliharaan Hubungan Perusahaan Terhadap Karyawan di PT. Djarum Kudus. https://www.researchgate.net/publication/361361878_PEMELIHARAAN_HUBUNGAN_PERUSAHAAN_TERHADAP_KARYAWAN_DI_PT_DJARUM_KUDUS. Diakses tanggal 17 Juli 2022.
- Yulianti, Y. 2019. Faktor-Faktor yang Berhubungan dengan Ibu Hamil di Wilayah Kerja Puskesmas Karangayar Kota Tasikmalaya Tahun 2019. <http://repositori.unsil.ac.id/535/7/11%20OBAB%20II.pdf>. Diakses tanggal 20 Maret 2022.

Description of Hemoglobin Levels for Pregnant Women Workers of The Djarum Cigarette Factory at The Mejobo Kudus Health Center

ORIGINALITY REPORT

14%

SIMILARITY INDEX

4%

INTERNET SOURCES

13%

PUBLICATIONS

3%

STUDENT PAPERS

PRIMARY SOURCES

- 1 "1st Annual Conference of Midwifery", Walter de Gruyter GmbH, 2020
Publication 4%
- 2 K. E. Chan, R. A. Lafayette, A. S. Whittemore, M. A. Hlatky, J. Moran. "Facility factors dominate the ability to achieve target haemoglobin levels in haemodialysis patients", Nephrology Dialysis Transplantation, 2008
Publication 4%
- 3 Elya Susmita, Astriana Astriana, Lady Octaviani Iqmy, Zarma H. "The Factors Related To The Incidence Of Anemia In Pregnant Women", Jurnal Kebidanan Malahayati, 2023
Publication 2%
- 4 elearning.medistra.ac.id
Internet Source 1%
- 5 www.researchgate.net

Internet Source

1 %

6

Funda Elmacioglu. "Is adequate and balanced nutrition during pregnancy more effective than iron and folic acid supplements?", Central European Journal of Medicine, 10/22/2009

Publication

1 %

7

Budi Santosa, Herlisa Anggraini, Ika Dyah Mawarni, Niza Muliaturun, Ardiansyah Nugroho, Uly Syarah, Siti Atika Risqoeni. "Erythrocyte index of residents exposed to lead in Tambaklorok, Semarang, Indonesia", Jurnal Teknologi Laboratorium, 2020

Publication

1 %

8

www.ncbi.nlm.nih.gov

Internet Source

1 %

9

journal.walisongo.ac.id

Internet Source

1 %

Exclude quotes On

Exclude matches < 1%

Exclude bibliography On