

Ethanollic extract of the natural product of Daun sirih (Piper betle) leaves may impede the effectiveness of the plasma jet contact style for acute wounds

by Fitri Nuroini

Submission date: 10-Jul-2023 03:03PM (UTC+0700)

Submission ID: 2129017296

File name: 1._Ethanollic_extract.pdf (1.94M)

Word count: 6332

Character count: 33620



ELSEVIER

17
Contents lists available at ScienceDirect

Clinical Plasma Medicine

journal homepage: www.elsevier.com/locate/cpme

Original research article

4
Ethanol extract of the natural product of *Daun sirih* (*Piper betle*) leaves may impede the effectiveness of the plasma jet contact style for acute wounds



Heni Setyowati Esti Rahayu^{a,k}, N. Nasruddin^{b,j,k,*}, Laela Hayu Nurani^{c,k}, Sri Darmawati^{b,k}, Afiana Rohmani^d, Heni Lutfiyati^e, Eka Sakti Wahyuningtyas^{a,j,k}, Isabella Meliawati Sikumbang^{e,j}, Zaenul Muhlisin^f, Andri Sukeksi^b, Fitri Nuroini^b, Tatsuo Ishijima^g, Junko Sugama^{h,i}, Toshio Nakatani^h

10
^a Department of Nursing, Faculty of Health Sciences, Universitas Muhammadiyah Magelang, Indonesia

^b Department of Medical Laboratory Science, Faculty of Nursing and Health Sciences, Universitas Muhammadiyah Semarang, Indonesia

^c Department of Pharmacy, Faculty of Pharmacy, Universitas Ahmad Dahlan, Yogyakarta, Indonesia

^d Faculty of Medicine, Universitas Muhammadiyah Semarang, Semarang, Indonesia

^e Department of Pharmacy, Faculty of Health Sciences, Universitas Muhammadiyah Magelang, Magelang, Indonesia

^f Department of Physics, Universitas Diponegoro, Semarang, Indonesia

^g Research Center for Sustainable Energy and Technology, Kanazawa University, Kanazawa-shi, Japan

^h Division of Nursing, Faculty of Health Sciences, Institute of Medical, Pharmaceutical, and Health Sciences, Kanazawa University, Kanazawa-shi, Japan

ⁱ Inced Health Care Science Research Unit, Institute for Frontier Science Initiative (InFiniti), Kanazawa University, Kanazawa-shi, Japan

^j Research Center for Experimental Wound Healing, Universitas Muhammadiyah Magelang, Indonesia

^k Muhammadiyah Research Network for Plasma Medicine (M-Plasmed), Semarang, Indonesia

ARTICLE INFO

Keywords:

Plasma medicine
ROS
RNS
PAW
Natural product
Wound

ABSTRACT

6
Purpose: An investigation was carried out to determine the effect of an ethanol extract of the natural product of *Daun sirih* or *Piper betle* leaves on the effectiveness of plasma jet treatment for cutaneous acute wound healing in a small animal model mimicking a clinical setting.

Method: An atmospheric plasma jet using medical grade argon gas as a carrier gas was developed. The ethanol extract of *Piper betle* leaf (EPB) formulated. Optical emission spectroscopy and chemical methods were applied to evaluate the presence of reactive oxygen species (ROS) and reactive nitrogen species (RNS) in the gas phase and in aqueous and ethanol media. Small animals were classified into 5 groups, namely, Control (C), Plasma jet (P), Ethanol extract of *Piper betle* leaf (EPB), Plasma jet followed by EPB (P-EPB) and EPB followed by plasma jet (EPB-P). The contact and meander styles of plasma jet treatment for wounds were applied daily on acute wounds for 1 min, either alone or before or after EPB treatments. Visual evaluation of wounds was conducted for 14 days. Microscopic evaluation was conducted on days 7, 11 and 14. General staining, namely, haematoxylin-eosin and Azan staining, was conducted to evaluate neopithelialisation and new collagen formation.

Results: This research showed that wound healing in the P group was faster than that in the other groups, while that in groups containing EPB was the same as that in C. In the P group, the number of days to reach peak inflammation was the fewest. On day 7, neopithelialisation and new collagen formation in P were significantly higher than those in other groups.

Conclusion: Plasma jet treatment alone is able to promote inflammation, neopithelialisation and new collagen formation to accelerate acute wound healing; however, its admixture with EPB may impede such effectiveness. Based on the characterization of the ROS and RNS results, the ethanol solvent may play a primary role in impeding its effectiveness.

15
* Corresponding author. N. Nasruddin, SSI, MSi, PhD. Department of Medical Laboratory Science, Faculty of Nursing and Health Sciences, Universitas Muhammadiyah Semarang, Jalan Kedungmundu Raya No.18, Semarang, 50273, Central Java, Indonesia.

E-mail address: nasruddin@unimus.ac.id (N. Nasruddin).

26
<https://doi.org/10.1016/j.cpm.2019.100090>

Received 9 October 2018; Received in revised form 18 May 2019; Accepted 12 June 2019
2212-8166/© 2019 Published by Elsevier GmbH.

1. Introduction

It is well known that there are two main types of wounds, namely, acute and chronic wounds. Conceptually, optimal healing of acute wounds has been divided into several chronologically overlapping phases, namely, inflammation, proliferation, and remodeling [1]. Reactive oxygen species (ROS) and reactive nitrogen species (RNS) play a pivotal role in the orchestration of such processes [2,3]. To date, there have been a number of therapeutic strategies based on modulation of ROS function, namely, topical H₂O₂ (or related ROS intermediates [4–10], recombinant glucose oxidase [11], honey [12–17], galvanic particles [18,19], recombinant PDGF [20–23] and recombinant galectin-1 [24]. Those strategies, unfortunately, have drawbacks. Sen [25] reported that the uncontrolled kinetics of ROS/RNS release is the primary barrier to using ROS/RNS-donating compounds for topical applications. The development of plasma medicine technology producing pharmacologically appropriate doses of ROS/RNS would open a new window of opportunity in new clinical treatment methods [26].

Plasma medicine is a multidisciplinary science involving plasma science, life sciences, pharmacy, biomedical and other health sciences aimed at applying physical plasma in the field of human health therapy. The plasma referred to in this context is not blood plasma but physical plasma as a phase of the fourth state of matter, in addition to solid, liquid and gas. Plasma is commonly known as ionized gas because in the plasma phase, there are stable parts (gases) and reactive parts (ions, energetic and radical particles). Conceptually, the medical aspects of plasma are related to the ability of plasmas to generate biological molecules, namely, reactive oxygen species (ROS) and reactive nitrogen species (RNS), which if carefully controlled and in the right doses can be efficacious for health therapy [27–29]. A recent study involving a multidisciplinary team of researchers and lawyers reported that the style of action of a plasma medicine source, specifically cold atmospheric pressure plasmas, is pharmacological and not physical in nature [30].

A primary goal in medical plasma research is to gain insight into how to optimize medical plasma as a health therapy while minimizing its side effects. One approach that now attracts substantial attention from medical plasma researchers is the combination of medical plasma with compounds in the liquid phase (solution or liquid), which is known as the plasma-activated water (PAW) approach. Furthermore, Jablonowski et al. [31] emphasized that the concept of plasma-liquid interaction is a determining factor for the application of plasma in the medical field.

Based on previous research, it was reported that the jet type of medical plasma can accelerate the healing of acute full-thickness wounds in an in vivo setting by promoting inflammation, neopithelialisation and wound contraction [32]. Efforts to optimize medical plasma performance for wound healing using a combination of plasma and liquid have previously been performed [33–35]. Nasruddin et al. [33] reported that a combination of plasma jet and water in a micro-litre volume was able to significantly accelerate the healing of acute wounds by promoting wound contraction, compared to plasma treatment alone. Nasruddin et al. [34] also studied the effects of medical plasma treatment combined with low-concentration honey solutions supported by microwell dressings for wound healing, but the result was not significant. Wahyunings et al. [35] reported that the combination of plasma jet and Manuka honey is more effective than a combination of plasma jet and Indonesian honey for the improvement of acute wound healing.

The effectiveness of natural products, such as hormones [36–38] and honey [39], as complementary and alternative medicine to support wound healing has been reported, while the leaves of the *Piper betle* plant (locally known as *Daun Sirih*) have long been in use for the relief of pain in Indonesian indigenous medicine. *Piper betle* leaves contain active compounds, such as alkaloids and tannins, that have anti-bacterial and anti-inflammatory effects [40]. The effectiveness of the

ethanolic extract of the leaves to improve ulcer healing was previously reported [41]. It was hypothesized that the antioxidant or free radical scavenging activity of the plant extract may be correlated with its healing action. The present study was conducted to evaluate the effectiveness of a combination treatment of plasma jet and ethanolic extract of *Piper betle* leaves for acute wound healing using an experimental design mimicking a clinical setting in an animal model. The contact style of plasma jet treatment was applied in this investigation.

2. Methods and materials

2.1. Atmospheric pressure plasma jet system

An atmospheric pressure plasma jet system that was developed based on Teschke et al. [42], and described previously [34] was applied in this experiment. A modification of the dimensions of the capillary quartz tube was conducted. In this experiment, the inner and outer diameters of the tube were to be 0.65 mm and 1.55 mm, respectively. This material was fabricated by Beijing Zhong Cheng Quartz Glass Co., Ltd (Beijing, China). Two electrodes from an aluminum foil ring were applied around the quartz tube for this system. Non-conductive material was applied to isolate those two electrodes.

Electrical and optical emission characterizations were conducted at the Research Center for Sustainable Energy and Technology, Institute of Science and Engineering, Kanazawa University, Japan. The discharge voltage and discharge current were measured with a high-voltage probe (P6015A; Tektronix, Inc., Tokyo, Japan) and a current probe (8585C; Pearson Electronics, Palo Alto, CA, USA). When argon gas at a flow rate of 1 standard litre per minute (slm) was allowed to flow from one of the quartz tube, a low-frequency (~ 18.32 kHz) high AC voltage with a peak-to-peak voltage of 9.58 kV and current of 55.2 mA was measured at the upper ring electrode. Hospital grade ultra-pure argon gas (99.999% purity) (Samator Company, Indonesia) was used as a carrier gas.

Optical emission spectroscopy (OES) measurements taken approximately 10 mm under the nozzle showed the emissions of the OH (A-X transition) near 309 nm, N₂ (C-B transition) (band head maximum at 337 nm) [43] and Ar I (maximum 763 nm). This measurement showed the presence of both hydroxyl radical (OH) and nitrogen-based reactive species in the gas phase during its generation (Fig. 1).




Fig. 1. Optical emission spectroscopy (OES) identification of atmospheric plasma jet near 10 mm under the nozzle of the plasma jet reactor without mouse. OH and nitrogen-based reactive species were detected.




Fig. 2. Left: *Piper betle* leaves. Right: Ethanol extract of *Piper betle* leaves ready to be used.

2.2. Preparation of an ethanolic extract of *Piper betle* leaf and determination of its H_2O_2 and its NO_2

2.2.1. Preparation of ethanolic extract of *Piper betle* leaves

Fresh *Piper betle* leaves were collected locally and authenticated by the Laboratory of Pharmacy, Universitas Muhammadiyah Magelang, Indonesia. After being air dried and powdered in a hand crusher, the dried, powdered leaves (600 g) were extracted three times with absolute ethanol (1800 mL each time) and the resultant filtrate was solvent-recovered by distillation in a Soxhlet apparatus [44]. The concentrated extract, referred to as ethanolic extract of *Piper betle* leaf (EPB), was used as a stock solution for this experiment (Fig. 2).

2.2.2. Determination of H_2O_2 and NO_2

Concentrations of H_2O_2 and NO_2 pure water (A), plasma jet-treated pure water (A-P), EPB diluted in pure water (EPB-A), plasma jet-treated EPB-A (EPB-A-P), ethanol (E), plasma jet-treated ethanol (E-P), ethanolic extract of *Piper betle* leaves diluted in ethanol (EPB-E) and plasma jet-treated EPB-E (EPB-E-P) were analyzed with a method employing a peroxidase enzyme for H_2O_2 and a naphthylethylenediamine visual colorimetric for NO_2 using commercial reagent (Kyoritsu Chemical-Check Lab., Japan; Model WAK-H2O2, range: 0.05–5.0 mg/L and Model WAK-NO2, range: 0.02–1.00 mg/L) after plasma jet treatment. This method was also previously used by other researchers [33]. EPB-A was 1 ml EPB diluted in 299 ml pure water. E was 96% ethanol. EPB-E was 1 ml EPB diluted in 299 ml ethanol 96%. Ten millilitres of A, EPB-A, E and EPB-E in 10-ml beakers were treated by plasma jet for 2 min, perpendicularly. Before treatment, the distances between the nozzle of the plasma jet reactor and the surfaces of the objects were approximately 10 mm. Digital pocket test devices (Kyoritsu Chemical-Check Lab., Japan; Model DPM-H2O2 and Model DPM-NO2) were used to measure the concentrations of H_2O_2 and NO_2 .

2.3. Animals and investigation protocol

The investigation protocol and animal care were conducted based on the Guidelines for the Care and Use of Laboratory Animals at Laboratorium Penelitian dan Pengujian Terpadu/Integrated Research and Testing Laboratory (LPPT UGM), Gadjah Mada University, Yogyakarta, Indonesia (certificate number:00004/04/LPPT/III/2018). This institution works under the standard of the ISO/IEC 17,025 and the National Accreditation Committee of Indonesia (Komite Akreditasi Nasional/KAN, Indonesia). Forty-five Balb/c male mice aged 8 weeks and weighing 21.0–24.0 g purchased from Histology Laboratory, Universitas Sebelas Maret, Surakarta, Indonesia were used. The mice

were housed individually in a room with a temperature of 28.0 ± 2.0 °C and light from 09:00 to 21:00 h. They were allowed *ad libitum* access to feed and water.

2.4. Wound healing model and plasma treatment

The experimental animals, namely, Balb mice, were anaesthetized via the peritoneal cavity by injection of 50 mg/kg ketamine (K) and 5 mg/kg xylazine (X) [45]. The protocol to make an acute wound on a small animal as described previously [34] was adopted. This protocol was applied to create 2 disc-shaped (4 mm in diameter), full-thickness skin wounds, including the panniculus, on both sides of the mouse dorsum using a disposable 4 mm biopsy punch.

In this experiment, wound samples were distributed in five groups:

- Hydrocolloid dressing alone (Control group or C)
- Extract of *Piper betle* alone (EPB)
- Plasma alone (P).
- Plasma pursued by extract of *Piper betle* (P-EPB)
- Extract of *Piper betle* pursued by Plasma (EPB-P)

The wounds of the control group animals were covered with a hydrocolloid dressing (HD). Ethanolic extract of *Piper betle* was applied daily on animals in the experimental groups using a sprayer. The contact style of plasma jet used daily to treat the wound for 1 min. The wound surface was placed approximately 10 mm under the nozzle tip. To ensure that all areas of the wound surfaces were exposed in the plasma jet-treated groups, the meander style of plasma jet treatment was implemented. The experimental protocol is shown in Fig. 3.

2.5. Macroscopic evaluation of wound

The protocol for visual or macroscopic wound inspection described previously [34] was adopted in this experiment. These inspections were conducted for 14 days, with Day 0 being the day when the acute wounds were created. A digital camera (Panasonic, Lumix FH6) was used daily to document wound conditions.

2.6. Estimated day of healing

Using a graph of the ratio of the current wound area to the initial wound area, an estimated day of wound healing was determined from the y-axis when the trend of reduction of the wound size leveled off, as reported previously [34], at $y = 0.4$ (Fig. 6). From that point, line z was drawn, crossing the lines of reduction for each group. Wound healing




Fig. 3. Experimental protocol from day 0 until day13.

days at points c*, p*, epb*, p-epb* and epb-p* were calculated based on their points of intersection with the x-axis.

2.7. Tissue processing and histological analysis

On days 7, 11 and 14 after wound creation, the experimental

animals were massively euthanized by injection with ketamine-xylazine, administered via IP injection. Tissue surgery was conducted on the wound and its surrounding skin. Tissue processing was then conducted by means of [7] previously described technique [34]. The harvested wound tissue was stapled onto polypropylene sheets to prevent hyp

contraction of the samples and then fixed for approximately 24 h in

neutral-buffered 10% formalin solution at pH 7.4. Tissue bisection was carried out at the wound center. Subsequently, sections were dehydrated in an alcohol series, cleaned in xylene, and embedded in paraffin to prepare serial 5 µm sections. The sections containing the wound center were stained with haematoxylin-eosin (HE) and Azan for general histological analysis. On the basis of the results of haematoxylin-eosin staining from histological samples on days 7, 11 and 14 the percentage of neoeithelialisation was calculated using the following formula [46]:

$$\text{Neo-epithelialisation (\%)} = \frac{\text{length of new epithelium}}{\text{length of wound between wound edges}} \times 100\%$$

Using the results of Azan staining from histological samples on days 7 and 11, the percentage of new collagen formation was calculated using the following formula:

$$\text{New collagen formation (\%)} = \frac{\text{portion of new collagen}}{\text{portion of wound between wound edges}} \times 100\%$$

2.8. Statistical analysis

Data were subjected to statistical analyses using SPSS 16.0. The ratios of the average wound area to the original wound area, neoeithelialisation and new collagen formation were evaluated by ANOVA followed by the Tukey-Kramer method; *P* values <0.05 were considered significant.

3. Results

3.1. Determination of H₂O₂ and NO₂

In non-plasma-treated aqueous sample EPB-A and the plasma-treated aqueous samples (A-P and EPB-A-P), NO₂ was identified with different concentrations, while in all ethanolic samples, NO₂ was almost unidentified. While NO₂ was detected in EPB-A, its concentration was almost the same as that in A-P. H₂O₂ was identified in all ethanolic samples, but its rates varied. The H₂O₂ levels in the plasma-treated ethanolic samples (E-P and EPB-E-P) were higher than those in non-plasma-treated ethanolic samples (E and EPB-E). H₂O₂ in EPB-E-P, however, was lower than that in E-P. Finally, two points were hypothesized. First, in the ethanolic extract of *Piper betle* samples, plasma jet treatment increases H₂O₂ concentrations, but the extract of *Piper betle* may suppress them. At the same time, ethanol may reduce or eliminate the RNS produced by the plasma jet. Second, in the ethanolic extract of *Piper betle* diluted in pure water, plasma jet treatment decreases H₂O₂ but increases RNS. Water may act as a mediator of plasma jet treatment's production of RNS.

3.2. Macroscopic results

Fig. 5 shows the observed wounds on days 0, 3, 7, 11 and 14 for every group. The healing in all groups has a similar pattern, in which the wounds increased in size during the inflammation phase and then decreased gradually during the proliferative and maturation phases. The wound size on day 11 in the P group is smaller than that in the other groups. The yellowish color shown in the EPB, EPB-P and P-EPB images consists of residue of the ethanolic extract of *Piper betle* leaf.

3.3. Wound area reduction and estimated day of healing

All groups healed with a similar pattern. During the inflammation phase (from days 0 until 3, 4, or 5), the wounds increased dramatically, and then they decreased gradually until day 14. The peak inflammation days for the P group, the groups with ethanolic extract of *Piper betle*

(EPB, P-EPB and EPB-P) and the C group were days 3, 4 and 5, respectively. The wound area on the peak inflammation day in the P group (see A with a dash-lined circle) was larger than that in the groups treated with ethanolic extract of *Piper betle* (see B with a dash-lined circle), while the wound area in the P and EPB-treated groups was larger than that in the C group (see C with dash-lined circle). From days 3 to 14, the observed wound area in the P group was significantly smaller than those in the C group on days 6 and 9; those in the P group were significantly smaller than those in the P-EPB group on days 7, 8 and 9; those in the P group were significantly smaller than those in the EPB group on days 8 and 9; and finally, those in the P group were significantly smaller than those in the EPB-P group.

The day of wound healing for the P (p*) group was estimated on Day 10, whilst those for the C group (c*) and the groups treated with ethanolic extract of *Piper betle* (epb*, p-epb* and epb-p*) were estimated to occur the same day: Day 12. The prediction showed that the day of wound healing in the P group was earlier compared with those of other groups. Finally, it was indicated that the ethanolic extract of *Piper betle* leaf may reduce the effectiveness of atmospheric plasma jet to accelerate acute wound healing.

3.4. Neo-epithelialization

On days 7, 11 and 14 after wound creation, the percentages of neoeithelialisation for every group were calculated (Fig. 7). On day 7, the superiority of the neoeithelialisation percentage in the P group was apparent. The neoeithelialisation percentage in the P group was significantly higher than those of the other groups (P vs C = *P* < 0.05; P vs EPB = *P* < 0.05; P vs P-EPB = *P* < 0.01; P vs EPB-P = *P* < 0.05), while those in groups containing ethanolic extract of *Piper betle* leaf were not significantly different from the Control (C vs EPB = *P* > 0.05; C vs P-EPB = *P* > 0.01; P vs EPB-P = *P* > 0.05). From days 7 to 11, the neoeithelialisation percentages for all groups increased dramatically; however, on day 11, those for all groups did not differ significantly. On day 14, the wounds in the C, EPB and P groups were completely covered by neo epithelium, whilst those in the P-EPB and EPB-P groups were more than 90% covered by neoeithelialium. The neoeithelialisation percentages for all groups on that day, however, did not differ significantly.

3.5. Collagen formation

On days 7 and 11 after wound creation, the percentages of collagen formation for every group were calculated (Fig. 8). On day 7, the percentage of new collagen formation in the P group was significantly higher than those in other groups (P vs C = *P* < 0.05; P vs EPB = *P* < 0.01; P vs P-EPB = *P* < 0.01; P vs EPB-P = *P* < 0.01), while those in groups containing ethanolic extract of *Piper betle* leaf were not significantly different from the Control (C vs EPB = *P* > 0.05; C vs P-EPB = *P* > 0.01; P vs EPB-P = *P* > 0.05). On day 11, the percentage of new collagen formation for the P group was significantly higher than that for the P-EPB group, but those for the C, P and EPB-P groups did not differ significantly.

4. Discussion

It has been reported that there is an intimate connection between parameters and styles of atmospheric plasma jet treatment and its healing effectiveness [34]. Although plasma-activated water (PAW) has recently gained a great deal of attention in the study of plasma medicine, well-tested models of PAW applied in animal or human clinical settings are still rare. To the best of our knowledge, this is the first report to study the effectiveness of combinatory treatment of atmospheric plasma jet and ethanolic extract of natural product *Piper betle* leaf for acute wound healing, using an animal model mimicking a modern clinical setting and supported by hydrocolloid dressing as a




Fig. 4. Concentrations of H₂O₂ and NO₂ in pure water (A), plasma jet-treated pure water (A-P), ethanolic extract of *Piper betle* diluted in pure water (EPB-A), plasma jet-treated EPB-A (EPB-A-P), ethanol (E), plasma jet-treated ethanol (E-P), ethanolic extract of *Piper betle* leaf diluted in ethanol (EPB-E) and plasma jet-treated EPB-E (EPB-E-P).

means of keeping the wound environment moist.

Indications regarding the contact style of plasma treatment for burn wound were investigated previously [47]. Our prior research, however, has shown that plasma jet treatment employing the spot and contact styles with a duration of 5 min causes detrimental effects on the normal skin of a mouse, which was previously marked by an increase of more than 4 °Celsius in the local temperature, as shown by an infrared thermal camera [33,34]. It was also reported that the use of the contact style of plasma jet at a distance of 5 mm for 3 min impedes wound healing [48].

In the present study, the contact style of plasma treatment was also applied, alone and before/after EPB, but in order to reduce its detrimental effect, the meander style of treatment was applied. Weltmann et al. [49] reported that plasma jet treatment using a moving style was able to significantly minimize the temperature elevation. This investigation showed that there was no significantly different effect between the two styles of plasma treatment combined with EPB, namely, plasma followed by extract of *Piper betle* (P-EPB) and extract of *Piper betle* followed by plasma (EPB-P). This investigation, however, indicated that plasma jet alone accelerates wound healing more effectively than either combination. Neo-epithelialization and new collagen formation was significantly enhanced in the plasma jet alone group,

Group	Day 0	Day 3	Day 7	Day 11	Day 14
C					
EPB					
P					
EPB-P					
P-EPB					

Fig. 5. Macroscopically observed wounds on days 0, 3, 7, 11, and 14.




Fig. 6. Graph of ratio of wound area to initial wound area and estimated days of healing.




Fig. 7. Neo-epithelialisation percentage on days 7, 11 and 14 after wound creation. On day 7, the superiority of neoeithelialisation percentage in P was observed.




Fig. 8. Percentage of new collagen formation.

while plasma jet treatment combined with EPB tended to be similar to the control group. Based on this finding, it is hypothesized that plasma jet with parameters as applied in this research design is effective in improving wound healing, but such effectiveness can possibly be impeded by the use of ethanolic extract of *Piper betle* leaf. In this context, however, the ethanol solvent may be acritical factor in the reduction in effectiveness. As shown in the results (Fig. 4), ethanol solvent reduces RNS and increases ROS excessively. It was reported that low concentrations of RNS [28] and ROS [3] have efficacy for wound healing, while high levels of ROS clearly have the potential to be a complicating factor in the regeneration and remodeling of nascent tissue. Detrimental effects on wound healing from the exposure to acute ethanol were also studied previously. Radek et al. [50] reported that one of the most significant effects of ethanol exposure on wound healing occurs during the inflammatory response, and altered cytokine production is a main component. Ethanol exposure also disturbs the proliferative response during healing, causing delays in neoeithelialisation, collagen formation, and angiogenesis.

This experiment also revealed that plasma jet treatment alone decreases the time that it takes a wound to reach its peak day in the inflammatory phase. In the context of the wound area, however, the shortness of the time required to reach the peak inflammatory day in the P group was followed by the highest level of expansion of the wound area. This phenomenon may be caused by the effect of plasma jet treatment on macrophage and fibroblast cells. It is well known that these two types of cells are critical to acute wound healing. Macrophage cells are considered as biomarkers for the late inflammatory phase, whilst fibroblasts play a main role in the proliferative phase, being especially associated with new collagen and myofibroblast cells formations. It was reported that plasma jet treatment has the ability to enhance phagocytotic activity of macrophages [51] and to increase the proliferation of fibroblasts. Liu et al. [20] reported that argon plasma jets have the ability to promote fibroblasts by activating the NF- κ B pathway and increasing the expression of cyclinD1 [52]. In summary, this investigation showed that topical treatment consisting of plasma jet alone has the ability to accelerate acute wound healing via the promotion of inflammation, neoeithelialisation and new collagen formation; however, its admixture with EPB may impede such effectiveness. High concentrations of ethanol solvent may play a role as an obstructive

factor.

Conflicts of interest

None

49 Acknowledgment

18 This research was supported by the research grant program Penelitian Dasar Unggulan Perguruan Tinggi (PDUPT) the Ministry of Research, Technology and Higher Education, Republic of Indonesia (contract number: 004/K6/KM/SP2H/PENELITIAN/2018).

14 Supplementary materials

Supplementary material associated with this article can be found, in the online version, at doi:10.1016/j.cpm.2019.100090.

References

- [1] T.J. Shaw, P. Martin, Wound repair at a glance, *J. Cell Sci.* 122 (2009) 3209–3213.
- [2] C. Dunnill, T. Patton, J. Brennan, J. Barrett, M. Dryden, J. Cooke, D. Leaper, N.T. Georgopoulos, Reactive oxygen species (ROS) and wound healing: the functional role of ROS and emerging ROS-modulating technologies for augmentation of the healing process, *Int. Wound J.* 14 (2015) 89–96, <https://doi.org/10.1111/iwj.12557>.
- [3] C.K. Sen, S. Roy, Redox signals in wound healing, *Biochimica et Biophysica Acta* 1780 (2008) 1348–1361.
- [4] Klyubin IV, K.M. Kirpichnikova, I.A. Gamaley, Hydrogen peroxide induced chemotaxis of mouse peritoneal neutrophils, *Eur. J. Cell Biol.* 70 (1996) 347–351.
- [5] J.R. Stone, T. Collins, The role of hydrogen peroxide in endothelial proliferative responses, *Endothelium* 9 (2002) 231–238.
- [6] G.A. Murrell, M.J. Francis, L. Bromley, Modulation of fibroblast proliferation by oxygen free radicals, *Biochem. J.* 265 (1990) 659–665.
- [7] M.M. Shi, J.J. Godleski, J.D. Paulauskis, Regulation of macrophage inflammatory protein-1alpha mRNA by oxidative stress, *J. Biol. Chem.* 271 (1996) 5878–5883.
- [8] M. Cho, T.K. Hunt, M.Z. Hussain, Hydrogen peroxide stimulates macrophage vascular endothelial growth factor release, *Am. J. Physiol. Heart Circ. Physiol.* 280 (2001) H2357–H2363.
- [9] A.E. Loo, R. Ho, B. Halliwell, Mechanism of hydrogen peroxide-induced keratinocyte emigration in a scratch-wound model, *Free Radic. Biol. Med.* 51 (2011) 884–892.
- [10] E. Tur, L. Bolton, B.E. Constantine, Topical hydrogen peroxide treatment of ischemic ulcers in the guinea pig: blood recruitment in multiple skin sites, *J. Am. Acad. Dermatol.* 33 (2 Pt 1) (1995) 217–221.
- [11] V. Arul, J. Masilamoni, E. Jesudason, P. Jaji, M. Inayathullah, D.D. John, S. Vignesh, R. Jayakumar, Glucose oxidase incorporated collagen matrices for dermal wound repair in diabetic rat models: a biochemical study, *J. Biomater. Appl.* 26 (2012) 917–938, <https://doi.org/10.1177/0885328210390402>.
- [12] P.H. Kwakman, S.A. Zaat, Antibacterial components of honey, *IUBMB Life* 64 (2012) 48–55.
- [13] V. Arul, J. Masilamoni, E. Jesudason, P. Jaji, M. Inayathullah, D.D. John, S. Vignesh, R. Jayakumar, Glucose oxidase incorporated collagen matrices for dermal wound repair in diabetic rat models: a biochemical study, *J. Biomater. Appl.* 26 (2012) 917–938, <https://doi.org/10.1177/0885328210390402>.
- [14] A.B. Jull, N. Cullum, J.C. Dumville, M.J. Westby, S. Deshpande, N. Walker, Honey as a topical treatment for wounds, *Cochrane Database Syst. Rev.* (3) (2015) 1–128, <https://doi.org/10.1002/14651858.CD005083.pub4>.
- [15] M. Dryden, X. Hudgell, K. Saeed, A.W.S. Dryden, J. Brooks, J. Cooke, Surgihoney – Honey Wound Treatment: First Report of In-Vitro Activity and Early Clinical Evaluation, Federation of Infection Societies, Birmingham, UK, 2013.
- [16] J. Cooke, M. Dryden, T. Patton, J. Brennan, J. Barrett, The antimicrobial activity of prototype modified honeys that generate reactive oxygen species (ROS) hydrogen peroxide, *BMC Res. Notes* 8 (2015) 20.
- [17] M. Dryden, C. Goddard, A. Madadi, M. Heard, K. Saeed, J. Cooke, Bioengineered Surgihoney as an antimicrobial wound dressing to prevent caesarean wound infection: a clinical and cost-effectiveness study, *Br. J. Midwifery* 22 (2014) 23–27.
- [18] M. Dryden, C. Goddard, A. Madadi, M. Heard, K. Saeed, J. Cooke, Bioengineered Surgihoney as an antimicrobial wound dressing to prevent caesarean wound infection: a clinical and cost-effectiveness study, *Br. J. Midwifery* 22 (2014) 23–27.
- [19] S. Kaur, P. Lyte, M. Garay, F. Liebel, Y. Sun, J.-C. Liu, M.D. Southall, Galvanic zinc-copper microparticles produce electrical stimulation that reduces the inflammatory and immune responses in skin, *Arch. Dermatol. Res.* 303 (2011) 551–562.
- [20] K. Kaltalioglu, S. Coskun-Cevher, F. Tugcu-Demiroz, N. Celebi, PDGF supplementation alters oxidative events in wound healing process: a time course study, *Arch. Dermatol. Res.* 305 (2013) 415–422.
- [21] M. Sundaresan, Z.X. Yu, V.J. Ferrans, K. Irani, T. Finkel, Requirement for generation of H2O2 for platelet-derived growth factor signal transduction, *Science* 270 (1995) 296–299.
- [22] H.-D. Beer, M.T. Longaker, S. Werner, Reduced expression of PDGF and PDGF receptors during impaired wound healing, *J. Invest. Dermatol.* 109 (1997) 132–138.
- [23] C.-H. Heldin, B. Westermark, Mechanism of action and in vivo role of platelet-derived growth factor, *Physiol. Rev.* 79 (1999) 1283–1316.
- [24] Y.-T. Lin, J.-S. Chen, M.-H. Wu, I.-S. Hsieh, C.-H. Liang, C.-L. Hsu, T.-M. Hong, Y.-L. Chen, Galectin-1 accelerates wound healing by regulating the neuropilin-1/Smad3/NOX4 pathway and ROS production in myofibroblasts, *J. Invest. Dermatol.* 135 (2015) 258–268.
- [25] C.K. Sen, The general case for redox control of wound repair, *Wound Repair Regen.* 11 (2003) 431–438.
- [26] M.G. Kong, G. Kroesen, G. Morfill, T. Nosenko, T. Shimizu, J. van Dijk, J.L. Zimmermann, Plasma medicine: an introductory review, *New. J. Phys.* 11 (2009) 115012.
- [27] B. Haertel, T. Voedtk, K.-D. Weltmann, U. Lindequist, Non-thermal atmospheric-pressure plasma possible application in wound healing, *Biomol. Ther.* 22 (6) (2014) 477–490.
- [28] A. Soneja, M. Drews, T. Malinski, Role of nitric oxide, nitroxidative and oxidative stress in wound healing, *Pharmacol. Rep.* 57 (Suppl.) (2005) S108–S119.
- [29] K.-D. Weltmann, Th. Voedtk, Plasma medicine—current state of research and medical application, *Plasma Phys. Control Fusion* 59 (2017) 014031–014042.
- [30] A. Kramer, B.R. Conway, K. Meissner, F. Scholz, et al., Cold atmospheric pressure plasma for treatment of chronic wounds: drug or medical device? *J. Wound Care* 26 (8) (2017) 470–475, <https://doi.org/10.12968/jowc.2017.26.8.470>.
- [31] H. Jablonowski, T. Voedtk, Research on plasma medicine-relevant plasma-liquid interaction: what happened in the past five years? *Clin. Plasma Med.* 3 (2) (2015) 42–52.
- [32] Nasruddin, Y. Nakajima, K. Mukai, H.S.E. Rahayu, M. Nur, T. Ishijima, H. Enomoto, Y. Uesugi, J. Sugama, T. Nakatani, Cold plasma on full-thickness cutaneous wound accelerates healing through promoting inflammation, re-epithelialisation and wound contraction, *Clin. Plasma Med.* 2 (2014) 28–35.
- [33] Nasruddin, Y. Nakajima, K. Mukai, E. Komatsu, H.S.E. Rahayu, M. Nur, T. Ishijima, H. Enomoto, Y. Uesugi, J. Sugama, T. Nakatani, A simple technique to improve contractile effect of cold plasma jet on acute mouse wound by dropping water, *Plasma Process. Polymers* 12 (2015) 1128–1138.
- [34] Nasruddin, I.K. Putri, S. Kamal, H.S.E. Rahayu, et al., Evaluation the effectiveness of combinative treatment of cold plasma jet, Indonesian honey, and micro-well dressing to accelerate wound healing, *Clin. Plasma Med.* (2017) 14–25 5–6pp.
- [35] E.S. Wahyuningtyas, A. Iswara, Y. Sari, S. Kamal, et al., Comparative study on manuka and Indonesian honeys to support the application of plasma jet during proliferative phase on wound healing, *Clin. Plasma Med.* 12 (2018) 1–9.
- [36] K. Mukai, Y. Nakajima, T. Urai, et al., The effect of 17 β -Estradiol administration on cutaneous wound healing in 24-Week ovariectomized female mice, *J. Horm.* (2014), <http://dx.doi.org/10.1155/2014/234632>.
- [37] K. Mukai, Y. Nakajima, T. Urai, E. Komatsu, Nasruddin, et al., 17 β -estradiol administration promotes delayed cutaneous wound healing in 40-week ovariectomized female mice, *Int. Wound J.* 13 (2014) 636–644.
- [38] K. Mukai, E. Komatsu, Y. Nakajima, T. Urai, Nasruddin, et al., The effect of 17 β -Estradiol on cutaneous wound healing in protein-malnourished ovariectomized female mouse model, *PLoS ONE* 9 (12) (2014) e115564, <https://doi.org/10.1371/journal.pone.0115564>.
- [39] Y. Nakajima, K. Mukai, Nasruddin, E. Komatsu, T. Iuchi, Y. Kitayama, J. Sugama, T. Nakatani, Evaluation of the effects of honey on acute-phase deep burn wounds, *Evid.-Based Complement. Altern. Med.* 2013 (2013) 1–20 Vol <http://dx.doi.org/10.1155/2013/784959>.
- [40] D. Pradhan, K.A. Suri, D.K. Pradhan, P. Biswasroy, Golden heart of the nature: *piper betle*, *J. Pharmacogn. Phytochem.* 1 (2013) 147.
- [41] B. Majumdar, S.G.R. Chaudhuri, A. Ray, S.K. Bandyopadhyay, Effect of ethanolic extract of Piper betle Linn leaf on healing of NSAID-induced experimental ulcer—a novel of free radical scavenging action, *Indian J. Exp. Biol.* 41 (2003) 311–315.
- [42] M. Teschke, J. Kedzierski, E. Finantu-Dinu, D. Korzec, J. Engemann, High-Speed photographs of a dielectric barrier atmospheric pressure plasma jet, *IEEE Trans. Plasma Sci.* 33 (2005) 310–311.
- [43] M. Tsuji, J. Kumagai, K. Nakano, T. Matsuzaki, T. Tsuji, Decomposition of N₂O in a microwave-absorbent assisted discharge of N₂ at atmospheric pressure, *Appl. Surf. Sci.* 217 (2003) 134–148.
- [44] S. Bhattacharya, M. Subramanian, S. Roychowdhury, et al., Radioprotective property of the ethanolic extract of *Piper betle* leaf, *J. Radiat. Res.* 46 (2005) 165–171.
- [45] J.W. Carpenter, *Exotic Animal Formulary*, 4th ed., Elsevier, St. Louis, 2013 (Ed.).
- [46] Haryanto, T. Urai, K. Mukai, J. Sugama, T. Nakatani, Effectiveness of Indonesian honey on the acceleration of cutaneous wound healing: an experimental study in mice, *Wounds* 24 (2012) 110–119.
- [47] A.V. Nastuta, I. Topala, C. Grigoras, V. Pohoata, G. Popa, Stimulation of wound healing by helium atmospheric pressure plasma treatment, *J. Phys. D Appl. Phys.* 44 (2011) 105204.
- [48] S. Darmawati, A. Rohmani, L.H. Nurani, et al., When plasma jet is effective for chronic wound bacteria inactivation, is it also effective for wound healing? *Clin. Plasma Med.* 14 (2019) 100085.
- [49] K.-D. Weltmann, Th.v. Voedtk, Basic requirements for plasma sources in medicine, *Eur. Phys. J. Appl. Phys.* 55 (01) (2011) 13807.
- [50] K.A. Radek, M.J. Ranzer, L.A. DiPietro, Brewing complications: the effect of acute ethanol exposure on wound healing, *J. Leukoc. Biol.* 86 (5) (2009) 1125–1134.
- [51] A.B. Shekhter, V.A. Serezhnikov, T.G. Rudenko, et al., Beneficial effect of gaseous nitric oxide on the healing of skin wounds, *Nitric Oxide* 12 (2005) 2010–2019.
- [52] J.-R. Liu, G.-M. Xu, X.-M. Shi, G.-J. Zhang, Low temperature plasma promoting fibroblast proliferation by activating the NF- κ B pathway and increasing cyclinD1 expression, *Sci. Rep.* 7 (2017) 11698, <https://doi.org/10.1038/s41598-017-12043-w>.

Ethanol extract of the natural product of Daun sirih (Piper betle) leaves may impede the effectiveness of the plasma jet contact style for acute wounds

ORIGINALITY REPORT

23%

SIMILARITY INDEX

18%

INTERNET SOURCES

18%

PUBLICATIONS

4%

STUDENT PAPERS

PRIMARY SOURCES

1	123chauffage.fr Internet Source	2%
2	www.ncbi.nlm.nih.gov Internet Source	1%
3	www.woundsresearch.com Internet Source	1%
4	ridb.kanazawa-u.ac.jp Internet Source	1%
5	ajpheart.physiology.org Internet Source	1%
6	thousands-passed.xyz Internet Source	1%
7	Kanae Mukai, Tamae Urai, Kimi Asano, Yukari Nakajima, Toshio Nakatani. "Evaluation of Effects of Topical Estradiol Benzoate Application on Cutaneous Wound Healing in Ovariectomized Female Mice", PLOS ONE, 2016	1%

8 Kazuaki Ninomiya, Tatsuo Ishijima, Masatoshi Imamura, Takayuki Yamahara et al. 1 %
"Evaluation of extra- and intracellular OH radical generation, cancer cell injury, and apoptosis induced by a non-thermal atmospheric-pressure plasma jet", Journal of Physics D: Applied Physics, 2013
Publication

9 advances.umw.edu.pl 1 %
Internet Source

10 media.neliti.com 1 %
Internet Source

11 Avijit Sarkar. "An ethanolic extract of leaves of Piper betle (Paan) Linn mediates its antileishmanial activity via apoptosis", Parasitology Research, 05/2008 <1 %
Publication

12 onlinelibrary.wiley.com <1 %
Internet Source

13 www.kansaiuniversityreports.com <1 %
Internet Source

14 dds.sciengine.com <1 %
Internet Source

15 www.inabj.org <1 %
Internet Source

16 NATO Science for Peace and Security Series A
Chemistry and Biology, 2012. <1 %
Publication

17 repositorio.unesp.br <1 %
Internet Source

18 Choirul Anam, Idam Arif, Freddy Haryanto,
Rena Widita, Fauzia P Lestari, Kusworo Adi,
Geoff Dougherty. "A SIMPLIFIED METHOD
FOR THE WATER-EQUIVALENT DIAMETER
CALCULATION TO ESTIMATE PATIENT DOSE
IN CT EXAMINATIONS", Radiation Protection
Dosimetry, 2018 <1 %
Publication

19 Donghai Li, Guiling Li, Jing Li, Zhi-Qiang Liu,
Xuman Zhang, Yu Zhang, He-Ping Li.
"Promotion of Wound Healing of Genetic
Diabetic Mice Treated by a Cold Atmospheric
Plasma Jet", IEEE Transactions on Plasma
Science, 2019 <1 %
Publication

20 ebin.pub <1 %
Internet Source

21 Chandan K. Sen. "The general case for redox
control of wound repair", Wound Repair and
Regeneration, 11/2003 <1 %
Publication

22 Peter Bruggeman. "Characterization of a direct dc-excited discharge in water by optical emission spectroscopy", Plasma Sources Science and Technology, 05/01/2009

Publication

<1 %

23 Piu Saha, Rupashree Sen, Chellaram Hariharan, Deepak Kumar, Padma Das, Mitali Chatterjee. " Berberine chloride causes a caspase-independent, apoptotic-like death in promastigotes ", Free Radical Research, 2009

Publication

<1 %

24 jurnal.uii.ac.id

Internet Source

<1 %

25 journals.utm.my

Internet Source

<1 %

26 parsproje.com

Internet Source

<1 %

27 www.lifesciencesite.com

Internet Source

<1 %

28 "American Society of Nephrology", Kidney International, 01/1989

Publication

<1 %

29 Gui-Min Xu, Yue Hao, Ming-Yang Sun, Jin-Ren Liu, Xing-Min Shi, Guan-Jun Zhang. "Characteristics of Plasma Activated Medium Produced by Atmospheric Pressure Helium

<1 %

Plasma Jet and Its Selective Effect on Malignant Melanoma and Normal Fibroblast Cells", IEEE Transactions on Plasma Science, 2020

Publication

30

ecomoment.org

Internet Source

<1 %

31

journal.unnes.ac.id

Internet Source

<1 %

32

zero.sci-hub.se

Internet Source

<1 %

33

"Physical and Chemical Basis of Nonthermal Plasma", Elsevier BV, 2018

Publication

<1 %

34

Avinash Kundadka Kudva, Suresh Rao, Pratima Rao, Romith Periera et al. "Chapter 14 Piper betle Linn. in Cancer: Past, Present, and Future", Springer Science and Business Media LLC, 2018

Publication

<1 %

35

Sander Bekeschus, Axel Kramer, Anke Schmidt. "Gas Plasma-Augmented Wound Healing in Animal Models and Veterinary Medicine", Molecules, 2021

Publication

<1 %

36

www.researchsquare.com

Internet Source

<1 %

37 Bárbara Campos Jorge, Armando Jorge Júnior, Julia Stein, Ana Carolina Casali Reis et al. "Safety assessment of the ethanolic extract from Piper vicosanum yunck leaves in male rats", Phytomedicine Plus, 2022
Publication <1 %

38 Endre Szili, Nishtha Gaur, Sung-Ha Hong, Hirofumi Kurita, Jun-Seok Oh, Masafumi Ito, Akira Mizuno, Akimitsu Hatta, Allison Cowin, David Graves, Robert Short. "The assessment of cold atmospheric plasma treatment of DNA in synthetic models of tissue fluid, tissue and cells", Journal of Physics D: Applied Physics, 2017
Publication <1 %

39 Kimi Shimamura. "Relationship between lymphangiogenesis and exudates during the wound-healing process of mouse skin full-thickness wound", Wound Repair and Regeneration, 07/2009
Publication <1 %

40 pdffox.com
Internet Source <1 %

41 www.cambridge.org
Internet Source <1 %

42 www.imedpub.com
Internet Source <1 %

43

www.researcherslinks.com

Internet Source

<1 %

44

Jian Zhang, Xiaohang Fu, Wenhui Li, He Li, Zhiwei Ying, Xinqi Liu, Liduan Yin.

"Enhancement of nutritional soy protein and peptide supplementation on skin repair in rats", *Journal of Functional Foods*, 2020

Publication

<1 %

45

Jiaxin Yang, WeiNan Zeng, Ping Xu, Xiaoxue Fu, Xiaojuan Yu, Lu Chen, Feng Leng, Chao Yu, Zhangyou Yang. "Glucose-responsive multifunctional metal-organic drug-loaded hydrogel for diabetic wound healing", *Acta Biomaterialia*, 2021

Publication

<1 %

46

Kanae Mukai, Yuriko Miyasaka, Kana Takata, Tamae Urai, Yukari Nakajima, Emi Komatsu, Junko Sugama, Toshio Nakatani. "Comparison of the Cutaneous Wound Healing of Ovariectomized Mouse at 12 Weeks with That of SHAM and Estrogen-Administered Mice", *Journal of Hormones*, 2014

Publication

<1 %

47

Lucas Buzeli de Souza, Jennyffer Ione de Souza Silva, Leonardo Bagne, Amanda Tavares Pereira et al. "Argon Atmospheric Plasma Treatment Promotes Burn Healing by

<1 %

Stimulating Inflammation and Controlling the Redox State", Inflammation, 2020

Publication

48

Sander Bekeschus, Anke Schmidt, Helena Jablonowski, Lydia Bethge et al.

"Environmental Control of an Argon Plasma Effluent and Its Role in THP-1 Monocyte Function", IEEE Transactions on Plasma Science, 2017

Publication

<1 %

49

Tetsuya Taima, Md. Shahiduzzaman, Takaaki Ishizeki, Kohei Yamamoto, Makoto Karakawa, Takayuki Kuwabara, Kohshin Takahashi.

"Sexithiophene-Based Photovoltaic Cells with High Light Absorption Coefficient via Crystalline Polymorph Control", The Journal of Physical Chemistry C, 2017

Publication

<1 %

50

doaj.org

Internet Source

<1 %

51

www.onmedu.edu.ua

Internet Source

<1 %

52

Kolb, Juergen F., Amber M. Mattson, Chelsea M. Edelblute, Xiaolong Hao, Muhammad Arif Malik, and Loree C. Heller. "Cold DC-Operated Air Plasma Jet for the Inactivation of Infectious Microorganisms", IEEE Transactions on Plasma Science, 2012.

<1 %

53

Wei Shen, Hai-Ming Shi, Wei-Hu Fan, Xin-Ping Luo, Bo Jin, Yong Li. "The effects of simvastatin on angiogenesis: studied by an original model of atherosclerosis and acute myocardial infarction in rabbit", *Molecular Biology Reports*, 2010

Publication

<1 %

54

pdfs.semanticscholar.org

Internet Source

<1 %

55

storage.googleapis.com

Internet Source

<1 %

56

ub-ed.ub.uni-greifswald.de

Internet Source

<1 %

57

www.ccmu.edu.cn

Internet Source

<1 %

58

www.ijpsr.info

Internet Source

<1 %

59

www.jci.org

Internet Source

<1 %

60

Christopher Dunnill, Thomas Patton, James Brennan, John Barrett, Matthew Dryden, Jonathan Cooke, David Leaper, Nikolaos T Georgopoulos. "Reactive oxygen species (ROS) and wound healing: the functional role of ROS and emerging ROS-modulating technologies

<1 %

for augmentation of the healing process",
International Wound Journal, 2017

Publication

61

"Comprehensive Clinical Plasma Medicine",
Springer Science and Business Media LLC,
2018

Publication

<1 %

62

Serdar Tort, Fatmanur Tuğcu Demiröz, Şule
Coşkun Cevher, Sanem Sarıbaş, Candan
Özoğul, Füsün Acartürk. "The effect of a new
wound dressing on wound healing:
Biochemical and histopathological
evaluation", Burns, 2020

Publication

<1 %

Exclude quotes Off

Exclude matches Off

Exclude bibliography On