

Turnitin InBj Polimorfisme MTNR1B

by Dr, Yanuarita

Submission date: 17-Jan-2023 01:13PM (UTC+0700)

Submission ID: 1993983793

File name: InBj_Polimorfisme_MTNR1B.pdf (1.37M)

Word count: 5660

Character count: 28501

RESEARCH ARTICLE

Association of rs10830963 *MTNR1B* and rs841853 *SLC2A1* Polymorphism with Obesity on Type 2 Diabetes Patients: An Overview of Melatonin Receptor and Transporter

Yanuarita Tursinawati^{1,*}, Arum Kartikadewi¹, Ari Yuniastuti², R Susanti²

¹Faculty of Medicine, Universitas Muhammadiyah, Semarang, Jl. Kedungmundu No.18, Semarang 50273, Indonesia
²Faculty of Mathematics and Natural Sciences, Universitas Negeri Semarang, Sekaran, Gunung Pati, Semarang 50229, Indonesia

*Corresponding author. E-mail: yanuarita_tursina@unimus.ac.id

Received date: Dec 30, 2020; Revised date: Mar 1, 2021; Accepted date: Mar 4, 2021

Abstract

BACKGROUND: One of the hormones that plays a role in glucose metabolism of Type 2 Diabetes Mellitus (T2DM) is melatonin. Its genetic variation is believed to play a significant role in the pathophysiology of obese and non-obese T2DM. The role of *MTNR1B* (melatonin receptor coding gene) and *SLC2A1* (Glucose transporter 1/GLUT 1 transporter coding gene) on the risk of obese and non-obese T2DM patients is controversial. This study aims to analyze the association between the rs10830963 *MTNR1B* and rs841853 *SLC2A1* polymorphism to the risk of Javanese obese T2DM.

METHODS: This was a cross-sectional study that involved 107 Javanese T2DM patients from primary health care in Semarang. Furthermore, obese T2DM was defined by a body mass index (BMI) more than 25 kg/m². The genetic variations examined were rs10830963 *MTNR1B* and rs841853 *SLC2A1* polymorphism by PCR-RFLP methods. Blood biochemistry parameters were also examined. Allele

and genotype frequencies of rs10830963 *MTNR1B* and rs841853 *SLC2A1* polymorphisms were analyzed using the χ^2 test with $p \leq 0.05$ and 95% CI.

RESULTS: There was a significant association between rs10830963 *MTNR1B* polymorphisms in obese and non-obese T2DM ($p=0.044$) and the CG genotype increased the risk of obese T2DM. Furthermore, the allele and genotype frequency of rs841853 *SLC2A1* polymorphism in both group had no significant difference, with a $p=0.756$ and $p=0.802$, respectively. There was also no significant difference in the biochemical parameters' in both groups of the two genetic variants studied.

CONCLUSION: The rs10830963 *MTNR1B* polymorphism is associated with the risk of obesity in Javanese T2DM patients but not for the rs841853 *SLC2A1* polymorphism.

KEYWORDS: polymorphism, *MTNR1B*, *SLC2A1*, obese, diabetes mellitus

Indones Biomed J. 2021; 13(2): 155-62

Introduction

Diabetes mellitus (DM), which is characterized by chronic hyperglycemia is a non-communicable disease and becomes major problem in Indonesia. According to data, the total number of DM patients in 2019 was around 120,835 people, these include those that received health services by a standard of around 2,066 or 1.71%. (1) Patients with type 2 DM (T2DM) experience abnormalities of blood glucose

level homeostasis characterized by an increase of fasting plasma glucose (FPG).

Melatonin is one of the hormones that play a role in the metabolism of carbohydrate and regulates plasma glucose. Moreover, a decrease in the level of melatonin and an increase of insulin in Goto Kakizaki rats was found in T2DM. (2) The melatonin hormone works according to its target organ by binding to melatonin receptors, namely MT1, which is coded for *MTNR1A* and *MTNR2* and encoded by *MTNR1B* gene. Furthermore, melatonin works across

the plasma membrane in cells mediated by a transporter, namely Glucose transporter 1 (GLUT 1) coded by the *SLC2A1* gene.(3,4)

Genome-wide Association Studies (GWAS) proved that there is a relationship between *MTNR1B* polymorphisms and FPG, insulin secretion, and the T2DM incidence.(5,6) This polymorphism occurs when C allele of the intron on of chromosome 11q21-q22 changes to G, which correlates with the risk of T2DM. The rs10830963 *MTNR1B* polymorphism was reported to be associated with increased FPG and T2DM in Caucasians, Chinese, Japanese, and Sri Lanka.(7,8,9) Another genetic variation regarding the role of melatonin in DM is the rs841853 *SLC2A1* polymorphism (XbaI G>T) which has been shown to double the risk of T2DM in Asia and Tunisian populations.(10)

The characteristics of T2DM patients vary widely. Classic T2DM patients usually have an obese condition. However, T2DM currently occurs in individuals with thin or normal body mass index (BMI). Lean T2DM patients show a faster loss of pancreatic beta-cell function than obese and therefore require earlier insulin treatment.(11,12) Studies in East Asia have shown that the onset of T2DM events is faster in individuals with lower BMI than in Caucasian. (11) Genetic involvement is believed to play a role in the T2DM pathogenesis of obese and non-obese patients. The genetic variations related to insulin secretion such as single-nucleotide polymorphism (SNP) *TCF7L2* are at risk of normal (lean) T2DM BMI. However, those related to insulin sensitivity such as SNP fat mass obesity (FTO) are at risk of T2DM obesity.(13-15) SNP *CDKN2BAS*, *CDKAL1* was associated with the risk of T2DM incidence in normal (lean) people.

Melatonin deficiency plays a role in obesity pathophysiology by regulating energy balance. It regulates energy intake, storage, and expenditure from adipose tissue, which determines a person's body weight. When the intake exceeds energy expenditure, it becomes overweight or obese.(16,17) Obesity is a risk factor for T2DM incidence. A study involving obese subjects proved that GG genotype of polymorphism rs10830963 *MTNR1B* gene was associated with an increase in fasting glucose levels.(18) There was an association between XbaI restriction fragment length polymorphism (RFLP) GLUT 1 in T2DM obese women. This indicates that gene plays a role in the incidence of T2DM in obese people.(19)

The involvement of rs10830963 *MTNR1B* polymorphisms and rs841853 *SLC2A1* in the risk of T2DM or obesity have been reported. However, there are no reports

on the association of these genetic variations in obese and non-obese T2DM individuals. This study aims to analyze the relationship between rs10830963 *MTNR1B* and the rs841853 *SLC2A1* polymorphism to the risk of obese and non-obese T2DM individuals of Javanese ethnicity, which is the largest ethnic group in Indonesia and the underlying clinical chemical parameters. The results will provide an overview of the contribution of genetic variation to the T2DM patients' heterogeneity based on BMI.

Methods

Subject Collection

This was an analytic observational study with a cross-sectional approach involved 107 T2DM patients from 3 health care centers in Semarang. T2DM patients that met the inclusion criteria were consecutively sampled, including Javanese T2DM patients, aged 30-70 years and had signed informed consent. The exclusion criteria include a history of cardiovascular disorders such as stroke, heart failure, and acute myocardial infarction. Obese T2DM was defined by a BMI of more than 25 kg/m², according to the BMI classification of the Asia Pacific population. Ethical permission was obtained from the Ethics Commission of the Faculty of Medicine, the Muhammadiyah University of Semarang with Number 040/EC/FK/2019.

Data Collection

The weight and height of the samples were measured to calculate the BMI. Blood pressure was measured at rest and classified into 5 classes, Normal: Systolic Blood Pressure (SBP) <120 mmHg or Diastolic Blood Pressure (DBP) <80 mmHg; Prehypertension: SBP 120-139 mmHg or DBP 80-89 mmHg; Hypertension grade 1: SBP 140-159 mmHg or DBP 90-99 mmHg; Hypertension grade 2: SBP ≥160 mmHg or DBP ≥100 mmHg, according to The Seventh Report of The Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure (JNC VII).(20) Venous samples were collected on ethylenediaminetetraacetic acid (EDTA) and plain tubes for the examination of gene polymorphism and biochemical parameters, respectively, such as fasting blood sugar, cholesterol, triglycerides, high-density lipoprotein cholesterol (HDL-C), low-density lipoprotein cholesterol (LDL-C), Cholesterol/HDL ratio (Chol/HDL), urea, and creatinine. Microalbumin examination was carried out using the subject's urine sample.

Biochemical Profile Examination

Respondents had fasted for 8 hours before collecting their venous blood to be examined for guanosine-diphosphatase (GDP) by an enzymatic method (GOD-PAP), ureum by Urease-gl²⁹amate dehydrogenase (GLDH), creatinine using the Jaffe, total cholesterol using CHOD-PAP, triglycerides using a GPO-PAP, HDL-C, and LDL-C by direct enzymatic method. Microalbumin was examined using respondents' urine samples with the immunoturbidimetric assay method and measured using a Cobas c311 autoanalyzer (Roche, Basel, Switzerland) at a wavelength of 340 nm.

6 Genotyping of *MTNR1B* Gene and *SLC2A1* Polymorphism

DNA was isolated from the blood using GeneJETM Genomic DNA Purification Kit. Amplification of the *MTNR1B* gene was done using Primer forward 5'-ATG CTA AGA ATT CAC ACC AG C T-3', reverse 5'-CAC AGT GCA GAC TGT TTT CTA ATC. Polymerase chain reaction (PCR) with predenaturation of 95°C, 7 minutes, 33 cycles: 95°C denaturation, 30 seconds, 60°C annealing, 30 seconds, 72°C elongation, 1 minute. The final extension was at 72°C for 3 minutes. RFLP method with PvuII enzyme was used to digest PCR product and visualized at 4% agarose gel. The electrophoresis results showed the C allele of 125 bp and G at 105 bp and 20 bp.

12 Amplification of *SLC2A1* was done using Primer forward 5'-TGC AAC CCA TGA GCT AAC AA-3' and reverse 5'-GAA CCC AGC ACT CTG TAG C-3'. PCR with predenaturation at 95°C, 5 minutes, 33 cycles: 95°C denaturation, 30 seconds, 60°C annealing, 30 seconds, 72°C elongation, 1 minute. The final extension was at 72°C for 3 minutes. Then, PCR product was digested with XbaI enzyme and visualized at 2% agarose gel. Electrophoresis results showed the T allele of 305 bp and G at 232 bp and 73 bp.

Data Analysis

The data for the existence of rs10830963 *MTNR1B* and rs841853 *SLC2A1* polymorphisms in the form of allele and genotype frequencies were analyzed by χ^2 test with $p < 0.05$ and CI=95%. The differences in the numerical characteristics of variables between groups were determined by the independent t-test.

Results

A total of 107 T2DM subjects, which consist of 45 obese and 62 non-obese were involved in the study. In the obese group, the majority of the sample was ≥ 60 years old (46.7%), female (82.2%), uncontrolled blood sugar levels

24 Table 1. Characteristic data of T2DM subjects in obese and non-obese groups.

Characteristics	n (%)		p-value*
	Obese (n=45)	Non-Obese (n=62)	
Age (year)			
30-34	1 (2.2%)	0 (0%)	0.660
35-39	0 (0%)	0 (0%)	
40-44	2 (4.4%)	1 (1.6%)	
45-49	3 (6.7%)	4 (6.5%)	
50-54	7 (15.6%)	7 (11.3%)	
55-59	11 (24.4%)	21 (33.9%)	
42 ≥ 60	21 (46.7%)	29 (46.8%)	
Gender			
Male	8 (17.8%)	17 (27.4%)	0.245
1 Female	37 (82.2%)	45 (72.6%)	
Fasting Plasma Glucose Controlling			
Uncontrolled (<90 or >130 mg/dL)	29 (64.4%)	47 (75.8%)	0.201
Controlled (90-130 mg/dL)	16 (35.6%)	15 (24.2%)	
Hypertension			
Normal	3 (6.7%)	7 (11.3%)	0.585
Prehypertension	20 (44.4%)	28 (45.2%)	
Hypertension grade 1	18 (40.0%)	18 (29.0%)	
Hypertension grade 2	4 (8.9%)	9 (14.5%)	

* Tested with Chi square test, $p < 0.05$ as significant result.

Table 2. Biochemical profile of T2DM subjects in obese and non-obese.

Biochemical Profile	Obese		Non-Obese		p-value*
	Mean±SD	Median (Min-Max)	Mean	Median (Min-Max)	
FPG (mg/dL)	155.02±9.94	139 (63-332)	164.24±10.63	141 (60-418)	0.752
HbA1c (mmol/mol)	7.82±0.27	7.2 (5.5-12.5)	7.88±0.26	7.45 (4.9-15.4)	0.847
Cholesterol (mg/dL)	165.09±8.99	160 (41-266)	194.08±5.99	199.5 (58-314)	0.018*
Triglycerides (mg/dL)	209.96±11.86	194 (78-402)	202.34±21.78	164 (61-1340)	0.060
HDL (mg/dL)	58.62±2.24	58 (33-97)	61.65±2.32	61 (33-113)	0.456
LDL (mg/dL)	201.31±8.47	199 (87-343)	196.50±6.96	204 (88-311)	0.786
Chol/HDL ratio	4.91±0.21	4.6 (2.8-9.7)	4.78±0.17	4.6 (2.6-9.2)	0.645
Ureum (mg/dL)	28.87±1.70	25 (15-61)	30.40±1.48	28 (15-80)	0.441
Creatinine (mg/dL)	0.88±0.06	0.76 (0.44-2.49)	0.96±0.63	0.89 (0.38-3.41)	0.362
Microalbumin (mg/dL)	241.89±54.93	50.2 (5-1582)	281.27±132.03	40.6 (4.2-8087.2)	0.202

* Tested with Mann Whitney U test, $p < 0.05$ as significant result.

(64.4%), and prehypertension (44.4%). Meanwhile, in the non-obese, the majority was ≥ 60 years old (46.8%), female (72.6%), uncontrolled blood sugar levels (75.8%), and prehypertension (45.2%) as shown in Table 1.

Based on the comparison results of the biochemical profile in Table 2, there was a significant difference between the obese and non-obese group in cholesterol levels with p -values of 0.018. However, there was no significant differences in FPG, HbA1c, Triglycerides, HDL, LDL Chol/HDL ratio, Ureum, creatinine, and microalbumin with $p > 0.05$.

Table 3 shows that in both groups, the majority of respondents with rs10830963 *MTNR1B* polymorphism had the C allele which was the wild type allele. Furthermore,

84.8% had CG genotype in the obese and 62.9% in the non-obese group. There was no significant difference between C and G alleles in both groups. However, the number of individuals with CG genotype was significantly higher than the CC in both groups with a p -value of 0.044. There was also a significant relationship between rs10830963 *MTNR1B* polymorphisms with obese and non-obese T2DM types where the CG genotype increased the risk of obesity. Meanwhile, in the rs841853 *SLC2A1* polymorphism, most of the allele was (wild type) and the most genotype was TT (45.2%) in the non-obese group. Moreover, the obese group found the same number in the TG and TT genotypes. There were no significant differences in the allele frequency and genotype of rs841853 *SLC2A1* polymorphism in the

Table 3. Allele distribution of rs10830963 *MTNR1B* and rs841853 *SLC2A1* polymorphisms.

		n (%)		p-value*
		Obese (n=45)	Non-Obese (n=62)	
rs10830963 <i>MTNR1B</i>				
Allele	C	52 (57.8%)	83 (66.9%)	0.171
	G	38 (42.2%)	41 (33.1%)	
Genotype	CC	7 (15.6%)	22 (35.5%)	0.044*
	CG	38 (84.4%)	39 (62.9%)	
	GG	0 (0%)	1 (1.6%)	
rs841853 <i>SLC2A1</i>				
Allele	G	54 (60.0%)	77 (62.1%)	0.756
	T	36 (40.0%)	47 (37.9%)	
Genotype	GG	9 (20.0%)	13 (21.0%)	0.802
	TG	18 (40.0%)	21 (33.9%)	
	TT	18 (40.0%)	28 (45.2%)	

* Tested with Chi square test, $p < 0.05$ as significant result.




Figure 1. PCR-RFLP electrophoresis result of rs10830963 *MTNR1B* polymorphism. A, B, G, and H were CG heterozygotes; C, D, E, F, and I were CC homozygotes, while J was GG homozygotes.

14 obese and non-obese groups, with a *p*-value of 0.756 and 0.802 respectively.

The electrophoresis results of PCR-RFLP were shown in Figure 1 and Figure 2 for the *MTNR1B* and *SLC2A1* gene, respectively. In Figure 1, it is seen that the letters A, B, G, and H had 125, 105 and 20 bp, respectively which were the CG (heterozygote) genotypes. C, D, E, F, and I had 125 bp (genotype CC) which were wild type alleles. J was GG homozygotes (105 and 20 bp). Meanwhile, in Figure 2, A, F, H, and I had 232 and 37 bp (genotype GG), B, C, E, G had 305, 232 and 37 bp (genotype TG) and D had 305 bp which was the TT genotype (homozygote risk allele T).

When comparing the biochemical profile results based on genotype on rs10830963 *MTNR1B* polymorphism between the obese and non-obese groups, it was found that there was no significant difference in biochemical parameters. However, there were higher but insignificant levels in the CG + GG genotype for FPG, HbA1c, Cholesterol, Chol/HDL ratio, ureum, and microalbumin in the obese group. Likewise, with non-obese group, levels of FPG, HbA1c, Cholesterol, HDL-C, LDL-C, ureum, creatinine, and microalbumin were higher in the CG + GG genotype (Table 4).

Table 5 showed the comparison of biochemical profile parameters based on the genotype of the rs841853 *SLC2A1* polymorphism and shows that there was no significant difference between the values of the GG and TG + TT parameters in both groups.




Figure 2. PCR-RFLP electrophoresis results of rs841853 *SLC2A1* polymorphism. A, F, H, I are the GG genotypes; B, C, E, G are the TG genotypes and D was the TT genotypes (homozygote risk allele T).

Discussion

14 clinical characteristics of T2DM patients may vary between obese and non-obese. This study analyzed the rs10830963 *MTNR1B* polymorphism in obese and non-obese T2DM individuals. The results indicate a significant relationship between rs10830963 *MTNR1B* polymorphisms in obese T2DM where the CG genotype increases the risk of T2DM obesity. Thus, this is contrary to research which identified the gene variations among DM obese and lean Han Chinese populations where this polymorphism was not associated with the risk of BMI in T2DM patients.(13) GG genotype, which has the risk of G allele is associated with a decreased beta-cell function (HOMA-B) and an increased insulin resistance (HOMA-IR) in adults.(18) Although there is a study showing that there is no correlation between HOMA-IR as a marker of insulin resistance and obesity.(21) Previous genomic loci associated with the risk of predisposing the clinical characteristics of obese and non-obese T2DM have been identified. A study conducted in Medan, Indonesia proved that the genetic variations such as FTO rs9939609 is associated with the risk of Chinese children obese subjects. (22) The pathophysiology in obese patients is more towards insulin resistance, whereas in lean patients or non-obese it is caused by defective-insulin release or impaired insulin secretion. Meanwhile, the genetic variations that regulate insulin secretion (CDC123/CAMK1D, CDKAL1, TCF7L2,

Table 4. The comparison of biochemical profile based on the genotype of rs10830963 *MTNR1B* polymorphism.

Biochemical Profile	Mean±SD Obese (n=45)		p-value	Mean±SD Non-Obese (n=62)		p-value
	CC	(CG) + (GG)		CC	(CG) + (GG)	
FPG (mg/dL)	123.43±16.49	160.84±11.19	0.167	146.86±15.03	173.80±14.17	0.219
HbA1c (mmol/mol)	6.94±0.47	7.99±0.31	0.177	7.29±0.35	8.21±0.35	0.053
Cholesterol (mg/dL)	160.86±25.49	165.87±9.71	0.795	186.18±11.97	198.43±6.59	0.365
Triglycerides (mg/dL)	211.43±20.62	209.68±13.62	0.725	209.68±54.77	198.30±16.14	0.151
HDL (mg/dL)	63.14±7.36	57.79±2.30	0.529	59.91±4.00	62.60±2.87	0.410
LDL (mg/dL)	224.57±29.48	197.03±8.48	0.415	189.18±11.34	200.53±8.85	0.431
Chol/HDL ratio	4.74±0.54	4.94±0.23	0.702	4.54±0.29	4.92±0.21	0.131
Ureum (mg/dL)	27.00±4.73	29.21±1.84	0.509	26.86±2.09	32.35±1.94	0.061
Creatinin (mg/dL)	0.95±0.26	0.87±0.06	0.635	0.90±0.09	0.99±0.08	0.556
Microalbumin (mg/dL)	97.61±46.98	268.46±63.72	0.347	112.96±47.11	373.84±202.45	0.332

HHEX, CDKN2BAS, TCF7L2, KCNJ11) are associated with T2DM risk in lean or non-obese subjects.(14,23-25) Another study which examined the SNP-19 Calpain-10 (CAPN10) polymorphisms proved that this polymorphism is associated with Javanese ethnic vulnerability to T2DM. (26) Abnormalities in CAPN10 can reduce secretion in pancreatic cells which according to research plays a role in the pathophysiology of T2DM in non-obese patients.

The results showed that those with CG genotype were significantly higher than CC in both groups. Obesity and insulin sensitivity have a modulatory effect on the prediction of genetic susceptibility in the T2DM incidence. (27) A cohort study among obese respondents demonstrated that there was a significant relationship between SNP rs10830963 *MTNR1B* and the prevalence of prediabetes or Impaired Glucose Tolerance (IGT).(28) Therefore, it is presumed that this polymorphism is significantly associated with T2DM-induced obesity.

Other results of this study prove that there is no significant difference of allele and genotype frequency of rs841853 *SLC2A1* polymorphism between 2 groups. The insignificant results in this study is due to ethnic differences, considering that the *SLC2A1* polymorphism is also influenced by ethnicity.(29) Allele 1 of GLUT 1 is associated with T2DM especially in overweight/obese women.(19) There is an increase in proinflammatory in the obese, while proinflammatory increases the GLUT1 (*SLC2A1*) expression in macrophages. This increases glucose uptake as well as metabolism and ultimately causing hyperinflammatory conditions.(30) Other studies have also shown a correlation between inflammatory markers such as high sensitivity C-reactive protein (hsCRP) in central obese elderly men.(31)

Other results also proved that the risk allele G contained in the CG + GG genotype increases the levels of FPG, HbA1c, cholesterol, ureum, and microalbumin in

Table 4. The comparison of biochemical profile based on the genotype of rs841853 *SLC2A1* polymorphism.

Biochemical Profile	Mean±SD Obese (n=45)		p-value	Mean±SD Non-Obese (n=62)		p-value
	GG	(TG) + (TT)		GG	(TG) + (TT)	
FPG (mg/dl)	159.22±14.86	152.22±13.48	0.677	161.39±17.55	166.59±13.17	0.515
HbA1c	8.09±0.42	7.65±0.37	0.215	7.61±0.39	8.11±0.36	0.520
Cholesterol (mg/dl)	174.50±12.42	158.81±12.52	0.292	194.14±8.88	194.03±8.26	0.832
Triglycerides (mg/dl)	215.72±20.16	206.11±14.77	0.719	199.00±17.82	211.68±37.13	0.904
HDL (mg/dl)	56.28±3.41	60.19±2.97	0.501	61.54±3.57	61.74±3.09	0.944
LDL(mg/dl)	199.28±14.88	202.67±10.28	0.899	203.75±10.23	190.53±9.51	0.380
Chol/HDL ratio	4.76±0.27	5.02±0.30	0.898	4.95±0.28	4.64±0.22	0.308
Ureum (mg/dl)	26.61±2.18	30.37±2.43	0.313	32.82±2.57	28.41±1.64	0.183
Creatinin (mg/dl)	0.76±0.06	0.96±0.09	0.211	1.00±0.11	0.92±0.07	0.529
Microalbumin (mg/dl)	197.45±89.48	271.511±70.25	0.271	162.53±43.54	379.06±238.41	0.651

both groups, although ⁵⁵ significant. The risk allele G, which increases FPG is consistent ⁶ with previous study. (18) In addition, the expression of the *MTNR1B* gene in pancreatic beta islet cells has a direct effect on regulating FPG levels. Therefore, when polymorphisms occur in this gene, it directly affects FPG. Furthermore, individuals at risk of allele G experience hepatic insulin resistance due to a decrease in suppression of hepatic glucose production during euglycemic hyperinsulinemia. This is also due to an increase in the hepatic glucose production rate, which increases gluconeogenesis. (8) In non-obese T2DM patients, there was also an increase in basal production of hepatic glucose, which ⁴⁷ resulted in an increase in FPG levels. In the obese group, cholesterol and LDL levels were higher in the allele G risk group. Other studies that analyze *MTNR1B* polymorphisms in relation to lipid profiles, stated that cholesterol and LDL have a relationship with the rs3781637 *MTNR1B* polymorphism. (32)

Furthermore, there is no significant difference in the biochemical parameters between GG and TG + TT of rs841853 *SLC2A1* polymorphism. However, the values of FPG, HbA1c, triglycerides, HDL, and microalbumin are higher in the TG + TT group. A high microalbumin levels in T2DM patients indicate a complication of DM nephropathy. The diagnosis of DM nephropathy is carried out when the urinary albumin level is >300 mg/24 hours with or without an increase in serum creatinine (>1.3 mg/dL). In the non-obese group, a mean microalbumin >300 mg/dL was found, which indicates the risk of ⁹ nephropathy. (33) Furthermore, several studies have shown a relationship between the rs841853 *SLC2A1* polymorphism and the risk of DN which is marked by an increase in microalbumin. (34) TT haplotype was found in nephropathy patients as reported ¹⁷ by previous study. (35) However, this is not proven by the results of this study.

Conclusion

There is a significant relationship between rs10830963 *MTNR1B* polymorphism in obese and non-obese T2DM, where the CG genotype increases the risk of ⁴⁰ obese T2DM. Furthermore, there is no differences in the allele and genotype frequency of rs841853 *SLC2A1* polymorphism in the obese and non-³⁵ obese groups. There is also no significant difference in the biochemical parameters in obese and non-obese groups on the two ⁹ genetic variants studied. In addition, rs10830963 *MTNR1B* polymorphism is associated with the risk of obesity in Javanese T2DM patients but not for the rs841853 *SLC2A1* polymorphism.

Acknowledgements

The authors are grateful to the Central Laboratory of Diponegoro University Semarang (Cebior), the Laboratory of the Mathematics and Natural Sciences Faculty, CITO laboratory and the Public ¹⁰ Health Center that helped with the data collection. The authors are also grateful to the Directorate of Research and Community Services, Directorate General of Research and Development, Ministry of Research, Technology, and Higher Education, under the scheme of Research on Inter-University Collaboration for funding this study.

Authors Contribution

YT, AK, AY and ⁷ RS carried out the research; YT organized research data, performed data analysis, interpretation, wrote the manuscript and prepared final version for publication; AK performed data analysis and interpretation of data; AY and RS helped review the data analysis and reviewed the manuscript.

References

1. Kementrian Kesehatan RI. Profil Kesehatan Indonesia 2019. Jakarta: Kementrian Kesehatan RI; 2020.
2. Peschke E, Frese T, Chankiewicz E, Peschke D, Preiss U, Schneyer U, et al. Diabetic Goto Kakizaki rats as well as type 2 diabetic patients show a decreased diurnal serum melatonin level and an increased pancreatic melatonin-receptor status. *J Pineal Res.* 2006; 40: 135-43.
3. Reiter RJ, Tan DX, Rosales-Corral S, Galano A, Jou MJ, Acuna-Castroviejo D. Melatonin mitigates mitochondrial meltdown: Interactions with SIRT3. *Int J Mol Sci.* 2018; 19: 2-28.
4. Hevia D, González-Menéndez P, Quiros-González I, Miara A, Rodríguez-García A, Tan DX, et al. Melatonin uptake through glucose transporters: A new target for melatonin inhibition of cancer. *J Pineal Res.* 2015; 58: 234-50.
5. Prokopenko I, Langenberg C, Florez JC, Saxena R, Soranzo N, Thorleifsson G, et al. Variants in the melatonin receptor 1B gene (*MTNR1B*) influence fasting glucose levels. *Nat Genet.* 2010; 41: 77-81.
6. Takeuchi F, Katsuya T, Chakrawarthy S, Yamamoto K, Fujioka A, Serizawa M, et al. Common variants at the GCK, GCKR, G6PC2-ABCB11 and *MTNR1B* loci are associated with fasting glucose in two Asian populations. *Diabetologia.* 2010; 53: 299-308.
7. Kwak SH, Jang HC, Park KS. Finding genetic risk factors of gestational diabetes. *Genomics Inf.* 2012; 10: 239-43.
8. Sparso T, Bonnefond A, Andersson E, Bouatia-Naji N, Holmkvist J, Wegner L, et al. G-allele of intronic rs10830963 in *MTNR1B* confers increased risk of impaired fasting glycemia and type 2 diabetes through an impaired glucose-stimulated insulin release: Studies involving 19,605 Europeans. *Diabetes.* 2009; 58: 1450-6.

9. Song JY, Wang HJ, Ma J, Xu ZY, Hinney A, Hebebrand J, *et al.* Association of the rs10830963 polymorphism in *MTNR1B* with fasting glucose levels in Chinese children and adolescents. *Obes Facts.* 2011; 4: 197-203.
10. Du B, Liu S, Cui C, Wang S, Cui W. Association between glucose transporter 1 rs841853 polymorphism and type 2 diabetes mellitus risk may be population specific. *J Diabetes.* 2013; 5: 291-9.
11. Ma RCW, Chan JCN. Type 2 diabetes in East Asians: Similarities and differences with populations in Europe and the United States. *Ann NY Acad Sci.* 2013; 1281: 64-91.
12. George AM, Jacob AG, Fogelfeld L. Lean diabetes mellitus: An emerging entity in the era of obesity. *World J Diabetes.* 2015; 6: 613-20.
13. Kong X, Xing X, Hong J, Zhang X, Yang W. Genetic variants associated with lean and obese type 2 diabetes in a Han Chinese population A case-control study. *Medicine (Baltimore).* 2016; 95: e3841. doi: 10.1097/MD.0000000000003841.
14. Timpson NJ, Lindgren CM, Weedon MN, Randall J, Ouwehand WH, Strachan DP, *et al.* Adiposity-related heterogeneity in patterns of type 2 diabetes susceptibility observed in genome-wide association data. *Diabetes.* 2009; 58: 505-10.
15. Boutouris K, Poupouridou N, Katsarou M, S, Gourzi TF, Tentolouris K, *et al.* Frequency distribution of TCF7L2, *MTNR1B*, *CDKAL1*, *SLC30A8* and *FTO* gene polymorphisms predisposing to type II diabetes mellitus and/or obesity in a greek population and their impact in personalized medicine. *Arch Balk Med Union.* 2016; 51: 157-63.
16. Cipolla-Neto J, Amaral FG, Afeche SC, Tan DX, Reiter RJ. Melatonin, energy metabolism, and obesity: A review. *J Pineal Res.* 2014; 56: 371-81.
17. Szewczyk-Golec K, Rajewski P, Gackowski M, Mila-Kierzenkowska C, Wesolowski R, Sutkowy P, *et al.* Melatonin supplementation lowers oxidative stress and regulates adipokines in obese patients on a calorie-restricted diet. *Oxid Med Cell Longev.* 2017; 2017: 8494107. doi: 10.1155/2017/8494107.
18. De Luis DA, Izaola O, Primo D, Aller R. Association of the rs10830963 polymorphism in *MTNR1B* with fasting glucose, serum adipokine levels and components of metabolic syndrome in adult obese subjects. *Nutr Hosp.* 2019; 36: 60-5.
19. Pontiroli AE, Capra F, Veglia F, Ferrari M, Xiang KS, Bell GI, *et al.* Genetic contribution of polymorphism of the *GLUT1* and *GLUT4* genes to the susceptibility to type 2 (non-insulin-dependent) diabetes mellitus in different populations. *Acta Diabetol.* 1996; 33: 193-7.
20. Chobanian AV, Bakris GL, Black HR, Cushman WC, Green LA, Izzo JL Jr, *et al.* Seventh report of the Joint National Committee on prevention, detection, evaluation, and treatment of high blood pressure. *Hypertension.* 2003; 42: 1206-52.
21. Budiyati R, Lukito W, Wijaya A. Correlation between visfatin, insulin resistance (homeostasis model assesment of insulin resistance), inflammation (high sensitivity C-reactive protein) and HDL cholesterol concentration in individuals with visceral obesity. *Indones Biomed J.* 2010; 2: 61-5.
22. Lubis SM, Fattah M, Damanik HA, Batubara JRL. Association of fat mass and obesity-associated Gene (*FTO*) rs9939609 variant with early onset obesity among bataknese and Chinese children in Indonesia: A case-control study. *Indones Biomed J.* 2017; 9: 147-52.
23. Cauchi S, Choquet H, Gutiérrez-Aguilar R, Capel F, Grau K, Proença C, *et al.* Effects of *TCF7L2* polymorphisms on obesity in European populations. *Obesity.* 2008; 16: 476-82.
24. Voight BF, Scott LJ, Steinthorsdottir V, Andrew P, Aulchenko YS, Thorleifsson G, *et al.* Twelve type 2 diabetes susceptibility loci identified through large-scale association analysis. *Nat Genet.* 2010; 42: 579-89.
25. Saxena R, Gianniny L, Burt NP, Lyssenko V, Giuducci C, Sjögren M, *et al.* Common single nucleotide polymorphisms in *TCF7L2* are reproducibly associated with type 2 diabetes and reduce the insulin response to glucose in nondiabetic individuals. *Diabetes.* 2006; 55: 2890-5.
26. Tursinawati Y, Hakim RF, Rohmani A, Kartikadewi A, Sandra F. *CAPN10* SNP-19 is associated with susceptibility of type 2 diabetes mellitus: A Javanese case-control study. *Indones Biomed J.* 2020; 12: 109-14.
27. Cauchi S, Nead KT, Choquet H, Horber F, Potoczna N, Balkau B, *et al.* The genetic susceptibility to type 2 diabetes may be modulated by obesity status: Implications for association studies. *BMC Med Genet.* 2008; 9: 45. doi: 10.1186/1471-2350-9-45.
28. Linder K, Wagner R, Hatzigelaki E, Ketterer C, Heni M, Machicao F, *et al.* Allele summation of diabetes risk genes predicts impaired glucose tolerance in female and obese individuals. *PLoS One.* 2012; 7: e38224. doi: 10.1371/journal.pone.0038224.
29. Amini S, Javanmardi M, Mekarizadeh A, Maroofi F, Jalali C, Azadi NA, *et al.* Association of *HaeIII* single nucleotide polymorphisms in the *SLC2A1* gene with risk of diabetic nephropathy: evidence from Kurdish patients with type 2 diabetes mellitus. *Qjm.* 2016; 109: 399-404.
30. Freerman AJ, Johnson AR, Sacks GN, Milner JJ, Kirk EL, Troester MA, *et al.* Metabolic reprogramming of macrophages: Glucose transporter 1 (*GLUT1*)-mediated glucose metabolism drives a proinflammatory phenotype. *J Biol Chem.* 2014; 289: 7884-96.
31. Triana R, Meiliana A, Halimah E, Wijaya A. Relationship between circulating protein p53 and high sensitivity C-reactive protein in central obesity men with inflammation. *Indones Biomed J.* 2019; 11: 59-62.
32. Ling Y, Li X, Gu Q, Chen H, Lu D, Gao X. A common polymorphism rs3781637 in *MTNR1B* is associated with type 2 diabetes and lipids levels in Han Chinese individuals. *Cardiovasc Diabetol.* 2011; 10: 27.
33. Stefanidis I, Tziastoudi M, Tsironi EE, Dardiotis E, Tachmitzi SV, Fotiadou A, *et al.* The contribution of genetic variants of *SLC2A1* gene in T2DM and T2DM-nephropathy: association study and meta-analysis. *Ren Fail.* 2018; 40: 561-76.
34. Stefanidis I, Kytoudis K, Papanthasiou AA, Zaragotas D, Melistas L, Kitsios GD, *et al.* *XbaI* *GLUT1* gene polymorphism and the risk of type 2 diabetes with nephropathy. *Dis Markers.* 2009; 27: 29-35.
35. Rizvi S. Association of genetic variants with diabetic nephropathy. *World J Diabetes.* 2014; 5: 809-16.

Turnitin InBJ Polimorfisme MTNR1B

ORIGINALITY REPORT

22%

SIMILARITY INDEX

16%

INTERNET SOURCES

17%

PUBLICATIONS

6%

STUDENT PAPERS

PRIMARY SOURCES

1	www.atlantis-press.com Internet Source	2%
2	Yani Lina, Andi Wijaya. "Adipose-Derived Stem Cells for Future Regenerative System Medicine", The Indonesian Biomedical Journal, 2012 Publication	2%
3	Submitted to University of Leeds Student Paper	1%
4	www.hindawi.com Internet Source	1%
5	mejc.sums.ac.ir Internet Source	1%
6	scielo.isciii.es Internet Source	1%
7	A. Abdullah, F. A. Amin, J. Stoelwinder, S. K. Tanamas, R. Wolfe, J. Barendregt, A. Peeters. "Estimating the risk of cardiovascular disease using an obese-years metric", BMJ Open, 2014 Publication	1%

8	academic-accelerator.com Internet Source	1 %
9	Alaa Hashim Abd Ali, Tatyana Pavlovna Shkurat, Maria Aleksandrovna Amelina. "Investigation of the rs9939609 polymorphism of the FTO gene in obese children and adolescents living", AIP Publishing, 2022 Publication	1 %
10	www.ncbi.nlm.nih.gov Internet Source	1 %
11	"Diabetes: from Research to Clinical Practice", Springer Science and Business Media LLC, 2021 Publication	1 %
12	file.scirp.org Internet Source	<1 %
13	"Minutes of the 44th Genral Assembly of the European Association for the Study of Diabetes", Diabetologia, 2009 Publication	<1 %
14	Jamal S.M. Sabir, Abdelfatteh El Omri, Noor A. Shaik, Babajan Banaganapalli et al. "The genetic association study of TP53 polymorphisms in Saudi obese patients", Saudi Journal of Biological Sciences, 2019 Publication	<1 %

media.neliti.com

15

Internet Source

<1 %

16

www.besjournal.com

Internet Source

<1 %

17

www.ijmtss.com

Internet Source

<1 %

18

"Posters", Journal of Diabetes, 04/2009

Publication

<1 %

19

www.thaiscience.info

Internet Source

<1 %

20

Stalis Norma Ethica, Sri Darmawati, Sri Sinto Dewi, Nurrahman Nurrahman, Ayu Rahmawati Sulistyaningtyas. "Streptolysin Encoding Genes sagC and sagD as Biomarkers of Fish Pathogen Streptococcus iniae: An In Silico Study", Squalen Bulletin of Marine and Fisheries Postharvest and Biotechnology, 2020

Publication

<1 %

21

Budi Santosa, Henna Ria Sunoko, Andri Sukeksi, Siti Thomas Zulaikhah. "IR Bagendit Paddy Leaves Extract Improves Liver Cell Morphology and Reduces The Activity of Transaminase Enzymes After Lead Exposure in Rat", The Indonesian Biomedical Journal, 2020

Publication

<1 %

- | | | |
|----|--|------|
| 22 | Jie Wu, Yiqing Zhu, Yunfei Jiang, Yaoping Cao. "Effects of metformin and ethinyl estradiol-cyproterone acetate on clinical, endocrine and metabolic factors in women with polycystic ovary syndrome", <i>Gynecological Endocrinology</i> , 2009
Publication | <1 % |
| 23 | journals.plos.org
Internet Source | <1 % |
| 24 | Submitted to Higher Education Commission Pakistan
Student Paper | <1 % |
| 25 | Submitted to Oxford Brookes University
Student Paper | <1 % |
| 26 | www.neliti.com
Internet Source | <1 % |
| 27 | "eHEART: www.heartjnl.com ", <i>Heart</i> , 10/1/2003
Publication | <1 % |
| 28 | Du, Bing, Shengmao Liu, Chengji Cui, Songyan Wang, and Wenpeng Cui. "Association between glucose transporter 1 rs841853 polymorphism and type 2 diabetes mellitus risk may be population specific (葡萄糖转移酶1的rs841853基因多态性与2型糖尿发生风险的关系可能具有人群特异性) : Is rs841853 a risk factor for T2D?", <i>Journal of Diabetes</i> , 2013. | <1 % |

29

Parra, Eliane, Natália Panzoldo, Denise Kaplan, Helena Coutinho de Oliveira, José dos Santos, Luiz Sérgio de Carvalho, Andrei Sposito, Magnus Gidlund, Ruy Nakamura, Vanessa de Souza Zago, Edna Nakandakare, Eder Carlos Quintão, and Eliana de Faria. "The I405V and Taq1B polymorphisms of the CETP gene differentially affect sub-clinical carotid atherosclerosis", *Lipids in Health and Disease*, 2012.

<1 %

Publication

30

Qi Wu, Yanmin Chen, Menglin Zhou, Mengting Liu, Lixia Zhang, Zhaoxia Liang, Danqing Chen. "An early prediction model for gestational diabetes mellitus based on genetic variants and clinical characteristics in China", *Diabetology & Metabolic Syndrome*, 2022

<1 %

Publication

31

Xinxia Zhang, Lanlan Xie, Min Zhong, Botong Yang, Qiyue Yang, Hui Yang, Chunguang Xie. "The association between melatonin receptor 1B gene polymorphisms and type 2 diabetes mellitus (T2DM) in Chinese populations: a meta-analysis", *Annals of Palliative Medicine*, 2020

<1 %

Publication

32

cms.jtgga.org
Internet Source

<1 %

33

pure.rug.nl
Internet Source

<1 %

34

www.bioseek.eu
Internet Source

<1 %

35

Dalia Essamy El Nashar, Kholoud Mohammad Alananbeh, Noora Al Hassan. "Genetic, dietary, and non-dietary risk factors of obesity among preparatory-year female students at Taibah University, Saudi Arabia", Journal of Taibah University for Science, 2018
Publication

<1 %

36

Parvaneh Keshavarz, Razie Habibipour, Malaek Ghasemi, Ehsan Kazemnezhad, Maryam Alizadeh, Mohammad Hasan Hedayati Omami. "Lack of genetic susceptibility of KCNJ11 E23K polymorphism with risk of type 2 diabetes in an Iranian population", Endocrine Research, 2014
Publication

<1 %

37

Sumeet Gupta, Vikas Jhawar, Bimal Kumar Agarwal, Partha Roy, Vipin Saini. "Alpha Adducin (ADD1) Gene Polymorphism and New Onset of Diabetes Under the Influence of Selective Antihypertensive Therapy in

<1 %

Essential Hypertension", Current Hypertension Reviews, 2019

Publication

38

Wang, X.. "APM1 gene variants -11377C/G and 4545G/C are associated respectively with obesity and with non-obesity in Chinese type 2 diabetes", Diabetes Research and Clinical Practice, 200906

Publication

<1 %

39

Y. Ishii, H. Hasegawa, H. Nishibori, M. Watanabe, M. Kitajima. "Impact of visceral obesity on surgical outcome after laparoscopic surgery for rectal cancer", British Journal of Surgery, 2005

Publication

<1 %

40

bmcrenotes.biomedcentral.com

Internet Source

<1 %

41

epublications.uaf.fi

Internet Source

<1 %

42

gbmrjournal.com

Internet Source

<1 %

43

jpbsonline.org

Internet Source

<1 %

44

jurnal.umj.ac.id

Internet Source

<1 %

45

onlinelibrary.wiley.com

Internet Source

<1 %

46

www.karger.com

Internet Source

<1 %

47

www.science.gov

Internet Source

<1 %

48

Christiane B. Sabourin, Marjorie Funk.
"Readmission of patients after coronary
artery bypass graft surgery", Heart & Lung:
The Journal of Acute and Critical Care, 1999

Publication

<1 %

49

Cipolla-Neto, J., F. G. Amaral, S. C. Afeche, D.
X. Tan, and R. J. Reiter. "Melatonin, energy
metabolism, and obesity: a review", Journal of
Pineal Research, 2014.

Publication

<1 %

50

Margaret R. Spitz, Michelle A. Detry, Patricia
Pillow, Yaohua Hu, Christopher I. Amos, Waun
Ki Hong, Xifeng Wu. "Variant alleles of the D2
dopamine receptor gene and obesity",
Nutrition Research, 2000

Publication

<1 %

51

O Mir. "Cyclooxygenase 2 selective inhibitor
induced bowel stricture: a case report", Gut,
1/1/2004

Publication

<1 %

52	Siska Mayasari Lubis, Miswar Fattah, Harun Alrasyid Damanik, Jose Rizal Latief Batubara. "Association of Fat Mass and Obesity-associated Gene (FTO) rs9939609 Variant with Early Onset Obesity among Bataknese and Chinese Children in Indonesia: A Case-control Study", The Indonesian Biomedical Journal, 2017 Publication	<1 %
53	bsj.uobaghdad.edu.iq Internet Source	<1 %
54	ejournal2.undip.ac.id Internet Source	<1 %
55	eprints.qut.edu.au Internet Source	<1 %
56	inabj.org Internet Source	<1 %
57	pr.hec.gov.pk Internet Source	<1 %
58	www.mdpi.com Internet Source	<1 %
59	www.scielo.br Internet Source	<1 %
60	Sara Pourdasthi, Nassim Faridi, Forouzandeh Monem-Homaie, S. Hamid Yaghooti,	<1 %

Ahmadreza Soroush, S. Zahra Bathaie.
"Subcutaneous adipocyte size but not adiposity was associated with inflammation, ER stress, and insulin resistance markers",
Research Square Platform LLC, 2022

Publication

61

Yan-yan Li, Hui Wang, Yang-yang Zhang.
"Melatonin receptor 1B gene rs10830963 C/G polymorphism associated with type 2 diabetes mellitus: An updated meta-analysis of 13,752 participants", Heliyon, 2022

Publication

62

Denis Foschini, Ronaldo V. T. dos Santos, Wagner L. Prado, Aline de Piano et al.
"Plaqueta e leptina em adolescentes com obesidade", Jornal de Pediatria, 2008

Publication

63

Jie-Yun Song, Hai-Jun Wang, Jun Ma, Zhi-Yuan Xu, Anke Hinney, Johannes Hebebrand, Yan Wang.
"Association of the rs10830963 Polymorphism in *MTNR1B* with Fasting Glucose Levels in Chinese Children and Adolescents", Obesity Facts, 2011

Publication

64

Xi Yang, YueFei Jiang, XinLin Hao, XiaoLing Cheng, Qin Wan.
"Correlation Between Cardiometabolic Index and Microalbuminuria

<1 %

<1 %

<1 %

<1 %

in Type 2 Diabetic Patients", Research Square Platform LLC, 2021

Publication

Exclude quotes Off

Exclude matches Off

Exclude bibliography On