

# Turnitin Mirna29 Scopus

*by Dr, Yanuarita*

---

**Submission date:** 17-Jan-2023 01:16PM (UTC+0700)

**Submission ID:** 1993985232

**File name:** Mirna29\_Scopus.pdf (2.55M)

**Word count:** 4680

**Character count:** 25484

## Overexpression of miR-29b in Plasma of Javanese Type 2 Diabetes Mellitus Patients in Semarang, Indonesia

### ABSTRACT

**Objective:** In type 2 diabetes, the miR-29b, also known as redoximirs, has been shown to regulate oxidative pathways. The aim of this study was to observe the circulation of miR-29b in plasma of Javanese type 2 diabetes mellitus patients at the Primary Health Care Center in Semarang, Indonesia, and its correlation with fasting blood glucose, cholesterol, and triglyceride.

**Methods:** The expression of miR-29b in type 2 diabetes mellitus groups and control groups was measured by reverse transcriptase polymerase chain reaction (RT-PCR). The  $2^{-\Delta\Delta Ct}$  method was used to determine the relative amount of miR-29b. The miR-29b expression was compared with an independent t-test. Furthermore, the correlation between miR-29b expression was analyzed by the Pearson correlation test using SPSS 17.0. Fold change was positive at 14.12 (i.e., upregulation).

**Results:** There was a significant negative correlation between age and miR-29b expression ( $P=0.000$ ,  $r=-0.627$ ). Moreover, fasting blood glucose and miR-29b also had a negative correlation ( $P=0.000$ ,  $r=-0.504$ ).

**Conclusion:** miR-29b overexpression occurred in plasma of Javanese type 2 diabetes mellitus patients at the Primary Health Care Center in Semarang, Indonesia, i.e., 14.12-fold overexpression. The level of miR-29b expression was negatively correlated with age and fasting blood glucose.

**Keywords:** Diabetes mellitus, Indonesia, micro-RNAs, miR-29b, plasma

### Introduction

In Indonesia, noncommunicable and communicable diseases (infectious disease) have the same high prevalence. More precisely, cases of noncommunicable diseases such as stroke, hypertension, diabetes mellitus (DM), tumor, and heart disease have continually increased. A total of 13% population in Indonesia has DM. Approximately 80%-90% of DM cases are type 2 diabetes mellitus (T2DM).<sup>1</sup> There were 10.7 million T2DM patients in Indonesia, according to The International Diabetes Federation (IDF) in 2019.<sup>2</sup> The Indonesian Ministry of Health has reported an increased number of T2DM cases from 1.1% in 2007 to 2.1% in 2013.<sup>3</sup> Javanese is the major population in Central Java province. DM incidence is the second highest after hypertension.<sup>4</sup> Javanese is the biggest ethnic group in Indonesia and has a cultural preference for consuming sweet foods that trigger a high risk of DM.

Diabetes mellitus is a disease characterized by a variety of symptoms such as hyperglycemia. Type 2 diabetes mellitus is a metabolic and endocrine disorder caused by insulin resistance and pancreatic  $\beta$ -cell dysfunction. Insulin resistance occurs due to inadequate insulin receptors in cells. Thus, cells are unable to use the glucose, and blood glucose levels remain high. Insulin resistance mainly occurs in insulin-sensitive tissues, such as muscle, adipose, liver, gastrointestinal tract, central nervous system, and pancreatic cells.<sup>5</sup> Insulin resistance pathophysiology involves several factors, including mitochondrial dysfunction, inflammation, obesity, and oxidative stress.<sup>6</sup>

The hyperglycemia condition in T2DM causes protein glycation, glucose autooxidation, lipid peroxidase, and glucose metabolic imbalance. The polyol pathway is involved in glucose metabolic imbalance. This mechanism causes the formation of free radicals that leads to the imbalance ratio of free radicals and antioxidants in the body. Moreover, it induces oxidative stress. Oxidative stress causes the damage of DNA and proteins in various tissues and is responsible for the development of T2DM and its complications.<sup>7</sup> micro-RNA has the capability to increase reactive oxygen species or decrease the antioxidant expression in T2DM.

Yanuarita Tursinawati<sup>1</sup>

Arum Kartikadewi<sup>1</sup>

Ari Yuniastuti<sup>1</sup>

Ratna Susanti<sup>2</sup>

<sup>1</sup>Department of Biomedics, Universitas Muhammadiyah Semarang Faculty of Medicine, Semarang, Indonesia

<sup>2</sup>State University of Semarang Faculty of Mathematics and Natural Sciences, Semarang, Indonesia

Corresponding author:

Yanuarita Tursinawati

✉ yanuarita\_tursina@unimus.ac.id

Received: April 10, 2022

Accepted: October 3, 2022

Publication Date: December 1, 2022

Cite this article as: Tursinawati Y, Kartikadewi A, Yuniastuti A, Susanti R. Overexpression of miR-29b in plasma of Javanese type 2 diabetes mellitus patients in Semarang, Indonesia. *Turk J Endocrinol Metab.* 2022;26(4):208-212.

DOI: 10.5152/tjem.2022.22016

Micro-RNA is a noncoding RNA consisting of 18–25 nucleotides that bind to the 3′-untranslated region of the target gene. Small micro-RNA molecules are actively or passively released into the circulation. It can be used as a biomarker for the onset and progression of some diseases such as DM. In diabetes, micro-RNA has a role in oxidative stress mechanism, inflammation, or insulin signaling pathway that responds to the development of insulin resistance. A previous study suggests the associations of miR-29 with insulin resistance in Goto Kakizaki mice. The expression of miR-29a and miR-29b significantly increases in 3T3-L1 cells after incubating with high glucose and various doses of insulin.<sup>8</sup> miR-29a and miR-29b are associated with caveolin 2, insulin-induced gene 1, and phosphatidylinositol 3-kinase regulatory subunit-α in the insulin signaling pathway.<sup>9,10</sup>

miR-29b, also known as redoximiRs, is a dominant micro-RNA in regulating gene expression and oxidative pathways associated with the incidence of DM.<sup>11–13</sup> In this study, we investigate the circulation of miR-29b in plasma of Javanese T2DM patients at the Primary Health Care Center in Semarang, Indonesia, and its correlation with fasting blood glucose (FBG), cholesterol, and triglyceride.

## Materials and Methods

### Research Design

The case–control study was performed by a consecutive sampling of T2DM patients in the Chronic Disease Management Program (PROLANS) program at the Primary Health Care Center in Semarang, Indonesia. The inclusion criteria were Javanese T2DM patients for the case group and non-T2DM with age 30–70 years for the control group. The exclusion criteria were having a history of cardiovascular disorders such as stroke, heart failure, and acute myocardial infarction. A total of 30 patients for the case group and 17 patients for the control group were used in this study. All respondents had signed the informed consent as a sign of willingness to participate in this study. This work was approved by the Ethics Commission of the Faculty of Medicine, Universitas Muhammadiyah Semarang, with Number 040/EC/FK/2019.

### Sample Analysis

Blood samples were obtained from the vein to examine the fasting blood sugar level and lipid profile (triglycerides and cholesterol). The plasma was used for micro-RNA examination. The RNA extraction was conducted using miRNeasy Serum/Plasma Advanced Kit (Qiagen, Hilden, Germany) (cat. no. 217204). RNA was transcribed into cDNA using the miRCURY LNA RT Kit (cat. no. 339340, Qiagen). Quantitative real-time PCR was performed using ExiLent SYBR Green master mix reagent (Qiagen, cat. no. 339345).

After RNA was extracted, total of 2 μL RNA was transcribed into cDNA mixing with 4.5 μL nuclease-free water, 2 μL RT Rx buffer, 1 μL RT enzyme mix, and 0.5 μL UniSp6 RNA spike. The solution was mixed with free water, 2 μL RT Rx buffer, 1 μL RT enzyme mix, and

0.5 μL UniSp6 RNA spike. For PCR amplification, the solution was conducted in a ProFlex thermal cycler at 42°C for 60 minutes, 95°C for 5 minutes, and immediately cool to 4°C. miR-29b expression was analyzed by reverse transcriptase polymerase chain reaction (RT-PCR), by mixing 3 μL cDNA, 1 μL free water, 5 μL ExiLent SYBR Green master mix, 1 μL PCR primary mix, and 0.05 μL Rox reference dye. The RT-PCR Program was performed on 7500 fast machines, with initial denaturation at 95°C for 2 minutes, followed by amplification for 40 cycles at 95°C for 10 seconds, 60°C for 60 seconds (melting curve 60°C–95°C). Hsa-mir-29b-3p and U6 snRNA were used as primers for RT-PCR.

micro-RNA amplification values were analyzed using the 7500 Fast RT-PCR system v2.3 software to obtain the value of Ct, the melt curve, and the melt peak curve. An increase or decrease in comparison in the target expression was analyzed using the Livak method is the  $2^{-\Delta\Delta C_t}$  method used to calculate the relative quantification of microRNA expression in plasma of T2DM patients compared to control with U6 snRNA as a reference gene.

### Statistical Analysis

miR-29b expression was presented as mean ± SD and compared to the T2DM group and control group using an independent t-test. The correlations between miR-29b expression with fasting blood sugar, cholesterol, and triglycerides were measured using a Pearson correlation test with a significance of  $P < .05$  and CI=95%. Multivariate analysis was conducted using a multiple regression test to determine the most important factor on miR-29b expression. All data were analyzed using the Statistical Package for Social Sciences, version 17.0 software (SPSS Inc.; Chicago, IL, USA).

## Results

A total of 30 T2DM patients and 17 patients for control was used in this study. The highest levels of FBG, cholesterol, and triglycerides were  $199.70 \pm 14.88$  mg/dL,  $200.27 \pm 7.69$  mg/dL, and  $201.97 \pm 22.42$  mg/dL in the T2DM group. We found that the highest body mass index (BMI) was  $24.55 \pm 0.92$  in the control group. There were significant differences between the T2DM and control groups with  $P < .005$ , that is, for the age parameters, FBG, and triglycerides. There was no significant difference between sex, BMI, and cholesterol (Table 1).

The miR-29b relative expression is presented in Table 2. We found that the fold change ( $2^{-\Delta\Delta C_t}$ ) value was positive with a value of 14.12

**Table 1. Sample Characteristics and Biochemical Data**

Variable	T2DM Group (n=30)	Control (n=17)	P
	Mean ± SD	Mean ± SD	
Age (years)	57.60 ± 1.39	45.35 ± 1.59	.000**
Gender			
Female	24	12	.493*
Male	6	5	
Body mass index	24.33 ± 0.57	24.55 ± 0.92	.929**
Fasting blood glucose (mg/dL)	199.70 ± 14.88	96.41 ± 2.55	.000**
Cholesterol (mg/dL)	200.27 ± 7.69	182.53 ± 11.06	.084**
Triglyceride (mg/dL)	201.97 ± 22.42	141.71 ± 18.87	.043**

T2DM, type 2 diabetes mellitus; SD, standard deviation.

\*Chi-square.

\*\*Mann–Whitney U-test.

## MAIN POINTS

- miR-29b overexpression occurred in the plasma of Javanese type 2 diabetes mellitus patients.
- miR-29b expressions were negatively correlated with age.
- miR-29b expressions were negatively correlated with fasting blood glucose.

**Table 2. Expression of miR-29b in T2DM Patients and Control Group**

Variable	T2DM (n = 30)	Control (n = 17)	P	$\Delta\Delta Ct$ miR-29b	Fold Change $2^{-\Delta\Delta Ct}$	Regulation
	Mean $\pm$ SD	Mean $\pm$ SD				
miR-29b	-1.38 $\pm$ 1.94	2.44 $\pm$ 2.28	.000	-3.82	14.12	Upregulation

T2DM, type 2 diabetes mellitus; SD, standard deviation.

according to the LIVAK method. It revealed that expression of plasma miR-29b in T2DM patients was increased compared to the control group. miR-29b levels between the T2DM and control groups were significantly different with  $P = .000$  (Figure 1).

Furthermore, there was a significant negative correlation between age and miR-29b variables ( $P = .000$ ,  $r = -0.627$ ) and between FBG and miR-29b variables ( $P = .000$ ,  $r = -0.504$ ). There were no significant correlations among BMI, cholesterol, and triglycerides (Table 3).

Based on the multivariate analysis results (Table 4), it was found that the variables with  $P < .005$  were age and FBG, with  $P = .000$  and  $P = .021$ , respectively. These results showed that age was the most important risk factor of the miR-29b levels.

#### Discussion

In the present study, we investigated the relationship between the miR-29b and DM incidence, especially in Javanese ethnicity. Based on the Household Life Aspect Survey data, the highest incidence of diabetes in Indonesia was Javanese and Madura ethnicity, so this study focuses on the incidence of diabetes in this ethnic group.<sup>14</sup>

micro-RNA has been shown to affect the oxidative stress in T2DM through the increase of reactive oxygen species or a decrease in antioxidant expression. Micro-RNA impact oxidative stress in diabetes pathogenesis through several mechanisms, including Nrf2 is (nuclear factor erythroid 2-related factor 2)–Keap1 (Kelch ECH associating protein 1) pathway, Sirtuin-1 (SIRT1), PGC is peroxisome proliferator-activated receptor-gamma coactivator-1 $\alpha$ , superoxide dismutase, catalase, and NADPH is nicotinamide adenine dinucleotide phosphate hydrogen oxidase 4 (NOX4).<sup>15-18</sup> Nuclear factor-erythroid-2-related factor 2 (Nrf-2) regulates the gene expression antioxidant Nuclear factor-erythroid-2 (NF-E2)-related factor 2 (Nrf-2) regulates the expression of antioxidant and cytoprotective

genes against oxidative stresses.<sup>19</sup> Nrf-2 interacts with Kelch-like golgi apparatus membrane protein-like protein ECHIDNA (ECH)-associated protein (KEAP1) that triggers the proteasomal degradation and increases along with the reduced cellular antioxidant responses. miR-29b is a KEAP1 expression regulator. Reduced expression of miR-29b occurs in diabetes and increases NRF-2 expression and oxidative stress.<sup>20,21</sup> These results are contradictory to our findings.

We demonstrated an increased expression of miR-29b in plasma of Javanese T2DM patients. These results were in line with a study conducted by Hung<sup>22</sup> using T2DM mice liver that observed miR-107, miR-148b, miR-222, miR-802, miR-34a, and miR-29a/b/c. Our study showed that miR-29b expression in T2DM was upregulated to 1.5-fold from normal conditions. In contrast, the study conducted by Zampetaki et al<sup>23</sup> showed a decrease of miR-29b, miR-126, and miR-223 expression in T2DM plasma.

miR-29b was overexpressed in this study. This is similar to a research by He et al that examined miR-29a/b/c in the muscle, fat, and liver of diabetic mice. An elevated level of miR-29b caused insulin resistance and decreased insulin-induced glucose uptake in adipocyte 3T3-L1 cells. Of the 3 miR-29 families, only miR-29a and miR-29b were modulated by hyperlipidemia and hyperglycemia in DM.<sup>8</sup> Another study demonstrated that miR-29a and miR-29c had a role in T2DM patients to regulate glucose of skeletal cells. An increase in expression only occurs in miR-29a and miR-29c, while miR-29b has no change in expression. This is contradictory to our results. An increase in miR-29a and miR-29c expression was the major cause of glucose reduction by 20% not in miR-29b.<sup>24</sup>

Furthermore, miR-29b expression decreases in diabetes and increases the expression of proteasome activator and its activity in




Figure 1.  $\Delta Ct$  comparison of miR-29 expression in plasma of T2DM group (n=30) and control group (n=17) with  $P = .000$ . T2DM, type 2 diabetes mellitus.

**Table 3. Correlation Between Parameters and miR-29b Expression**

Variable	miR-29b P-value	r
Age	.000	-0.627
Body mass index	.749	0.048
Fasting blood glucose	.000	-0.504
Cholesterol	.056	-0.280
Triglyceride	.108	-0.238

**Table 4. Multivariate Analysis to Determine Risk Factors of miR-29b Expression**

Variable	B	SE	P
Age	-0.143	0.037	.000
Body mass index	-0.041	0.094	.662
Fasting blood glucose	-0.010	0.004	.021
Cholesterol	-0.007	0.007	.331
Triglyceride	0.000	0.003	.744

SE, standard error.

endothelial cells. Proteasome has an important role in the degradation of nuclear factor kappa B (NF $\kappa$ B  $\alpha$ ) inhibitor. The elevated activity of proteasome will increase the activity of transcript factors NF $\kappa$ B and NOX. They trigger oxidative stress.<sup>25</sup>

Another study using Cmah-null mice showed that miR-29b-3p was significantly upregulated in the pancreas and liver through PI3K/AKT in insulin signaling pathway.<sup>26</sup> Overexpression of miR-29b inhibits the insulin-induced phosphorylation of AKT in the hepatocyte. It reveals that micro-RNA acts as a negative regulator of insulin signaling.<sup>27</sup> 5-Aminoimidazole-4-carboxamide 1- $\beta$ -D-ribofuranoside is also involved in insulin signaling that regulates hepatic glucose and lipid metabolism. This pathway involves miR-29b downregulated against the p85 target genes that inhibit AKT insulin-induced phosphorylation in wild-type C57BL/6 mouse hepatocytes.<sup>28</sup> This is in line with our study that the overexpression of miR-29b may inhibit the insulin signaling pathway in T2DM patients.

The role of miR-29b in insulin resistance involves a facilitated glucose transporter member 4 (GLUT4) along with hexokinase-2 (HK2). In skeletal muscle and white adipose tissue, glucose uptake is facilitated by GLUT4 through Akt activation.<sup>29,30</sup> If glucose gets into the cells, it will be phosphorylated by HK2 into glucose-6-phosphate and affect further metabolism. A decreased expression of GLUT4 in the muscle cells was commonly found in diabetic mice and humans.<sup>31,32</sup> Moreover, there was a decreased expression of SLC2A3, GLUT4 and Hk2/HK2 and increased expression of miR-29b-3p.<sup>33</sup> An increased expression of miR-29b-3p and miR-29c-3p was found in streptozotocin (STZ), a compound that has a toxicity toward pancreatic  $\beta$  cells-diabetic rats. GLUT4 and HK2 were negatively correlated with miR-29b-3p.

The miR-29 roles in DM are also associated with other proteins, including Secreted Protein Acidic Rich in Cysteine (SPARC). Secreted Protein Acidic Rich in Cysteine is a protein produced by adipocytes that regulate adipogenesis, adipocyte differentiation, and hyperplasia.<sup>34</sup> Obesity and insulin resistance have an impact on SPARC expression in T2DM.<sup>35</sup> miR-29s also regulates SPARC by binding with 3'-UTR. Overexpression of SPARC increases insulin secretion in beta cells and GLUT4 production in the 3T3-L1 adipocyte.<sup>36</sup> Song et al<sup>37</sup> suggest that the overexpression of miR-29 (consisted of miR-29 a,b,c) downregulates SPARC that leads to GLUT4 downregulation in the 3T3-L1 adipocyte. It induces insulin resistance due to the decrease in glucose uptake. This could be one of the theories that are in line with our findings. SPARC is involved in miR-29b overexpression in T2DM.

This study also analyzed the correlation of miR-29 levels with different characteristics and clinical chemical parameters. The results showed that there was a negative correlation between age and miR-29b (the most important factor) and between fasting blood glucose and miR-29b. The miR-29b level is correlated with fasting blood glucose. The higher the expression of miR-29b, the fasting blood glucose level will decrease. This is in line with Liang's study that demonstrates that overexpression of miR-29a-c in rat liver decreases fasting blood glucose through negative regulation of hepatic gluconeogenesis. The decrease is mediated by the downregulation of PGC-1 $\alpha$  and G6Pase proteins that are targets of miR-29a-c. The decrease in PGC-1 $\alpha$  and G6Pase proteins causes a reduction in liver glucose production.<sup>38</sup> However, Zampetaki et al<sup>23</sup> showed that many micro-RNAs were inversely correlated with FBG, except miR-29b and miR-320.

miR-29b levels are considered to decrease with increase in age. Circulating miRNA is simultaneously detected in the blood as a biomarker, and its level is influenced by several factors, including age. Moreover, some miRNAs such as miR-24 and miR-130b-5p are negatively correlated with age.<sup>39</sup> However, levels of plasma miR-29b related to age have never been reported, although increasing levels of miR-29b with age in mice and human brains have been discovered.<sup>40</sup> Therefore, this study can be used as a reference that miR-29b levels are correlated with the age of T2DM patients.

This study did not simultaneously measure miR-29b with the target protein such as NRF2-KEAP1, Sirtuin 1 (SIRT1), Toll-like Receptor 7 (TLR7), GLUT, PI3K/AKT, SPARC, PGC-1 $\alpha$ , and G6Pase so that further research may consider specific protein targets.

## Conclusions

miR-29b overexpression occurred in plasma of Javanese T2DM patients at the Primary Health Care Center in Semarang, Indonesia, that is, 14.12-fold overexpression. The level of miR-29b expression was negatively correlated with age and FBG.

## Funding

Directorate of Research and Community Services, Directorate General of Research and Development, Ministry of Research, Technology, and Higher Education, the Republic of Indonesia, funded this study.

## Source of Finance

During this study, no financial or spiritual support was received neither from any pharmaceutical company that has a direct connection with the research subject nor from a company that provides or produces medical instruments and materials which may negatively affect the evaluation process of this study.

**Ethics Committee Approval:** This research has received ethical approval from the Ethics Commission of the Faculty of Medicine, Universitas Muhammadiyah Semarang, with Number 040/EC/FK/2019.

**Informed Consent:** Written informed consent was obtained from the patients who participated in this study.

**Peer-review:** Externally peer-reviewed.

**Author Contributions:** Data Collection and/or Processing – Y.T.; Analysis and/or Interpretation – Y.T., A.K., A.Y., R.S.; Literature Search – ; Writing Manuscript – Y.T.; Critical Review – A.Y., R.S.

**Acknowledgments:** The authors would like to the head of the Semarang City Health Center for allowing researchers to conduct studies in their area.

**Declaration of Interests:** The authors declare that they have no competing interest.

**Funding:** This study is funded by the Directorate of Research and Community Services, Directorate General of Research and Development, Ministry of Research, Technology, and Higher Education, the Republic of Indonesia, under the research scheme of Research Collaboration between Universities, 2019 No. 7/E/KPT/2019.

## References

1. Idris H, Hasyim H, Utama F. Analysis of diabetes mellitus determinants in Indonesia. A study from the Indonesian basic health research 2013. *Acta Med Indones.* 2017;49(4):291-298.
2. International Diabetes Federation. *IDF Diabetes Atlas.* 8th ed. Brussels: International Diabetes Federation; 2017.

3. Data and Information Center IM of H. *Situasi dan Analisis Diabetes*. Jakarta: Data and Information Center IM of H; 2014.
4. Dinas Kesehatan Provinsi Jawa Tengah. *Profil Kesehatan Profinsi Jawa Tengah Tahun 2017*. Indonesia: Central Java Government; 2017.
5. Nigi L, Grieco GE, Ventriglia G, et al. MicroRNAs as regulators of insulin signaling. Research updates and potential therapeutic perspectives in type 2 diabetes. *Int J Mol Sci*. 2018;19(12):1-24. [\[CrossRef\]](#)
6. Hirsch GE, Heck TG. Inflammation, oxidative stress, and altered heat shock response in type 2 diabetes: the basis for new pharmacological and non-pharmacological interventions. *Arch Physiol Biochem*. 2019;20:1-15. [\[CrossRef\]](#)
7. Qadir MMF, Klein D, Álvarez-Cubela S, Domínguez-Bendala J, Pastori RL. The role of microRNAs in diabetes-related oxidative stress. *Int J Mol Sci*. 2019;20(21):1-25. [\[CrossRef\]](#)
8. He A, Zhu L, Gupta N, Chang Y, Fang F. Overexpression of micro ribonucleic acid 29, highly up-regulated in diabetic rats, leads to insulin resistance in 3T3-L1 adipocytes. *Mol Endocrinol*. 2007;21(11):2785-2794. [\[CrossRef\]](#)
9. Deng J, Guo M. MicroRNAs and type 2 diabetes. *ExRNA*. 2019;1:1. [\[CrossRef\]](#)
10. Pandey AK, Verma G, Vig S, Srivastava S, Srivastava AK, Datta M. MiR-29a levels are elevated in the db/db mice liver and its overexpression leads to attenuation of insulin action on PEPCCK gene expression in HepG2 cells. *Mol Cell Endocrinol*. 2011;332(1-2):125-133. [\[CrossRef\]](#)
11. Roggli E, Gattesco S, Caille D, et al. Changes in microRNA expression contribute to pancreatic  $\beta$ -cell dysfunction in prediabetic NOD mice. *Diabetes*. 2012;61(7):1742-1751. [\[CrossRef\]](#)
12. Pullen TJ, da Silva Xavier GA, Kelsey G, Rutter GA. miR-29a and miR-29b contribute to pancreatic cell specific silencing of monocarboxylate transporter 1 (Mct1). *Mol Cell Biol*. 2011;31(15):3182-3194. [\[CrossRef\]](#)
13. Cheng X, Ku CH, Siow RCM. Regulation of the Nrf2 antioxidant pathway by microRNAs. New players in micromanaging redox homeostasis. *Free Radic Biol Med*. 2013;64:4-11. [\[CrossRef\]](#)
14. Garnita D, Risiko F. *Diabetes Mellitus di Indonesia (Analisis Data SAKERTI 2007)*. Jakarta: University of Indonesia; 2012.
15. Julian FGK. *Diabetes: Oxidative Stress and Dietary Antioxidants*. 2nd ed. London: Academic Press; 2020.
16. Mortuza R, Feng B, Chakrabarti S. MiR-195 regulates SIRT1-mediated changes in diabetic retinopathy. *Diabetologia*. 2014;57(5):1037-1046. [\[CrossRef\]](#)
17. Zhao Y, Dong D, Reece EA, Wang AR, Yang P. Oxidative stress-induced miR-27a targets the redox gene Nrf2 in diabetic embryopathy. *Am J Obstet Gynecol*. 2018;218(1):136.e1-136.e10. [\[CrossRef\]](#)
18. Fu Y, Zhang Y, Wang Z, et al. Regulation of NADPH oxidase activity is associated with miRNA-25-mediated NOX4 expression in experimental diabetic nephropathy. *Am J Nephrol*. 2010;32(6):581-589. [\[CrossRef\]](#)
19. Reichard JF, Motz GT, Puga A. Heme oxygenase-1 induction by NRF2 requires inactivation of the transcriptional repressor BACH1. *Nucleic Acids Res*. 2007;35(21):7074-7086. [\[CrossRef\]](#)
20. Wei J, Zhang Y, Luo Y, et al. Aldose reductase regulates miR-200a-3p/141-3p to coordinate Keap1-Nrf2, Tgf $\beta$ 1/2, and Zeb1/2 signaling in renal mesangial cells and the renal cortex of diabetic mice. *Free Radic Biol Med*. 2014;67:91-102. [\[CrossRef\]](#)
21. Zhou L, Xu DY, Sha WG, et al. High glucose induces renal tubular epithelial injury via Sirt1/NF-kappaB/microR-29/Keap1 signal pathway. *J Transl Med*. 2015;13:352. [\[CrossRef\]](#)
22. Hung YH, Kanke M, Kurtz CL, et al. Acute suppression of insulin resistance-associated hepatic miR-29 in vivo improves glycemic control in adult mice. *Physiol Genomics*. 2019;51(8):379-389. [\[CrossRef\]](#)
23. Zampetaki A, Kiechl S, Drozdov I, et al. Plasma microRNA profiling reveals loss of endothelial MiR-126 and other microRNAs in type 2 diabetes. *Circ Res*. 2010;107(6):810-817. [\[CrossRef\]](#)
24. Massart J, Sjögren RJO, Lundell LS, et al. Altered miR-29 expression in type 2 diabetes influences glucose and lipid metabolism in skeletal muscle. *Diabetes*. 2017;66(7):1807-1818. [\[CrossRef\]](#)
25. Wang S, Zhang M, Liang B, Americans for Medical Progress. K $\alpha$ 2 Deletion causes aberrant expression and activation of NAD(P)H oxidase and consequent endothelial dysfunction in vivo: role of 26S proteasomes. *Circ Res*. 2010;106:1117-1128. [\[CrossRef\]](#)
26. Salama A, Fichou N, Allard M, et al. MicroRNA-29b modulates innate and antigen-specific immune responses in mouse models of autoimmunity. *PLoS One*. 2014;9(9):e106153. [\[CrossRef\]](#)
27. Ventriglia G, Nigi L, Sebastiani G, Dotta F. MicroRNAs: novel players in the dialogue between pancreatic islets and immune system in autoimmune diabetes. *BioMed Res Int*. 2015;2015:749734. [\[CrossRef\]](#)
28. Kwon DN, Chang BS, Kim JH. MicroRNA dysregulation in liver and pancreas of CMP-Neu5Ac hydroxylase null mice disrupts insulin/PI3K-AKT signaling. *BioMed Res Int*. 2014;2014:236385. [\[CrossRef\]](#)
29. Liu J, Ye C, Liu W, et al. AICAR enhances insulin signaling via downregulation of miR-29. *Can J Physiol Pharmacol*. 2016;94(2):199-205. [\[CrossRef\]](#)
30. van Dam EM, Govers R, James DE. Akt activation is required at a late stage of insulin-induced GLUT4 translocation to the plasma membrane. *Mol Endocrinol*. 2005;19(4):1067-1077. [\[CrossRef\]](#)
31. Kampmann U, Christensen B, Nielsen TS, et al. GLUT4 and UBC9 protein expression is reduced in muscle from type 2 diabetic patients with severe insulin resistance. *PLoS One*. 2011;6(11):e27854. [\[CrossRef\]](#)
32. Yonamine CY, Pinheiro-Machado E, Michalani ML, et al. Resveratrol improves glycemic control in type 2 diabetic obese mice by regulating glucose transporter expression in skeletal muscle and liver. *Molecules*. 2017;22(7):1-13. [\[CrossRef\]](#)
33. Esteves JV, Yonamine CY, Pinto-Junior DC, Gerlinger-Romero F, Enguita FJ, Machado UF. Diabetes modulates microRNAs 29b-3p, 29c-3p, 199a-5p and 532-3p expression in muscle. Possible role in GLUT4 and HK2 repression. *Front Endocrinol*. 2018;9:536. [\[CrossRef\]](#)
34. Chavey C, Boucher J, Monthouël-Kartmann MN, et al. Regulation of secreted protein acidic and rich in cysteine during adipose conversion and adipose tissue hyperplasia. *Obesity (Silver Spring)*. 2006;14(11):1890-1897. [\[CrossRef\]](#)
35. Kos K, Wilding JPH. SPARC: A key player in the pathologies associated with obesity and diabetes. *Nat Rev Endocrinol*. 2010;6(4):225-235. [\[CrossRef\]](#)
36. Nie J, Bradshaw AD, Delany AM, Sage EH. Inactivation of SPARC enhances high-fat diet-induced obesity in mice. *Connect Tissue Res*. 2011;52(2):99-108. [\[CrossRef\]](#)
37. Song H, Ding L, Zhang S, Wang W. MiR-29 family members interact with SPARC to regulate glucose metabolism. *Biochem Biophys Res Commun*. 2018;497(2):667-674. [\[CrossRef\]](#)
38. Liang J, Liu C, Qiao A, et al. MicroRNA-29a-c decrease fasting blood glucose levels by negatively regulating hepatic gluconeogenesis. *J Hepatol*. 2013;58(3):535-542. [\[CrossRef\]](#)
39. Huan T, Chen G, Liu C, et al. Age-associated microRNA expression in human peripheral blood is associated with all-cause mortality and age-related traits. *Aging Cell*. 2018;17(1):1-10. [\[CrossRef\]](#)
40. Fenn AM, Smith KM, Lovett-Racke AE, Guerau-de-Arellano M, Whitacre CC, Godbout JP. Increased micro-RNA 29b in the aged brain correlates with the reduction of insulin-like growth factor-1 and fractalkine ligand. *Neurobiol Aging*. 2013;34(12):2748-2758. [\[CrossRef\]](#)

# Turnitin Mirna29 Scopus

## ORIGINALITY REPORT

**23%**  
SIMILARITY INDEX

**13%**  
INTERNET SOURCES

**21%**  
PUBLICATIONS

**3%**  
STUDENT PAPERS

## PRIMARY SOURCES

1	Hui Wang, Tian-Tian Wang, Xiu-Ping Lv. "Expression and prognostic value of miRNA-29b in peripheral blood for endometrial cancer", Future Oncology, 2018 Publication	1%
2	<a href="http://bjp.sagepub.unboundmedicine.com">bjp.sagepub.unboundmedicine.com</a> Internet Source	1%
3	<a href="http://journals.sagepub.com">journals.sagepub.com</a> Internet Source	1%
4	Submitted to Drexel University Student Paper	<1%
5	<a href="http://cora.ucc.ie">cora.ucc.ie</a> Internet Source	<1%
6	<a href="http://scholar.ui.ac.id">scholar.ui.ac.id</a> Internet Source	<1%
7	Ning Li, Juanlian Cui, Xuanchu Duan, Huihui Chen, Fang Fan. "Suppression of Type I Collagen Expression by miR-29b via PI3K, Akt, and Sp1 Pathway in Human Tenon's	<1%

# Fibroblasts", Investigative Ophthalmology & Visual Science, 2012

Publication

8

[www.pacjmedsci.com](http://www.pacjmedsci.com)

Internet Source

<1 %

9

BRUNO GROLLEMUND, Caroline Dissaux, Pascale Gavelle, Carla Perez Martinez, Jimmy Mullaert, Toni Alfaiate, Antoine Guedeney. "The relational development of infants with CLP: impact on parents and on the parent-infant relationship. Results from the first French prospective multicenter CLP study.", Research Square Platform LLC, 2020

Publication

<1 %

10

Jichao Liang, Changzheng Liu, Aijun Qiao, Ying Cui et al. "MicroRNA-29a-c decrease fasting blood glucose levels by negatively regulating hepatic gluconeogenesis", Journal of Hepatology, 2013

Publication

<1 %

11

Meyliana, Erick Fernando, Surjandy, Marjuky. "The Business Process of Good Manufacturing Practice Based on Blockchain Technology in the Pharmaceutical Industry", 2021 Fifth International Conference on Information Retrieval and Knowledge Management (CAMP), 2021

Publication

<1 %



12	<a href="http://bio-bigdata.hrbmu.edu.cn">bio-bigdata.hrbmu.edu.cn</a> Internet Source	<1 %
13	<a href="http://ses.library.usyd.edu.au">ses.library.usyd.edu.au</a> Internet Source	<1 %
14	Julian Friedrich, Guido Krenning. "MicroRNAs linking oxidative stress and diabetes", Elsevier BV, 2020 Publication	<1 %
15	Rakhmetkazhy Bersimbaev, Akmaral Aripova, Olga Bulgakova, Assya Kussainova, Almira Akparova, Alberto Izzotti. "The Plasma Levels of hsa-miR-19b-3p, hsa-miR-125b-5p, and hsamiR- 320c in Patients with Asthma, COPD and Asthma-COPD Overlap Syndrome (ACOS)", MicroRNA, 2021 Publication	<1 %
16	Xinli Huang, Yun Gao, Jianjie Qin, Sen Lu. "The Role of miR-34a in the Hepatoprotective Effect of Hydrogen Sulfide on Ischemia/Reperfusion Injury in Young and Old Rats", PLoS ONE, 2014 Publication	<1 %
17	<a href="http://em3foamed.squarespace.com">em3foamed.squarespace.com</a> Internet Source	<1 %
18	Joanna Szydełko, Beata Matyjaszek-Matuszek. "MicroRNAs as Biomarkers for Coronary Artery Disease Related to Type 2 Diabetes Mellitus—From Pathogenesis to Potential	<1 %

# Clinical Application", International Journal of Molecular Sciences, 2022

Publication

---

19

Long Zhang, Charbel S. Hobeika, Damir Khabibullin, Deyang Yu et al. "Hypersensitivity to ferroptosis in chromophobe RCC is mediated by a glutathione metabolic dependency and cystine import via solute carrier family 7 member 11", Proceedings of the National Academy of Sciences, 2022

Publication

---

<1 %

20

Silvia Garavelli, Sara Bruzzaniti, Elena Tagliabue, Francesco Prattichizzo et al. "Blood Co-Circulating Extracellular microRNAs and Immune Cell Subsets Associate with Type 1 Diabetes Severity", International Journal of Molecular Sciences, 2020

Publication

---

<1 %

21

Submitted to University of Stellenbosch, South Africa

Student Paper

---

<1 %

22

[fjps.springeropen.com](https://fjps.springeropen.com)

Internet Source

---

<1 %

23

[vtechworks.lib.vt.edu](https://vtechworks.lib.vt.edu)

Internet Source

---

<1 %

24

E. van Rooij, L. B. Sutherland, J. E. Thatcher, J. M. DiMaio, R. H. Naseem, W. S. Marshall, J. A.

<1 %

Hill, E. N. Olson. "Dysregulation of microRNAs after myocardial infarction reveals a role of miR-29 in cardiac fibrosis", Proceedings of the National Academy of Sciences, 2008

Publication

---

25

Qianqian Zhang, Xiaofeng Yu, Linghan Tian, Yanjun Cong, Linfeng Li. "Therapeutic effects of epigallocatechin and epigallocatechin gallate on the allergic reaction of s1-casein sensitized mice", Food Science and Human Wellness, 2023

Publication

---

<1 %

26

Submitted to University of Western Sydney

Student Paper

---

<1 %

27

Xue-Qiang Zhu, Pei Lu, Zhong-Lin Xu, Qiang Zhou, Jun Zhang, Zhi-Biao Wang, Feng Wu. "Alterations in Immune Response Profile of Tumor-Draining Lymph Nodes after High-Intensity Focused Ultrasound Ablation of Breast Cancer Patients", Cells, 2021

Publication

---

<1 %

28

Jiayu Zhang, Liang Wu, Jiawei Chen, Sisi Lin, Daqiu Cai, Chengwei Chen, Zhenguo Chen. "Downregulation of MicroRNA 29a/b exacerbated diabetic retinopathy by impairing the function of Müller cells via Forkhead box protein O4", Diabetes and Vascular Disease Research, 2018

<1 %

29 [assets.researchsquare.com](https://assets.researchsquare.com) <1 %  
Internet Source

---

30 [mafiadoc.com](https://mafiadoc.com) <1 %  
Internet Source

---

31 "Handbook of Immunosenescence", Springer Science and Business Media LLC, 2019 <1 %  
Publication

---

32 Guohua Ding, Qian Yang, Xueyu Ruan, Tingting Si, Boxuan Yuan, Wenwei Zheng, Quanliang Xie, Ousmane Ahmat Souleymane, Xuchu Wang. "Proteomics analysis of the effects for different salt ions in leaves of true halophyte *Sesuvium portulacastrum*", *Plant Physiology and Biochemistry*, 2022 <1 %  
Publication

---

33 Onder Tulumbaci, M. Anil Onan, Sedat Turkoglu, Mertihan Kurdoglu, Bulent Boyaci, M. Bulent Tiras. "Effects of ritodrine hydrochloride tocolysis on echocardiographic parameters", *The Journal of Maternal-Fetal & Neonatal Medicine*, 2009 <1 %  
Publication

---

34 Rui Bi, Chunrong Bao, Lianyong Jiang, Hao Liu, Yang Yang, Ju Mei, Fangbao Ding. "MicroRNA-27b plays a role in pulmonary arterial hypertension by modulating peroxisome

proliferator-activated receptor  $\gamma$  dependent  
Hsp90-eNOS signaling and nitric oxide  
production", Biochemical and Biophysical  
Research Communications, 2015

Publication

---

35

Sukanya Srivastava, Richa Rathor, Som Nath Singh, Geetha Suryakumar. "Emerging role of MyomiRs as biomarkers and therapeutic targets in skeletal muscle diseases", American Journal of Physiology-Cell Physiology, 2021

Publication

---

36

Tang, X.. "Role of microRNAs in diabetes", BBA - Gene Regulatory Mechanisms, 200811

Publication

---

37

[mail.scialert.net](mailto:mail.scialert.net)

Internet Source

---

38

[theoncologist.alphamedpress.org](http://theoncologist.alphamedpress.org)

Internet Source

---

39

[veterinaryresearch.biomedcentral.com](http://veterinaryresearch.biomedcentral.com)

Internet Source

---

40

[www.karger.com](http://www.karger.com)

Internet Source

---

41

Edy Siswantoro, Nasrul Hadi Purwanto, Sutomo . "Effectiveness of Alkali Water Consumption to Reduce Blood Sugar Levels in Diabetes Mellitus Type 2", Journal of Diabetes Mellitus, 2017

<1 %

<1 %

<1 %

<1 %

<1 %

<1 %

<1 %

42

Katarzyna Szkudelska, Marzanna Deniziak, Maciej Sassek, Ignacy Szkudelski, Wojciech Noskowiak, Tomasz Szkudelski. "Resveratrol Affects Insulin Signaling in Type 2 Diabetic Goto-Kakizaki Rats", International Journal of Molecular Sciences, 2021

Publication

---

<1 %

43

Katarzyna Szkudelska, Tomasz Szkudelski. "The anti-diabetic potential of betaine. Mechanisms of action in rodent models of type 2 diabetes", Biomedicine & Pharmacotherapy, 2022

Publication

---

<1 %

44

Kayla R Steele, Michelle L Townsend, Brin F.S. Grenyer. "Parenting stress and competence in borderline personality disorder: The influence of mental health, trauma history, attachment and reflective capacity.", Research Square Platform LLC, 2020

Publication

---

<1 %

45

Marta Greco, Eusebio Chiefari, Maria Mirabelli, Alessandro Salatino et al. "Chapter 34-1 Insights into the World of MicroRNAs", Springer Science and Business Media LLC, 2022

Publication

---

<1 %

46

N E Priadi, H Anggraini, E T W Maharani. "The Differences of Salivary and Urine Acetone Levels in Diabetes Mellitus Patients", IOP Conference Series: Earth and Environmental Science, 2019

Publication

<1 %

47

Ruixia Zhu, Xu Liu, Zhiyi He. "Association of miR-122, miR-124 miR-126 and miR-143 gene polymorphisms with ischemic stroke in the northern Chinese Han population", International Journal of Neuroscience, 2019

Publication

<1 %

48

Yuan, Li, Chang Zhou, Yanxia Lu, Min Hong, Zuoyang Zhang, Zheyang Zhang, Yaya Chang, Chao Zhang, and Xuenong Li. "IFN- $\gamma$ -mediated IRF1/miR-29b feedback loop suppresses colorectal cancer cell growth and metastasis by repressing IGF1", Cancer Letters, 2015.

Publication

<1 %

49

[docksci.com](http://docksci.com)

Internet Source

<1 %

50

[eprints.gla.ac.uk](http://eprints.gla.ac.uk)

Internet Source

<1 %

51

[escholarship.org](http://escholarship.org)

Internet Source

<1 %

52

[gene-quantification.biz](http://gene-quantification.biz)

Internet Source

<1 %

53

[www.tandfonline.com](http://www.tandfonline.com)

Internet Source

&lt;1 %

54

Aishwarya P. Dasare, Piyush Gondaliya, Akshay Srivastava, Kiran Kalia. "A therapeutic approach towards microRNA29 family in vascular diabetic complications: A boon or curse?", *Journal of Diabetes & Metabolic Disorders*, 2019

Publication

&lt;1 %

55

Hyung-Keun Kim, Joo Dong Park, Seung Hee Choi, Dong Jun Shin, Sohyun Hwang, Hae-Yun Jung, Kyung-Soon Park. "Functional Link between miR-200a and ELK3 Regulates the Metastatic Nature of Breast Cancer", *Cancers*, 2020

Publication

&lt;1 %

56

Laura Nigi, Giuseppina Grieco, Giuliana Ventriglia, Noemi Brusco et al. "MicroRNAs as Regulators of Insulin Signaling: Research Updates and Potential Therapeutic Perspectives in Type 2 Diabetes", *International Journal of Molecular Sciences*, 2018

Publication

&lt;1 %

57

Lora-Pozo, Lucena-Anton, Salazar, Galán-Mercant, Moral-Munoz. "Anthropometric, Cardiopulmonary and Metabolic Benefits of the High-Intensity Interval Training Versus

&lt;1 %



Moderate, Low-Intensity or Control for Type 2 Diabetes: Systematic Review and Meta-Analysis", International Journal of Environmental Research and Public Health, 2019

Publication

---

58

Manlio Vinciguerra, Michelangelo Foti. "PTEN and SHIP2 phosphoinositide phosphatases as negative regulators of insulin signalling", Archives of Physiology and Biochemistry, 2008

Publication

---

<1 %

59

Pamungkas, Chamroonsawasdi, Vatanasomboon, Charupoonphol. "Barriers to Effective Diabetes Mellitus Self-Management (DMSM) Practice for Glycemic Uncontrolled Type 2 Diabetes Mellitus (T2DM): A Socio Cultural Context of Indonesian Communities in West Sulawesi", European Journal of Investigation in Health, Psychology and Education, 2019

Publication

---

<1 %

60

Paola Mirra, Cecilia Nigro, Immacolata Prevezano, Alessia Leone et al. "The Destiny of Glucose from a MicroRNA Perspective", Frontiers in Endocrinology, 2018

Publication

---

<1 %

61

Ran Tao, Xiang-Xue Fan, Hai-Jing Yu, Guo Ai, Hong-Yue Zhang, Hong-Yan Kong, Qi-Qin

<1 %

Song, Yu Huang, Jia-Quan Huang, Qin Ning. "MicroRNA-29b-3p prevents -induced liver fibrosis by targeting COL1A1 and COL3A1 ", Journal of Cellular Biochemistry, 2018

Publication

---

62

Xianwei Cui, Jingmin Tan, Yujie Shi, Chen Sun, Yun Li, Chenbo Ji, Jun Wu, Zhao Zhang, Siyu Chen, Xirong Guo, Chang Liu. "The long non-coding RNA Gm10768 activates hepatic gluconeogenesis by sequestering microRNA-214 in mice", Journal of Biological Chemistry, 2018

Publication

---

<1 %

63

Zhen He, Jian Jun Yang, Rong Zhang, Hua Ting Li, Liang Wu, Feng Jiang, Wei Ping Jia, Cheng Hu. "Circulating miR - 29b positively correlates with non - alcoholic fatty liver disease in a Chinese population", Journal of Digestive Diseases, 2019

Publication

---

<1 %

64

[encyclopedia.pub](http://encyclopedia.pub)

Internet Source

---

<1 %

65

[obgyn.pericles-prod.literatumonline.com](http://obgyn.pericles-prod.literatumonline.com)

Internet Source

---

<1 %

66

[researchspace.ukzn.ac.za](http://researchspace.ukzn.ac.za)

Internet Source

---

<1 %

67

[www.domo-systems.com](http://www.domo-systems.com)

Internet Source

<1 %

68

[www.pure.ed.ac.uk](http://www.pure.ed.ac.uk)

Internet Source

<1 %

69

Chandrasagar B Dugani, Amira Klip. "Glucose transporter 4: cycling, compartments and controversies", EMBO reports, 2005

Publication

<1 %

70

Chenggui Miao, Guoxue Zhang, Zhongwen Xie, Jun Chang. "MicroRNAs in the pathogenesis of type 2 diabetes: new research progress and future direction", Canadian Journal of Physiology and Pharmacology, 2018

Publication

<1 %

71

Francesco Prattichizzo, Angelica Giuliani, Artan Ceka, Maria Rita Rippo et al. "Epigenetic mechanisms of endothelial dysfunction in type 2 diabetes", Clinical Epigenetics, 2015

Publication

<1 %

72

Giuseppina Emanuela Grieco, Zein Mersini Besharat, Giada Licata, Daniela Fignani et al. "Circulating microRNAs as clinically useful biomarkers for Type 2 Diabetes Mellitus: miRNomics from bench to bedside", Translational Research, 2022

Publication

<1 %

73 Jin - Dian Tan, Mei - Feng Zhou, Song Yang, Jian - Ping Lin. " Long noncoding promotes osteosarcoma cell proliferation, invasion, and migration via the axis ", The Kaohsiung Journal of Medical Sciences, 2022  
Publication

---

74 Julie Massart, Rasmus J.O. Sjögren, Leonidas S. Lundell, Jonathan M. Mudry et al. "Altered miR-29 Expression in Type 2 Diabetes Influences Glucose and Lipid Metabolism in Skeletal Muscle", Diabetes, 2017  
Publication

---

75 MicroRNAs and Other Non-Coding RNAs in Inflammation, 2015.  
Publication

---

76 Mohammed Alfaifi, Amit Kumar Verma, Mohammed Yahya Alshahrani, Prakash C Joshi et al. "Cell-free long non-coding RNA-H19 and miRNA-29a, miRNA-29b expression in Type 2 Diabetes Mellitus patients.", Research Square, 2020  
Publication

---

77 [www.cell.com](http://www.cell.com)  
Internet Source

---

78 "Explaining Health Across the Sciences", Springer Science and Business Media LLC, 2020  
Publication

---

79

Mustafa Abdo Saif Dehwah, Aimin Xu,  
Qingyang Huang. "MicroRNAs and Type 2  
Diabetes/Obesity", Journal of Genetics and  
Genomics, 2012

Publication

---

<1 %

---

Exclude quotes Off

Exclude matches Off

Exclude bibliography On