



Search mail



Compose

5,974

Inbox

Starred

Snoozed

Important

Sent

Categories

More

Labels

Unwanted

270 of 9,513

Thank you for your submission to Journal of Sport Science and Fitness

Inbox x



Editorial Team of Journal of Sport Science and Fitness jssf@mail.unnes.a... Sat, Jun 15, 10:19 AM
to me

Dear Yuliana,

Thank you for your submission to Journal of Sport Science and Fitness. We have received your submission, EFFECTIVENESS OF WHEY PROTEIN AND SOYMILK ON MUSCLE MASS AND HEMATOCRIT LEVELS OF VOLLEYBALL ATHLETES, and a member of our editorial team will see it soon. You will be sent an email when an initial decision is made, and we may contact you for further information.

You can view your submission and track its progress through the editorial process at the following location:

Submission URL: <https://journal.unnes.ac.id/journals/jssf/authorDashboard/submission/7229>

If you have been logged out, you can login again with the username ulvieanna.

If you have any questions, please contact me from your [submission dashboard](#).

Thank you for considering Journal of Sport Science and Fitness as a venue for your work.



Search mail



Compose

104 of 9,513

Inbox

5,974

Your submission has been reviewed and we encourage you to submit revisions

Inbox x



Starred

Snoozed

Important

Sent

Categories

More

Labels

Unwanted



Fuadah Nor Wiqoyatul Milla Fuadahnw@gmail.com via my.unnes.id
to me

Sat, Jul 27, 10:14 AM

Dear Yuliana,

Your submission EFFECTIVENESS OF WHEY PROTEIN AND SOYMILK ON MUSCLE MASS AND HEMATOCRIT LEVELS OF VOLLEYBALL ATHLETES has been reviewed and we would like to encourage you to submit revisions that address the reviewers' comments. An editor will review these revisions and if they address the concerns adequately, your submission may be accepted for publication.

The reviewers' comments are included at the bottom of this email. Please respond to each point in the reviewers' comments and identify what changes you have made. If you find any of the reviewer's comments to be unjustified or inappropriate, please explain your perspective.

When you have completed your revisions, you can upload revised documents along with your response to the reviewers' comments at your [submission dashboard](#). If you have been logged out, you can login again with the username ulvieanna.

If you have any questions, please contact me from your [submission dashboard](#).

We look forward to receiving your revised submission.

Kind regards,

	<p>JSSF (vol) (issue) (year)</p> <p>Journal of Sport Sciences and Fitness</p> <p>http://journal.unnes.ac.id/sju/index.php/jssf</p>		
<p>EFFECTIVENESS OF WHEY PROTEIN AND SOYMILK ON MUSCLE MASS AND HEMATOCRIT LEVELS OF VOLLEYBALL ATHLETES</p>			
<p>Article's Info</p> <hr/> <p><i>Article's History:</i></p> <p>Received Month Year</p> <p>Accepted Month Year</p> <p>Published Month Year</p> <hr/> <p>Keywords: whey protein, soy milk, muscle mass, hematocrit levels, volleyball athlete</p>		<p>Abstract</p> <hr/> <p>Volleyball is one of the team sports that requires robust muscle mass for exceptional performance and endurance. Protein intake contributes to the formation of muscle mass and red blood cells that are tightly linked to hematocrit levels. The objective of this study was to determine the effectiveness of whey protein and soymilk on muscle mass and hematocrit levels in volleyball athletes. The study was a true experiment with a randomized pre-test and post-test design, involving 20 adolescent, male volleyball athletes divided into 2 groups: whey protein and the soymilk group selected by purposive sampling. The research variables were muscle mass and hematocrit levels. Statistical analysis used independent t-test and paired t-test. The results showed increased muscle mass in the whey protein group (3.50 ± 15.04), but the opposite was found in the soymilk (-1.82 ± 6.46), resulting in significant difference in muscle mass changes between groups ($p=0.020$). Both groups exhibited reduced hematocrit levels (whey protein group = -3.50 ± 7.69; soymilk group = -1.20 ± 6.30), with a significantly higher decrease shown in the whey protein group ($p=0.019$).</p>	
<p>Correspondence E-mail: *ulvieanna@gmail.com</p>			<p>ISSN 2252-6528</p>

Introduction

Volleyball is a very popular team sport in Indonesia because volleyball is a fun sport and can be a place to channel talent (Oktaviani, 2017). Volleyball athletes need good muscle strength to provide a high-quality play (Fauzan, 2019). Muscle tissue makes up 40-50% of total body weight. Adequate muscle mass is required in volleyball games to reduce risks of joint injury during training. The more muscle mass in the body, the better the muscle capacity is (Harahap, 2014). Optimal muscle mass is achieved from a combination of intense training and sufficient protein intake (Azhar, 2013).

Protein plays a major role in muscle building (Harahap, 2014). Protein contains essential branched-chain amino acids (BCAA)—leucine, isoleucine, and valine—which are metabolized in skeletal muscle. Setiowati (2013) reported that increased protein intake led to elevated muscle mass of 3.8%. Accordingly, Morton (2017) stated that increasing daily protein intake, along with regular exercise training, will cause changes in muscle mass. In addition, protein promotes blood cell formation (hemopoiesis), particularly erythrocytes (Andarina and Sumarni, 2006).

The volume of red blood cells in 100 ml of blood is expressed as hematocrit (Rahmatillah, 2019). Hematocrit reflects the ratio of red blood cells compared to total blood volume (Lita et al, 2016). Physical activity incites elevated lung capacity and overall blood circulation, implying increased number of circulating red blood cells which transports oxygen to cells and tissues (Norwidiyanti et al, 2022). Chai et al (2019) found a simple but significant correlation between $VO_2\max$ and hematocrit levels. $VO_2\max$ is the amount of oxygen available to exercising muscles to produce energy (Hariyanti et al, 2020). An athlete's $VO_2\max$ can be influenced by nutrient intake, particularly protein (Kameswara P.S, 2014).

Whey protein is the protein component in whey milk (Hutama, 2019). It has a high concentration of BCAA (26%) and has been proven to be beneficial for muscle building (Mardiana et al, 2022). However, the high product cost makes it unfit for athletes that would require regular consumption. As an alternative, soybeans may be a more affordable and accessible option.

Soybean (*Glycine max*) is known for its high protein content complemented with the complete essential amino acids and excellent digestibility. Lynch et al (2020) discovered that consuming soybeans positively affected muscle mass. Soy protein (*Glycine max*) contains isoflavones (sub-class of flavonoids) which function as antioxidants to prevent muscle damage (Fitriana et al, 2014). This study aims to determine the effectiveness of whey protein and soymilk on muscle mass gain and hematocrit levels in volleyball athletes.

METHOD

The study was a true experiment with a randomized pre-test post-test design. A total of 16 young, male volleyball athletes were selected by purposive sampling based on the following inclusion criteria: adolescents aged 14 to 18-year-old, have been an athlete for at least a year, not currently taking muscle mass enhancing supplements, and not participating in any programs to build muscle mass. Samples were equally divided into 2 groups and given a glass of either whey protein milk or soymilk after training for 12 days within 4 weeks. Muscle mass measurements were taken at the beginning and end of the study using a skinfold caliper while hematocrit levels were measured at the beginning and end of the study using For a Kit Test 6 Plus.

Primary data collected were training status, intake, physical activity, and nutritional status. Intake was assessed with 4 x 24-hour recall. Measurements of muscle mass and hematocrit levels were carried out on athletes before and after treatment. Univariate analysis was done to describe the mean, standard deviation, maximum and minimum value of the data. Normality test was done with Shapiro Wilk, then bivariate analysis used Independent Sample T-Test and mean difference was analyzed using Paired t-test.

RESULT AND DISCUSSION

Sample Characteristics

Sixteen male volleyball athletes were divided into two groups, whey protein and soy milk group, had characteristics as described in Table 1. The level of compliance in consuming whey protein and soy milk was 100%.

Commented [Ma1]: Explanation about role of protein in hematocrit will be better mentioned in the beginning of the paragraph

Commented [Ma2]: Explain about soymilk (1-2 sentences)

Commented [Ma3]: How many (in ml)

Table 1 Sample characteristics

Characteristic	Whey Protein	Soymilk	<i>p</i> value
BMI-for-age (kg/m ²)	0.73±0.45	0.63±0.62	0.693
Age (year)	16.50±1.17	16.30±1.49	0.744
Training status (year)	3.90±1.66	2.30±1.76	0.052
Physical activity level (PAL)	1.68±0.57	1.50±0.18	0.051
Energy intake (%)	102.07±1.50	91.25±1.06	0.082
Protein intake (%)	83.14±1.27	78.34±1.08	0.377
Fat intake (%)	85.80±1.78	75.00±1.71	0.185
Carbohydrate intake (%)	110.67±1.74	102.00±2.06	0.323
Calcium intake (%)	31.4±1.06	21.6±1.33	0.084
Vitamin C intake (%)	38.4±2.68	30.8±2.45	0.516
Iron intake (%)	78.6±1.17	70.21±1.84	0.244

No difference was found between groups for any of the sample characteristics listed in Table 1 ($p > 0.05$) so that BMI-for-age, age, exercise status, physical activity (PAL), adequacy level did not become confounding variables.

Body composition is the sum of all parts of the body consisting of adipose or fat tissue and lean tissue mass. Lean body mass includes muscle mass, bone, skin, non-fat body tissue and other body tissues, with muscle mass constituting to 40-50% of lean body mass (Browers & Fox, 1988). The formation of body fat mass in comparison to lean body mass is related to athlete performance.

Muscle Mass

Table 2. Muscle mass changes during the intervention

Muscle mass	Mean±SD			<i>p</i> ^a
	Pre (%)	Post (%)	Δ muscle mass	
Whey Protein	67.87±15.2	71.37±10.4	3.50±15.04	0.048
Sari Kedelai	66.82±9.52	65.00±8.59	-1.82±6.46	0.039
<i>p</i> ^b	0.856	0.022	0.020	

Notes :

p^a : paired t-test

p^b : independent t-test

Significant difference was found in muscle mass changes in both groups. Paired t-test analysis confirmed that whey protein group exhibited a notable increase ($p = 0.048$) of muscle mass at the end of the intervention, whereas the soymilk group experienced a significant reduction ($p = 0.039$). Therefore, it could be inferred that the two protein sources had opposite effects on muscle mass.

This study proved that whey protein is more effective for increasing muscle mass than soymilk, possibly due to the difference in BCAA amino acid contents. One gram of whey protein contains 116 mg leucine, 64.1 mg isoleucine and 60.9 mg valine; while BCAA content for every gram of soybean protein is 86.08 mg leucine, 54.4 mg isoleucine and 55.84 mg valine (Yanti et al, 2021). Furthermore, one serving of whey protein contains 11.22 grams of protein while one serving of soymilk contains 10.5 grams of protein. Setiowati and Hadi (2013) also affirmed that high protein intake was significantly correlated with muscle mass.

The post-intervention disparity of muscle mass changes among groups were most likely linked to BCAA which plays an important role in energy metabolism for muscle performance (Mardiana et al, 2022). The increased muscle mass was seemingly attributed to the high BCAA content in whey protein (26%). This type of amino acids is typically absorbed by the intestines to later produce NH₃ and other organic compounds that will eventually culminate in the urea cycle. However, during exercise, constant muscle damage occurs,

inciting immediate body response including the conversion of amino acids into new muscle cells (Harahap, 2014).

On the other hand, the decreased muscle mass in the soymilk group could be caused by the presence of antinutritional compounds in soybeans. Antinutrient compounds are secondary metabolites produced by plants as a form of defense against biological attacks (Aviles-Gaxiola et al, 2018). The metabolites produced are tannin, phytic acid, and antitrypsin, of which antitrypsin is the main factor causing limited protein utilization in soybeans (Khattab and Arnfield, 2009). Antitrypsin is the main antinutrient compound in soybeans that can inhibit the performance of the trypsin enzyme by forming an indigestible enzyme-inhibitor complex, thus interfering protein absorption. The antitrypsin content in soybeans reported the highest content of 94.1 U/mg (Khattab and Arnfield, 2009).

Hematocrit levels

Tabel 3. Hematocrit level changes during the intervention

Hematocrit	Mean±SD			<i>p</i> ^a
	Pre (%)	Post (%)	Δ hematokrit	
Whey Protein	39.30±4.62	35.80±9.17	-3.50±7.69	0.021
Soymilk	37.10±3.81	35.90±5.85	-1.20±6.30	0.040
<i>p</i> ^b	0.261	0.049	0.047	

Notes :

p^a : paired t-test

p^b : independent t-test

There was a notable difference in hematocrit levels across groups post-intervention. Paired t-test analysis revealed significant effects on hematocrit levels in both the whey protein ($p=0.021$) and soymilk group ($p=0.040$).

This study proves that soymilk can maintain hematocrit levels in athletes compared to whey protein because soymilk contains isoflavones that function as antioxidants. Antioxidants can play a role in preventing oxidative stress due to exposure to free radicals that affect hematocrit levels. In addition, soymilk protein plays a role in increasing hematocrit levels which affect $VO_2\max$. Similarly, Chai et al (2019) also found a simple but significant correlation between $VO_2\max$ and hematocrit. An athlete's $VO_2\max$ can be influenced by macronutrient intake such as protein (Kameswara, P.S, 2014).

Both groups experienced decreased hematocrit levels post-intervention, suggesting that neither whey protein nor soymilk were effective in increasing hematocrit levels. One of the factors related to this is "sports anemia" or anemia that occurs in athletes. Sports anemia is caused by increased plasma volume in the blood. Physical exercise with heavy intensity that is carried out continuously and with a long duration can also cause a decrease in red blood cell mass by intravascular hemolysis, causing hematocrit levels to decrease.

The protein in whey protein and soymilk affects iron absorption. However, whey protein and soymilk also contain calcium, which negatively impacts the absorption of heme and non-heme iron. Calcium inhibits iron transport through the basolateral membrane and enterocytes to the plasma. If iron absorption is inhibited, hematocrit levels will be low. The calcium content in one serving of whey protein is 171.6 mg while soy milk is 57.72 mg. Therefore, soy milk is more effective in maintaining hematocrit levels than soy milk (Roziq and Nuryanto, 2016).

CONCLUSION

Whey protein has better effects than soymilk in building and increasing muscle mass. Soymilk is better for maintaining hematocrit levels than whey protein.

REFERENCES

Andarina, D., Sumarmi S. 2006. Hubungan Konsumsi Protein Hewani dan Zat Besi dengan Kadar Hemoglobin pada Balita Usia 13 – 36 Bulan. *The Indonesian Journal of Public Health*. 3 (1): 19 – 23.

- Azhar, D.S., 2013. Perbandingan Pengaruh Susu Sapi Murni Dengan Susu Kedelai Terhadap Peningkatan Massa Otot Pada Latihan Beban. *Skripsi Universitas Pendidikan Indonesia*. Bandung.
- Chai, G., Qiu, J., Chen, S., Pan, Q., Shen, X., and Kang, J. 2019. Hematological, Hormonal and Fitness Indices in Youth Swimmers: Gender-Related Comparisons. *Journal of Human Kinetics*. 70 (2): 69 – 80.
- Davies, R.W., Carson, B.P., and Jakeman, P.M. 2018. The Effect of Whey Protein Supplementation on the Temporal Recovery of Muscle Function Following Resistance Training: A Systematic Review and Meta-Analysis. *Journal Nutrients*. 10 (2): 1 – 10.
- Fauzan, F.A. 2019. Pengaruh Pemberian Puding Kacang Merah (*Phaseolus vulgaris. L*) Terhadap Massa Otot dan Ketahanan Otot Pada Atlet Voli Remaja di Persatuan Bola Voli Binataruna Kota Semarang. *Skripsi Universitas Muhammadiyah Semarang*. Semarang.
- Fitriana, A., Rosidi, A., Pakpaham, T.R. 2014. Gambaran Asupan Vitamin Sebagai Zat Antioksidan Atlet Sepakbola di Pusat Pendidikan dan Latihan Pelajar Jawa Tengah di Salatiga. *Jurnal Gizi Universitas Muhammadiyah Semarang*. 3 (1): 16 – 21.
- Harahap, N.S. 2014. Protein dalam Nutrisi Olahraga. *Jurnal Ilmu Keolahragaan*. 13 (2): 45 – 54.
- Hariyanti, M.S., Rahayu, N.I., and Pitriani, P. 2020. Hubungan Kadar Hemoglobin dan Vo₂max Pada Atlet Softball Putra. *Jurnal Terapan Ilmu Keolahragaan*. 5 (1): 16 – 21.
- Hutama, R.F., and Andoyo, R. 2019. Produksi Whey Protein Concentrate (WPC) Terdenaturasi Menggunakan Beberapa Metode Preparasi. *Universitas Padjadjaran*. 3(1): 39 – 46.
- Kameswara, P.S.I. 2014. Perbedaan Nilai VO₂max dan Jarak Tempuh Lari Antara Pemberian Susu Rendah Lemak dan Minuman Olahraga Komersial Pada Atlet Sepak Bola. *Skripsi Universitas Diponegoro*. Semarang
- Kritikos, S., Papanikolaou, K., Draganidis, D., Poulos, A., Georgakouli, K., Tsimeas, P., Tzatzakis, T., Batsilas, D., Batrakoulis, A., Deli, C.K., Chatzinikolaou, A., Mohr, M., Jamurtas, A.Z., and Fatouros, L.G. 2021. Effect of Whey vs. Soy Protein Supplementation on Recovery Kinetics Following Speed Endurance Training in Competitive Male Soccer Players: a Randomized Controlled Trial. *Journal of the International Society of Sports Nutrition*. 18 (23): 1 – 15.
- Lita, E., Santosa, B., and Sukeksi, A. 2016. Perbedaan Kadar Hematokrit Berdasarkan Waktu Penundaan. *Karya Tulis Ilmiah Universitas Muhammadiyah Semarang*. Semarang.
- Liu, Y., Du, H., Li, P., Shen, Y., Peng, H., Liu, S., Zhou, G.A., Zhang, H., Liu, Z., Shi, M., Huang, X., Li, Y., Zhang, M., Wang, Z., Zhu, B., Han, B., Liang, C., and Tian, Z. 2020. Pan-Genome of Wild and Cultivated Soybeans. *Cell Resource*. 182: 162 – 176.
- Lynch, H.M., Buman, M.P., Dickinson, J.M., Ransdell, L.B., Johnston, C.S., and Wharton, C.M. 2020. No Significant Differences in Muscle Growth and Strength Development When Consuming Soy and Whey Protein Supplements Matched for Leucine Following a 12 week Resistance Training Program in Men and Women: A Randomized Trial. *International Journal of Environmental Research and Public Health*. 17 (3871): 1 – 14.
- Mardiana, Rachmawati, L., Sari, N.P., and Al Amien, T.N., 2022. Whey Protein, Daun Kelor, Kurma, dan Kelelahan Otot. *Book Chapter Kesehatan Masyarakat Jilid 2*. Universitas Negeri Semarang. Semarang.
- Messina, M., Lynch, H., Dickinson, J.M., and Reed, K.E. 2018. No Difference Between the Effects of Supplementing With Soy Protein Versus Animal Protein on Gains in Muscle Mass and Strength in Response to Resistance Exercise. *Internasional Journal of Sport Nutrition and Exercise Metabolism*. 28 (6): 674 – 685.

- Morton, R.W., Murphy, K.T., McKellar, S.R., Schoenfeld, B.J., Henselmans, M., Helms, E., Aragon, A.A., Devries, M.C., Banfield, L., Krieger, J.W., and Phillips, S.M. 2017. A Systematic review, meta-analysis and meta-regression of The Effect of Protein Supplementation on Resistance Training-Induced Gains in Muscle Mass and Strength in Healthy Adults. *Br J Sport Med.* 52: 376 – 384.
- Norwidianti, T., Kaidah, S., and Huldani. 2022. Literature Review: Hubungan Kadar Hematokrit dengan Nilai VO₂max Pada Atlet. *Jurnal Homeostatis.* 5 (1): 203 – 210.
- Oktaviany, D.D. 2017. Perbedaan Pengaruh Latihan *Rope Jump* dan *Squat Jump* Dengan Metode *Interval* Terhadap Daya Ledak Otot Tungkai Pemain Bola Voli. *Universitas 'Aisyiyah Yogyakarta.* Yogyakarta.
- Patil, G., Mian, R., Vuong, T., Pantalone, V., Song, Q., Chen, P., Shannon, G.J., Carter, T.C., and Nguyen, H.T. 2017. Molecular Mapping and Genomics of Soybean Seed Protein: a Review and Perspective For The Future. *Theor Appl Genect.* 130 (10): 1975 – 1991.
- Rahman, A., Kaidah, S., Huldani. 2021. Literature Review: Pengaruh Latihan Aerobik Intensitas Sedang Terhadap Kadar Hematokrit. *Jurnal Homeostatis.* 4 (2): 435 – 446.
- Rahmatillah, A.A. 2018. Pemeriksaan Hematokrit Pada Petani Garam di Dusun Ageng Desa Pinggir Papas Sumenep. *Skripsi Universitas Muhammadiyah Surabaya.* Surabaya.
- Rutherford, S.M., Fanning, A.C., Miller, B.J., and Moughan, P.J. 2015. Protein Digestibility-Corrected Amino Acid Scores and Digestible Indispensable Amino Acid Scores Differentially Describe Protein Quality in Growing Male Rats. *The Journal of Nutrition.* (145): 372 – 379.
- Siska, M.T., Zahtamal, and Putri, F. 2019. Pengaruh Kombinasi Latihan Beban dengan Metode *Pyramid set* dan Konsumsi Susu Tinggi Protein Terhadap Peningkatan Massa Otot. *Jurnal Ilmu Kesehatan.* 13 (2): 107 – 116.
- Setiowati, A., Hadi. 2013. Pengaruh Suplementasi Protein Terhadap Komposisi Tubuh pada Atlet. *Jurnal Media Ilmu Keolahragaan Indonesia.* 3 (2): 67 – 71.
- Tokede, O.A., Onabanjo, T.A., Yansane, A., Gaziano, J.M., and Djousse, L. 2015. Soya Products and Serum Lipids: a Meta-Analysis of Randomised Controlled Trials. *British Journal of Nutrition.* 144 (6): 831 – 843.
- Wim H.M., Saris and Marleen A. van Baak. 2018. *Glucagon and insulin responses after ingestion of different amounts of intact and hydrolysed proteins.* *British Journal of Nutrition.* 100, 61 – 69.
- Yanti, R., Angkasa, D., and Jus'at, I. 2021. Pengembangan Produk Snack Bar Tinggi BCAA (*Branched-chain Amino Acid*) Berbahan Tepung Kapri (*Pisum sativum*), Kecapir (*Psophocarpus tetragonolbus*) dan Kedelai (*Glycine max*) Sebagai Makanan Alternatif Untuk Daya Tahan Atlet. *The Journal of Nutrition and Food Research.* 44 (1): 21 – 30.

Author's Name / Journal of Sport Sciences and Fitness (vol) (issue) (year)



Search mail



Compose

107 of 9,513

Inbox

5,974

Your submission has been sent for another round of review

Inbox x



Starred



Fuadah Nor Wiqoyatul Milla Fuadahnw@gmail.com via my.unnes.id
to me

Fri, Jul 26, 5:03 PM

Snoozed

Important

Sent

Categories

More

Dear Yuliana,

Your revised submission, EFFECTIVENESS OF WHEY PROTEIN AND SOYMILK ON MUSCLE MASS AND HEMATOCRIT LEVELS OF VOLLEYBALL ATHLETES, has been sent for a new round of peer review. You will hear from us with feedback from the reviewers and information about the next steps.

If you have any questions, please contact me from your [submission dashboard](#).

Kind regards,

Fuadah Nor Wiqoyatul Milla

Labels

Unwanted

Reply
Forward
😊



Search mail



Compose

83 of 9,513

Inbox

5,974

Your submission has been accepted to Journal of Sport Science and Fitness



Starred

Snoozed

Important

Sent

Categories

More

Labels

Unwanted



Gustiana Mega Anggita mega.anggita@mail.unnes.ac.id via my.unnes.id to me

Wed, Jul 31, 1:47 PM

Dear Yuliana,

I am pleased to inform you that we have decided to accept your submission without further revision. After careful review, we found your submission, EFFECTIVENESS OF WHEY PROTEIN AND SOYMILK ON MUSCLE MASS AND HEMATOCRIT LEVELS OF VOLLEYBALL ATHLETES, to meet or exceed our expectations. We are excited to publish your piece in Journal of Sport Science and Fitness and we thank you for choosing our journal as a venue for your work.

Your submission is now forthcoming in a future issue of Journal of Sport Science and Fitness and you are welcome to include it in your list of publications. We recognize the hard work that goes into every successful submission and we want to congratulate you on reaching this stage.

Your submission will now undergo copy editing and formatting to prepare it for publication.

You will shortly receive further instructions.

If you have any questions, please contact me from your [submission dashboard](#).

Kind regards,

Gustiana Mega Anggita



Search mail



Compose

5,974

Just published: Vol. 10 No. 1 (2024): Journal of Sport Science and Fitness of Journal of Sport Science and Fitness

Inbox x



Gustiana Mega Anggita mega.anggita@mail.unnes.ac.id via my.unnes.id to me

Wed, Jul 31, 10:03 PM

Dear Yuliana,

We are pleased to announce the publication of [Vol. 10 No. 1 \(2024\): Journal of Sport Science and Fitness](#) of Journal of Sport Science and Fitness. We invite you to read and share this work with your scholarly community.

Many thanks to our authors, reviewers, and editors for their valuable contributions, and to our readers for your continued interest.

Thank you,

Gustiana Mega Anggita

[Unsubscribe](#) from emails sent by [Journal of Sport Science and Fitness](#).

Reply
Forward
😊

- Inbox
- Starred
- Snoozed
- Important
- Sent
- Categories
- More

- Labels
- Unwanted



EFFECTIVENESS OF WHEY PROTEIN AND SOYMILK ON MUSCLE MASS AND HEMATOCRIT LEVELS OF VOLLEYBALL ATHLETES

Yuliana Noor Setiawati Ulvie [✉], Aprodhit Justicia Nabila¹, Ali Rosidi¹, Zahra Labtrobdiba¹

¹Universitas Muhammadiyah Semarang

ulvieanna@gmail.com¹

Article's Info

Abstract

Article's History:

Received June 2024

Accepted July 2024

Published Juli 2024

Keywords:

whey protein, soy milk, muscle mass, hematocrit levels, volleyball athlete

Volleyball is one of the team sports that requires robust muscle mass for exceptional performance and endurance. Protein intake contributes to the formation of muscle mass and red blood cells that are tightly linked to hematocrit levels. The objective of this study was to determine the effectiveness of whey protein and soymilk on muscle mass and hematocrit levels in volleyball athletes. The study was a true experiment with a randomized pre-test and post-test design, involving 20 adolescent, male volleyball athletes divided into 2 groups: whey protein and the soymilk group selected by purposive sampling. The research variables were muscle mass and hematocrit levels. Statistical analysis used independent t-test and paired t-test. The results showed increased muscle mass in the whey protein group (3.50 ± 15.04), but the opposite was found in the soymilk (-1.82 ± 6.46), resulting in significant difference in muscle mass changes between groups ($p=0.020$). Both groups exhibited reduced hematocrit levels (whey protein group = -3.50 ± 7.69 ; soymilk group = -1.20 ± 6.30), with a significantly higher decrease shown in the whey protein group ($p=0.019$).

INTRODUCTION

Volleyball is a very popular team sport in Indonesia because volleyball is a fun sport and can be a place to channel talent (Oktaviany, 2017). Volleyball athletes need good muscle strength to provide a high-quality play (Fauzan, 2019). Muscle tissue makes up 40-50% of total body weight. Adequate muscle mass is required in volleyball games to reduce risks of joint injury during training. The more muscle mass in the body, the better the muscle capacity is (Harahap, 2014). Optimal muscle mass is achieved from a combination of intense training and sufficient protein intake (Azhar, 2013).

Protein plays a major role in muscle building (Harahap, 2014). Protein contains essential branched-chain amino acids (BCAA)—leucine, isoleucine, and valine—which are metabolized in skeletal muscle. Setiowati (2013) reported that increased protein intake led to elevated muscle mass of 3.8%. Accordingly, Morton (2017) stated that increasing daily protein intake, along with regular exercise training, will cause changes in muscle mass. In addition, protein promotes blood cell formation (hemopoiesis), particularly erythrocytes (Andarina and Sumarni, 2006).

An athlete's VO_2 max can be influenced by nutrient intake, particularly protein (Kameswara P.S, 2014). VO_2 max is the amount of oxygen available to exercising muscles to produce energy (Hariyanti et al, 2020). Chai et al (2019) found a simple but significant correlation between VO_2 max and hematocrit levels. The volume of red blood cells in 100 ml of blood is expressed as hematocrit (Rahmatillah, 2019). Hematocrit reflects the ratio of red blood cells compared to total blood volume (Lita et al, 2016). Physical activity incites elevated lung capacity and overall blood circulation,

implying increased number of circulating red blood cells which transports oxygen to cells and tissues (Norwidianti et al, 2022).

Whey protein is the protein component in whey milk (Hutama, 2019). It has a high concentration of BCAA (26%) and has been proven to be beneficial for muscle building (Mardiana et al, 2022). However, the high product cost makes it unfit for athletes that would require regular consumption. As an alternative, soybeans may be a more affordable and accessible option.

Soybean (*Glycine max*) is known for its high protein content complemented with the complete essential amino acids and excellent digestibility. Lynch et al (2020) discovered that consuming soybeans positively affected muscle mass. Soy protein (*Glycine max*) contains isoflavones (sub-class of flavonoids) which function as antioxidants to prevent muscle damage (Fitriana et al, 2014). One of the processed soybean products is soy milk. Soy milk according to SNI 01-3830-1995 is a product derived from soy bean seed extract with water or a solution of soy flour in water, with or without the addition of other permitted food ingredients (Adawiyah et al, 2018). This study aims to determine the effectiveness of whey protein and soymilk on muscle mass gain and hematocrit levels in volleyball athletes.

METHOD

The study was a true experiment with a randomized pre-test post-test design. A total of 16 young, male volleyball athletes were selected by purposive sampling based on the following inclusion criteria: adolescents aged 14 to 18-year-old, have been an athlete for at least a year, not currently taking muscle mass enhancing supplements, and not participating in any programs to build muscle mass. Samples were equally

divided into 2 groups and given a glass of (210 ml) either whey protein milk or soymilk after training for 12 days within 4 weeks. Muscle mass measurements were taken at the beginning and end of the study using a skinfold caliper while hematocrit levels were measured at the beginning and end of the study using For a Kit Test 6 Plus.

Primary data collected were training status, intake, physical activity, and nutritional status. Intake was assessed with 4 x 24-hour recall. Measurements of muscle mass and hematocrit levels were carried out on athletes before and after treatment. Univariate analysis was done to describe the mean, standard deviation, maximum and minimum value of the data. Normality test was done with Shapiro Wilk, then bivariate analysis used Independent Sample T-Test and mean difference was analyzed using Paired t-test.

RESULT AND DISCUSSION

The result of the research is presented in the form of chart, table, or description. The analysis and interpretation of the result is needed to be elaborated before being discussed. The discussion is focused of elaborating the data and result of the analysis with the problems of the research or the purpose of the research in a broader theoretical context. In the discussion, also, providing answer of why based on the facts taken from the data. The discussion is written according to the data that is being discussed and supposedly not biased from the data result.

Table 1 Sample characteristics

Characteristic	Whey Protein	Soymilk	<i>p</i>
BMI-for-age (kg/m ²)	0.73±0.45	0.63±0.62	0.693
Age (year)	16.50±1.17	16.30±1.49	0.744
Training status (year)	3.90±1.66	2.30±1.76	0.052
	1.68±0.57	1.50±0.18	0.051

	Mean±SD	Mean±SD	<i>p</i> ^a
Physical activity level (PAL)	102.07±1.50	91.25±1.06	0.082
Energy intake (%)	83.14±1.27	78.34±1.08	0.377
Protein intake (%)	85.80±1.78	75.00±1.71	0.185
Fat intake (%)	110.67±1.74	102.00±2.06	0.323
Carbohydrate intake (%)	31.4±1.06	21.6±1.33	0.084
Calcium intake (%)	38.4±2.68	30.8±2.45	0.516
Vitamin C intake (%)	78.6±1.17	70.21±1.84	0.244
Iron intake (%)			

Tabel 2. Muscle mass changes during the intervention

Muscle mass	Mean±SD			<i>p</i> ^a
	Pre (%)	Post (%)	Δ muscle mass	
Whey Protein	67.87±15.2	71.37±10.4	3.50±15.04	0.048
Sari Kedelai	66.82±9.52	65.00±8.59	-1.82±6.46	0.039
<i>p</i> ^b	0.856	0.022	0.020	

Notes :

p^a : paired t-test

p^b : independent t-test

Tabel 3. Hematocrit level changes during the intervention

Hematocrit	Mean±SD			<i>p</i> ^a
	Pre (%)	Post (%)	Δ hematokrit	
Whey Protein	39.30±4.62	35.80±9.17	-3.50±7.69	0.021
Soymilk	37.10±3.81	35.90±5.85	-1.20±6.30	0.040
<i>p</i> ^b	0.261	0.049	0.047	

Notes :

p^a : paired t-test

p^b : independent t-test

CONCLUSION

Whey protein has better effects than soymilk in building and increasing muscle mass. Soymilk is better for maintaining hematocrit levels than whey protein.

REFERENCES

- Adawiyah, D.R., Andarwulam, N., Triana, R.N., Agustin, D., Gitaprawati, D. 2018. Evaluation of Soybean Varieties for Quality of Soymilk. *Jurnal Mutu Pangan*. 5 (1): 10 – 16.
- Andarina, D., Sumarmi S. 2006. Hubungan Konsumsi Protein Hewani dan Zat Besi dengan Kadar Hemoglobin pada Balita Usia 13 – 36 Bulan. *The Indonesian Journal of Public Health*. 3 (1): 19 – 23.
- Azhar, D.S., 2013. Perbandingan Pengaruh Susu Sapi Murni Dengan Susu Kedelai Terhadap Peningkatan Massa Otot Pada Latihan Beban. *Skripsi Universitas Pendidikan Indonesia*. Bandung.
- Chai, G., Qiu, J., Chen, S., Pan, Q., Shen, X., and Kang, J. 2019. Hematological, Hormonal and Fitness Indices in Youth Swimmers: Gender-Related Comparisons. *Journal of Human Kinetics*. 70 (2): 69 – 80.
- Davies, R.W., Carson, B.P., and Jakeman, P.M. 2018. The Effect of Whey Protein Supplementation on the Temporal Recovery of Muscle Function Following Resistance Training: A Systematic Review and Meta-Analysis. *Journal Nutrients*. 10 (2): 1 – 10.
- Fauzan, F.A. 2019. Pengaruh Pemberian Puding Kacang Merah (*Phaseolus vulgaris*. L) Terhadap Massa Otot dan Ketahanan Otot Pada Atlet Voli Remaja di Persatuan Bola Voli Binaruna Kota Semarang. *Skripsi Universitas Muhammadiyah Semarang*. Semarang.
- Fitriana, A., Rosidi, A., Pakpaham, T.R. 2014. Gambaran Asupan Vitamin Sebagai Zat Antioksidan Atlet Sepakbola di Pusat Pendidikan dan Latihan Pelajar Jawa Tengah di Salatiga. *Jurnal Gizi Universitas Muhammadiyah Semarang*. 3 (1): 16 – 21.
- Harahap, N.S. 2014. Protein dalam Nutrisi Olahraga. *Jurnal Ilmu Keolahragaan*. 13 (2): 45 – 54.
- Hariyanti, M.S., Rahayu, N.I., and Pitriani, P. 2020. Hubungan Kadar Hemoglobin dan Vo2max Pada Atlet Softball Putra. *Jurnal Terapan Ilmu Keolahragaan*. 5 (1): 16 – 21.
- Hutama, R.F., and Andoyo, R. 2019. Produksi Whey Protein Concentrate (WPC) Terdenaturasi Menggunakan Beberapa Metode Preparasi. *Universitas Padjadjaran*. 3(1): 39 – 46.
- Kameswara, P.S.I. 2014. Perbedaan Nilai VO₂max dan Jarak Tempuh Lari Antara Pemberian Susu Rendah Lemak dan Minuman Olahraga Komersial Pada Atlet Sepak Bola. *Skripsi Universitas Diponegoro*. Semarang
- Kritikos, S., Papanikolaou, K., Draganidis, D., Poulos, A., Georgakouli, K., Tsimeas, P., Tzatzakis, T., Batsilas, D., Batrakoulis, A., Deli, C.K., Chatzinikolaou, A., Mohr, M., Jamurtas, A.Z., and Fatouros, L.G. 2021. Effect of Whey vs. Soy Protein Supplementation on Recovery Kinetics Following Speed Endurance Training in Competitive Male Soccer Players: a Randomized Controlled Trial. *Journal of the International Society of Sports Nutrition*. 18 (23): 1 – 15.
- Lita, E., Santosa, B., and Sukeksi, A. 2016. Perbedaan Kadar Hematokrit Berdasarkan Waktu Penundaan. *Karya Tulis Ilmiah Universitas Muhammadiyah Semarang*. Semarang.
- Liu, Y., Du, H., Li, P., Shen, Y., Peng, H., Liu, S., Zhou, G.A., Zhang, H., Liu, Z., Shi, M., Huang, X., Li, Y., Zhang, M., Wang, Z., Zhu, B., Han, B., Liang, C., and Tian, Z. 2020. Pan-Genome of Wild and Cultivated Soybeans. *Cell Resource*. 182: 162 – 176.
- Lynch, H.M., Buman, M.P., Dickinson, J.M., Ransdell, L.B., Johnston, C.S., and Wharton, C.M. 2020. No Significant Differences in Muscle Growth and Strength Development When Consuming Soy and Whey Protein Supplements Matched for Leucine Following a 12 week Resistance Training Program in Men and Women: A Randomized Trial. *International Journal of Environmental Research and Public Health*. 17 (3871): 1 – 14.
- Mardiana, Rachmawati, L., Sari, N.P., and Al Amien, T.N., 2022. Whey Protein, Daun Kelor, Kurma, dan Kelelahan Otot. *Book Chapter Kesehatan Masyarakat Jilid 2*. Universitas Negeri Semarang. Semarang.
- Messina, M., Lynch, H., Dickinson, J.M., and Reed, K.E. 2018. No Difference Between the Effects of Supplementing With Soy Protein Versus Animal Protein on Gains in Muscle Mass and Strength in Response to Resistance Exercise. *Internasional Journal of Sport Nutrition and Exercise Metabolism*. 28 (6): 674 – 685.
- Morton, R.W., Murphy, K.T., McKellar, S.R., Schoenfeld, B.J., Henselmans, M., Helms, E., Aragon, A.A., Devries, M.C., Banfield, L., Krieger, J.W., and Phillips, S.M. 2017. A Systematic review, meta-


- analysis and meta-regression of The Effect of Protein Supplementation on Resistance Training-Induced Gains in Muscle Mass and Strength in Healthy Adults. *Br J Sport Med.* 52: 376 – 384.
- Norwidianti, T., Kaidah, S., and Huldani. 2022. Literature Review: Hubungan Kadar Hematokrit dengan Nilai VO_{2max} Pada Atlet. *Jurnal Homeostatis.* 5 (1): 203 – 210.
- Oktaviany, D.D. 2017. Perbedaan Pengaruh Latihan *Rope Jump* dan *Squat Jump* Dengan Metode *Interval* Terhadap Daya Ledak Otot Tungkai Pemain Bola Voli. *Universitas 'Aisyiyah Yogyakarta.* Yogyakarta.
- Patil, G., Mian, R., Vuong, T., Pantalone, V., Song, Q., Chen, P., Shannon, G.J., Carter, T.C., and Nguyen, H.T. 2017. Molecular Mapping and Genomics of Soybean Seed Protein: a Review and Perspective For The Future. *Theor Appl Genect.* 130 (10): 1975 – 1991.
- Rahman, A., Kaidah, S., Huldani. 2021. Literature Review: Pengaruh Latihan Aerobik Intensitas Sedang Terhadap Kadar Hematokrit. *Jurnal Homeostatis.* 4 (2): 435 – 446.
- Rahmatillah, A.A. 2018. Pemeriksaan Hematokrit Pada Petani Garam di Dusun Ageng Desa Pinggir Papas Sumenep. *Skripsi Universitas Muhammadiyah Surabaya.* Surabaya.
- Rutherford, S.M., Fanning, A.C., Miller, B.J., and Moughan, P.J. 2015. Protein Digestibility-Corrected Amino Acid Scores and Digestible Indispensable Amino Acid Scores Differentially Describe Protein Quality in Growing Male Rats. *The Journal of Nutrition.* (145): 372 – 379.
- Siska, M.T., Zahtamal, and Putri, F. 2019. Pengaruh Kombinasi Latihan Beban dengan Metode *Pyramid set* dan Konsumsi Susu Tinggi Protein Terhadap Peningkatan Massa Otot. *Jurnal Ilmu Kesehatan.* 13 (2): 107 – 116.
- Setiowati, A., Hadi. 2013. Pengaruh Suplementasi Protein Terhadap Komposisi Tubuh pada Atlet. *Jurnal Media Ilmu Keolahragaan Indonesia.* 3 (2): 67 – 71.
- Tokede, O.A., Onabanjo, T.A., Yansane, A., Gaziano, J.M., and Djousse, L. 2015. Soya Products and Serum Lipids: a Meta-Analysis of Randomised Controlled Trials. *British Journal of Nutrition.* 144 (6): 831 – 843.
- Wim H.M., Saris and Marleen A. van Baak. 2018. *Glucagon and insulin responses after ingestion of different amounts of intact and hydrolysed proteins.* *British Journal of Nutrition.* 100, 61 – 69.
- Yanti, R., Angkasa, D., and Jus'at, I. 2021. Pengembangan Produk Snack Bar Tinggi BCAA (*Branched-chain Amino Acid*) Berbahan Tepung Kapri (*Pisum sativum*), Kecipir (*Psophocarpus tetragonolbus*) dan Kedelai (*Glycine max*) Sebagai Makanan Alternatif Untuk Daya Tahan Atlet. *The Journal of Nutrition and Food Research.* 44 (1): 21 – 30

D3 Gizi

EFFECTIVENESS OF WHEY PROTEIN AND SOYMILK ON MUSCLE MASS AND HEMATOCRIT LEVELS OF VOLLEYBALL A...

 Jurnal 3

 D3 Gizi

 Universitas Muhammadiyah Semarang

Document Details

Submission ID

trn:oid::1:2986452619

Submission Date

Aug 20, 2024, 8:41 AM GMT+7

Download Date

Aug 20, 2024, 8:49 AM GMT+7

File Name

Copyedit_Yuliana_20082024.doc

File Size

141.5 KB

6 Pages

2,861 Words

16,609 Characters

12% Overall Similarity

The combined total of all matches, including overlapping sources, for each database.

Filtered from the Report

- ▶ Bibliography

Match Groups

- 15 Not Cited or Quoted 11%**
Matches with neither in-text citation nor quotation marks
- 0 Missing Quotations 0%**
Matches that are still very similar to source material
- 0 Missing Citation 0%**
Matches that have quotation marks, but no in-text citation
- 0 Cited and Quoted 0%**
Matches with in-text citation present, but no quotation marks

Top Sources

- 5% Internet sources
- 2% Publications
- 11% Submitted works (Student Papers)

Integrity Flags





0 Integrity Flags for Review

No suspicious text manipulations found.




Our system's algorithms look deeply at a document for any inconsistencies that would set it apart from a normal submission. If we notice something strange, we flag it for you to review.

A Flag is not necessarily an indicator of a problem. However, we'd recommend you focus your attention there for further review.

Match Groups

-  **15 Not Cited or Quoted 11%**
Matches with neither in-text citation nor quotation marks
-  **0 Missing Quotations 0%**
Matches that are still very similar to source material
-  **0 Missing Citation 0%**
Matches that have quotation marks, but no in-text citation
-  **0 Cited and Quoted 0%**
Matches with in-text citation present, but no quotation marks

Top Sources

- 5%  Internet sources
- 2%  Publications
- 11%  Submitted works (Student Papers)

Top Sources

The sources with the highest number of matches within the submission. Overlapping sources will not be displayed.

1	Student papers	
	Universitas Muhammadiyah Semarang	8%
2	Internet	
	journal.unnes.ac.id	3%
3	Internet	
	garuda.kemdikbud.go.id	2%



JSSF (9) (2) (2023)

Journal of Sport Sciences and Fitness<http://journal.unnes.ac.id/sju/index.php/jssf>**EFFECTIVENESS OF WHEY PROTEIN AND SOYMILK ON MUSCLE MASS AND HEMATOCRIT LEVELS OF VOLLEYBALL ATHLETES****Yuliana Noor Setiawati Ulvie^{1✉}, Aprodhit Justicia Nabila², Ali Rosidi³, Zahra Labtrobdiba⁴**Universitas Muhammadiyah Semarang^{1,2,3,4}ulvieanna@gmail.com¹aprodhitjusticia@gmail.com²alirrosidi@unimus.ac.id³zahra_latrobdiba@unimus.ac.id⁴**Article's Info***Article's History:*

Received June 2024

Accepted July 2024

Published July 2024

*Keywords:**whey protein, soy milk, muscle mass, hematocrit levels, volleyball athlete***Abstract**

Volleyball is one of the team sports that requires robust muscle mass for exceptional performance and endurance. Protein intake contributes to the formation of muscle mass and red blood cells that are tightly linked to hematocrit levels. The objective of this study was to determine the effectiveness of whey protein and soymilk on muscle mass and hematocrit levels in volleyball athletes. The study was a true experiment with a randomized pre-test and post-test design, involving 20 adolescent, male volleyball athletes divided into 2 groups: whey protein and the soymilk group selected by purposive sampling. The research variables were muscle mass and hematocrit levels. Statistical analysis used independent t-test and paired t-test. The results showed increased muscle mass in the whey protein group (3.50 ± 15.04), but the opposite was found in the soymilk (-1.82 ± 6.46), resulting in significant difference in muscle mass changes between groups ($p=0.020$). Both groups exhibited reduced hematocrit levels (whey protein group = -3.50 ± 7.69 ; soymilk group = -1.20 ± 6.30), with a significantly higher decrease shown in the whey protein group ($p=0.019$).

Correspondence E-mail✉: ulvieanna@gmail.com

Jl. Kedungmundu No.18, Kedungmundu, Kec. Tembalang, Kota Semarang, Jawa Tengah 50273

ISSN 2252-6528

INTRODUCTION

Volleyball is a very popular team sport in Indonesia because volleyball is a fun sport and can be a place to channel talent (Oktaviany, 2017). Volleyball athletes need good muscle strength to provide a high-quality play (Fauzan, 2019). Muscle tissue makes up 40-50% of total body weight. Adequate muscle mass is required in volleyball games to reduce risks of joint injury during training. The more muscle mass in the body, the better the muscle capacity is (Harahap, 2014). Optimal muscle mass is achieved from a combination of intense training and sufficient protein intake (Azhar, 2013).

Protein plays a major role in muscle building (Harahap, 2014). Protein contains essential branched-chain amino acids (BCAA)—leucine, isoleucine, and valine—which are metabolized in skeletal muscle. Setiowati (2013) reported that increased protein intake led to elevated muscle mass of 3.8%. Accordingly, Morton (2017) stated that increasing daily protein intake, along with regular exercise training, will cause changes in muscle mass. In addition, protein promotes blood cell formation (hemopoiesis), particularly erythrocytes (Andarina and Sumarni, 2006).

An athlete's VO_2 max can be influenced by nutrient intake, particularly protein (Kameswara P.S, 2014). VO_2 max is the amount of oxygen available to exercising muscles to produce energy (Hariyanti et al, 2020). Chai et al (2019) found a simple but significant correlation between VO_2 max and hematocrit levels. The volume of red blood cells in 100 ml of blood is expressed as hematocrit (Rahmatillah, 2019). Hematocrit reflects the ratio of red blood cells compared to total blood volume (Lita et al, 2016). Physical activity

incites elevated lung capacity and overall blood circulation, implying increased number of circulating red blood cells which transports oxygen to cells and tissues (Norwidianti et al, 2022).

Whey protein is the protein component in whey milk (Hutama, 2019). It has a high concentration of BCAA (26%) and has been proven to be beneficial for muscle building (Mardiana et al, 2022). However, the high product cost makes it unfit for athletes that would require regular consumption. As an alternative, soybeans may be a more affordable and accessible option.

Soybean (*Glycine max*) is known for its high protein content complemented with the complete essential amino acids and excellent digestibility. Lynch et al (2020) discovered that consuming soybeans positively affected muscle mass. Soy protein (*Glycine max*) contains isoflavones (sub-class of flavonoids) which function as antioxidants to prevent muscle damage (Fitriana et al, 2014). One of the processed soybean products is soy milk. Soy milk according to SNI 01-3830-1995 is a product derived from soy bean seed extract with water or a solution of soy flour in water, with or without the addition of other permitted food ingredients (Adawiyah et al, 2018). This study aims to determine the effectiveness of whey protein and soymilk on muscle mass gain and hematocrit levels in volleyball athletes.

METHOD

The study was a true experiment with a randomized pre-test post-test design. A total of 16 young, male volleyball athletes were selected by purposive sampling based on the following inclusion criteria: adolescents aged 14 to 18-

year-old, have been an athlete for at least a year, not currently taking muscle mass enhancing supplements, and not participating in any programs to build muscle mass. Samples were equally divided into 2 groups and given a glass of (210 ml) either whey protein milk or soymilk after training for 12 days within 4 weeks. Muscle mass measurements were taken at the beginning and end of the study using a skinfold caliper while hematocrit levels were measured at the beginning and end of the study using For a Kit Test 6 Plus.

Primary data collected were training status, intake, physical activity, and nutritional status. Intake was assessed with 4 x 24-hour recall. Measurements of muscle mass and hematocrit levels were carried out on athletes before and after treatment. Univariate analysis was done to describe the mean, standard deviation, maximum and minimum value of the data. Normality test was done with Shapiro Wilk, then bivariate analysis used Independent Sample T-Test and mean difference was analyzed using Paired t-test.

RESULT AND DISCUSSION

Sample Characteristics

Table 1 Sample characteristics

Characteristic	Whey Protein	Soymilk	<i>p</i>
BMI-for-age (kg/m ²)	0.73±0.45	0.63±0.62	0.693
Age (year)	16.50±1.17	16.30±1.49	0.744
Training status (year)	3.90±1.66	2.30±1.76	0.052
Physical activity level (PAL)	1.68±0.57	1.50±0.18	0.051
Energy intake (%)	102.07±1.50	91.25±1.06	0.082
Protein intake (%)	83.14±1.27	78.34±1.08	0.377
Fat intake (%)	85.80±1.78	75.00±1.71	0.185

Carbohydrate intake (%)	110.67±1.74	102.00±2.06	0.323
Calcium intake (%)	31.4±1.06	21.6±1.33	0.084
Vitamin C intake (%)	38.4±2.68	30.8±2.45	0.516
Iron intake (%)	78.6±1.17	70.21±1.84	0.244

Sixteen male volleyball athletes were divided into two groups, whey protein and soy milk group, had characteristics as described in Table 1. The level of compliance in consuming whey protein and soy milk was 100%.

No difference was found between groups for any of the sample characteristics listed in Table 1 (*p*>0.05) so that BMI-for-age, age, exercise status, physical activity (PAL), adequacy level did not become confounding variables.

Body composition is the sum of all parts of the body consisting of adipose or fat tissue and lean tissue mass. Lean body mass includes muscle mass, bone, skin, non-fat body tissue and other body tissues, with muscle mass constituting to 40-50% of lean body mass (Browers & Fox, 1988). The formation of body fat mass in comparison to lean body mass is related to athlete performance.

Muscle Mass

Table 2. Muscle mass changes during the intervention

Muscle mass	Mean±SD			<i>p^a</i>
	Pre (%)	Post (%)	Δ muscle mass	
Whey Protein	67.87±15.2	71.37±10.4	3.50±15.04	0.048
Sari Kedelai	66.82±9.52	65.00±8.59	-1.82±6.46	0.039
<i>p^b</i>	0.856	0.022	0.020	

Notes :
p^a : paired t-test
p^b : independent t-test

Significant difference was found in muscle mass changes in both groups. Paired t-

test analysis confirmed that whey protein group exhibited a notable increase ($p = 0.048$) of muscle mass at the end of the intervention, whereas the soymilk group experienced a significant reduction ($p = 0.039$). Therefore, it could be inferred that the two protein sources had opposite effects on muscle mass.

This study proved that whey protein is more effective for increasing muscle mass than soymilk, possibly due to the difference in BCAA amino acid contents. One gram of whey protein contains 116 mg leucine, 64.1 mg isoleucine and 60.9 mg valine; while BCAA content for every gram of soybean protein is 86.08 mg leucine, 54.4 mg isoleucine and 55.84 mg valine (Yanti et al, 2021). Furthermore, one serving of whey protein contains 11.22 grams of protein while one serving of soymilk contains 10.5 grams of protein. Setiowati and Hadi (2013) also affirmed that high protein intake was significantly correlated with muscle mass.

The post-intervention disparity of muscle mass changes among groups were most likely linked to BCAA which plays an important role in energy metabolism for muscle performance (Mardiana et al, 2022). The increased muscle mass was seemingly attributed to the high BCAA content in whey protein (26%). This type of amino acids is typically absorbed by the intestines to later produce NH_3 and other organic compounds that will eventually culminate in the urea cycle. However, during exercise, constant muscle damage occurs, inciting immediate body response including the conversion of amino acids into new muscle cells (Harahap, 2014).

On the other hand, the decreased muscle mass in the soymilk group could be caused by the presence of antinutritional compounds in

soybeans. Antinutrient compounds are secondary metabolites produced by plants as a form of defense against biological attacks (Aviles-Gaxiola et al, 2018). The metabolites produced are tannin, phytic acid, and antitrypsin, of which antitrypsin is the main factor causing limited protein utilization in soybeans (Khattab and Arnfield, 2009). Antitrypsin is the main antinutrient compound in soybeans that can inhibit the performance of the trypsin enzyme by forming an indigestible enzyme-inhibitor complex, thus interfering protein absorption. The antitrypsin content in soybeans reported the highest content of 94.1 U/mg (Khattab and Arnfield, 2009).

Hematocrit levels

Tabel 3. Hematocrit level changes during the intervention

Hematocrit	Mean±SD			p^a
	Pre (%)	Post (%)	Δ hematokrit	
Whey Protein	39.30±4.62	35.80±9.17	-3.50±7.69	0.021
Soymilk	37.10±3.81	35.90±5.85	-1.20±6.30	0.040
p^b	0.261	0.049	0.047	

Notes :
 p^a : paired t-test
 p^b : independent t-test

There was a notable difference in hematocrit levels across groups post-intervention. Paired t-test analysis revealed significant effects on hematocrit levels in both the whey protein ($p=0.021$) and soymilk group ($p=0.040$).

This study proves that soymilk can maintain hematocrit levels in athletes compared to whey protein because soymilk contains isoflavones that function as antioxidants. Antioxidants can play a role in preventing oxidative stress due to exposure to free radicals

that affect hematocrit levels. In addition, soymilk protein plays a role in increasing hematocrit levels which affect $VO_2\max$. Similarly, Chai et al (2019) also found a simple but significant correlation between $VO_2\max$ and hematocrit. An athlete's $VO_2\max$ can be influenced by macronutrient intake such as protein (Kameswara, P.S, 2014).

Both groups experienced decreased hematocrit levels post-intervention, suggesting that neither whey protein nor soymilk were effective in increasing hematocrit levels. One of the factors related to this is "sports anemia" or anemia that occurs in athletes. Sports anemia is caused by increased plasma volume in the blood. Physical exercise with heavy intensity that is carried out continuously and with a long duration can also cause a decrease in red blood cell mass by intravascular hemolysis, causing hematocrit levels to decrease.

The protein in whey protein and soymilk affects iron absorption. However, whey protein and soymilk also contain calcium, which negatively impacts the absorption of heme and non-heme iron. Calcium inhibits iron transport through the basolateral membrane and enterocytes to the plasma. If iron absorption is inhibited, hematocrit levels will be low. The calcium content in one serving of whey protein is 171.6 mg while soy milk is 57.72 mg. Therefore, soy milk is more effective in maintaining hematocrit levels than soy milk (Roziqo and Nuryanto, 2016).

CONCLUSION

Whey protein has better effects than soymilk in building and increasing muscle mass. Soymilk is better for maintaining hematocrit levels than whey protein.

REFERENCES

- Adawiyah, D.R., Andarwulam, N., Triana, R.N., Agustin, D., Gitapratwi, D. 2018. Evaluation of Soybean Varieties for Quality of Soymilk. *Jurnal Mutu Pangan*. 5 (1): 10 – 16.
- Andarina, D., Sumarmi S. 2006. Hubungan Konsumsi Protein Hewani dan Zat Besi dengan Kadar Hemoglobin pada Balita Usia 13 – 36 Bulan. *The Indonesian Journal of Public Health*. 3 (1): 19 – 23.
- Azhar, D.S., 2013. Perbandingan Pengaruh Susu Sapi Murni Dengan Susu Kedelai Terhadap Peningkatan Massa Otot Pada Latihan Beban. *Skripsi Universitas Pendidikan Indonesia*. Bandung.
- Chai, G., Qiu, J., Chen, S., Pan, Q., Shen, X., and Kang, J. 2019. Hematological, Hormonal and Fitness Indices in Youth Swimmers: Gender-Related Comparisons. *Journal of Human Kinetics*. 70 (2): 69 – 80.
- Davies, R.W., Carson, B.P., and Jakeman, P.M. 2018. The Effect of Whey Protein Supplementation on the Temporal Recovery of Muscle Function Following Resistance Training: A Systematic Review and Meta-Analysis. *Journal Nutrients*. 10 (2): 1 – 10.
- Fauzan, F.A. 2019. Pengaruh Pemberian Puding Kacang Merah (*Phaseolus vulgaris. L*) Terhadap Massa Otot dan Ketahanan Otot Pada Atlet Voli Remaja di Persatuan Bola Voli Binataruna Kota Semarang. *Skripsi Universitas Muhammadiyah Semarang*. Semarang.
- Fitriana, A., Rosidi, A., Pakpaham, T.R. 2014. Gambaran Asupan Vitamin Sebagai Zat Antioksidan Atlet Sepakbola di Pusat Pendidikan dan Latihan Pelajar Jawa Tengah di Salatiga. *Jurnal Gizi Universitas Muhammadiyah Semarang*. 3 (1): 16 – 21.
- Harahap, N.S. 2014. Protein dalam Nutrisi Olahraga. *Jurnal Ilmu Keolahragaan*. 13 (2): 45 – 54.
- Hariyanti, M.S., Rahayu, N.I., and Pitriani, P. 2020. Hubungan Kadar Hemoglobin dan $Vo_2\max$ Pada Atlet Softball Putra. *Jurnal Terapan Ilmu Keolahragaan*. 5 (1): 16 – 21.
- Hutama, R.F., and Andoyo, R. 2019. Produksi Whey Protein Concentrate (WPC) Terdenaturasi Menggunakan Beberapa Metode Preparasi. *Universitas Padjadjaran*. 3(1): 39 – 46.
- Kameswara, P.S.I. 2014. Perbedaan Nilai $VO_2\max$ dan Jarak Tempuh Lari Antara Pemberian Susu Rendah Lemak dan Minuman Olahraga Komersial Pada Atlet Sepak Bola. *Skripsi Universitas Diponegoro*. Semarang

- Kritikos, S., Papanikolaou, K., Draganidis, D., Poulos, A., Georgakouli, K., Tsimeas, P., Tzatzakis, T., Batsilas, D., Batrakoulis, A., Deli, C.K., Chatzinikolaou, A., Mohr, M., Jamurtas, A.Z., and Fatouros, L.G. 2021. Effect of Whey vs. Soy Protein Supplementation on Recovery Kinetics Following Speed Endurance Training in Competitive Male Soccer Players: a Randomized Controlled Trial. *Journal of the International Society of Sports Nutrition*. 18 (23): 1 – 15.
- Lita, E., Santosa, B., and Sukeksi, A. 2016. Perbedaan Kadar Hematokrit Berdasarkan Waktu Penundaan. *Karya Tulis Ilmiah Universitas Muhammadiyah Semarang*. Semarang.
- Liu, Y., Du, H., Li, P., Shen, Y., Peng, H., Liu, S., Zhou, G.A., Zhang, H., Liu, Z., Shi, M., Huang, X., Li, Y., Zhang, M., Wang, Z., Zhu, B., Han, B., Liang, C., and Tian, Z. 2020. Pan-Genome of Wild and Cultivated Soybeans. *Cell Resource*. 182: 162 – 176.
- Lynch, H.M., Buman, M.P., Dickinson, J.M., Ransdell, L.B., Johnston, C.S., and Wharton, C.M. 2020. No Significant Differences in Muscle Growth and Strength Development When Consuming Soy and Whey Protein Supplements Matched for Leucine Following a 12 week Resistance Training Program in Men and Women: A Randomized Trial. *International Journal of Environmental Research and Public Health*. 17 (3871): 1 – 14.
- Mardiana, Rachmawati, L., Sari, N.P., and Al Amien, T.N., 2022. Whey Protein, Daun Kelor, Kurma, dan Kelelahan Otot. *Book Chapter Kesehatan Masyarakat Jilid 2*. Universitas Negeri Semarang. Semarang.
- Messina, M., Lynch, H., Dickinson, J.M., and Reed, K.E. 2018. No Difference Between the Effects of Supplementing With Soy Protein Versus Animal Protein on Gains in Muscle Mass and Strength in Response to Resistance Exercise. *International Journal of Sport Nutrition and Exercise Metabolism*. 28 (6): 674 – 685.
- Morton, R.W., Murphy, K.T., McKellar, S.R., Schoenfeld, B.J., Henselmans, M., Helms, E., Aragon, A.A., Devries, M.C., Banfield, L., Krieger, J.W., and Phillips, S.M. 2017. A Systematic review, meta-analysis and meta-regression of The Effect of Protein Supplementation on Resistance Training-Induced Gains in Muscle Mass and Strength in Healthy Adults. *Br J Sport Med*. 52: 376 – 384.
- Norwidianti, T., Kaidah, S., and Huldani. 2022. Literature Review: Hubungan Kadar Hematokrit dengan Nilai VO_{2max} Pada Atlet. *Jurnal Homeostatis*. 5 (1): 203 – 210.
- Oktaviani, D.D. 2017. Perbedaan Pengaruh Latihan *Rope Jump* dan *Squat Jump* Dengan Metode *Interval* Terhadap Daya Ledak Otot Tungkai Pemain Bola Voli. *Universitas 'Aisyiyah Yogyakarta*. Yogyakarta.
- Patil, G., Mian, R., Vuong, T., Pantalone, V., Song, Q., Chen, P., Shannon, G.J., Carter, T.C., and Nguyen, H.T. 2017. Molecular Mapping and Genomics of Soybean Seed Protein: a Review and Perspective For The Future. *Theor Appl Genet*. 130 (10): 1975 – 1991.
- Rahman, A., Kaidah, S., Huldani. 2021. Literature Review: Pengaruh Latihan Aerobik Intensitas Sedang Terhadap Kadar Hematokrit. *Jurnal Homeostatis*. 4 (2): 435 – 446.
- Rahmatillah, A.A. 2018. Pemeriksaan Hematokrit Pada Petani Garam di Dusun Ageng Desa Pinggir Papas Sumenep. *Skripsi Universitas Muhammadiyah Surabaya*. Surabaya.
- Rutherford, S.M., Fanning, A.C., Miller, B.J., and Moughan, P.J. 2015. Protein Digestibility-Corrected Amino Acid Scores and Digestible Indispensable Amino Acid Scores Differentially Describe Protein Quality in Growing Male Rats. *The Journal of Nutrition*. (145): 372 – 379.
- Siska, M.T., Zahtamal, and Putri, F. 2019. Pengaruh Kombinasi Latihan Beban dengan Metode *Pyramid set* dan Konsumsi Susu Tinggi Protein Terhadap Peningkatan Massa Otot. *Jurnal Ilmu Kesehatan*. 13 (2): 107 – 116.
- Setiowati, A., Hadi. 2013. Pengaruh Suplementasi Protein Terhadap Komposisi Tubuh pada Atlet. *Jurnal Media Ilmu Keolahragaan Indonesia*. 3 (2): 67 – 71.
- Tokede, O.A., Onabanjo, T.A., Yansane, A., Gaziano, J.M., and Djousse, L. 2015. Soy Products and Serum Lipids: a Meta-Analysis of Randomised Controlled Trials. *British Journal of Nutrition*. 144 (6): 831 – 843.
- Wim H.M., Saris and Marleen A. van Baak. 2018. *Glucagon and insulin responses after ingestion of different amounts of intact and hydrolysed proteins*. *British Journal of Nutrition*. 100, 61 – 69.
- Yanti, R., Angkasa, D., and Jus'at, I. 2021. Pengembangan Produk Snack Bar Tinggi BCAA (*Branched-chain Amino Acid*) Berbahan Tepung Kapri (*Pisum sativum*), Kecapir (*Psophocarpus tetragonolbus*) dan Kedelai (*Glycine max*) Sebagai Makanan Alternatif Untuk Daya Tahan Atlet. *The Journal of Nutrition and Food Research*. 44 (1): 21 – 30